

2016-2441  
2016-05-20

**KILMOS PF  
MOSQUITO  
COILS**

**DOMESTIC**

**Each Coil Burns for 6 Hours**

**GUARANTEE:**

d-cis trans Allethrin ..... 0.25 %

Warning, contains the allergen wheat.

REG. NO. 22671 P.C.P. ACT

**KEEP OUT OF REACH OF CHILDREN READ THE LABEL BEFORE USING**

Net Contents: Package containing 10 coils with 2 metal holders - Each Coil weighs approximately 12 grams.

Manufactured by: Blood Protection Co., Ltd.  
110-112 Austin Road, Kowloon,  
Hong Kong for;

J Mudryj Regulatory Consultants Ltd.  
17 OAKFIELD DRIVE  
ST. ANDREWS, MB  
R1A 2W9

ph: (204) 615-2197

**DIRECTIONS FOR USE:** Use stand as a holder for the coil. Pull the center piece to an upright position. Loosen outer rings of coil until ends move freely. Grasp each "comma-shaped" portion in the center between thumb and forefinger of each hand. Separate the comma-shaped portions by gently pressing them in opposite directions 5 millimeters apart. Hold one comma-shaped end and shake lightly or gently pry apart to separate coils. Insert tip of coil stand into slot in comma-shaped end of coil. Light outer end of coil until glowing. Blow out any flame and allow to smolder. Place coils 2-3 m (6-10 ft) apart around area to be protected. Use outdoors.

For best results, do not use in wind. Areas such as backyards, patios, decks, etc. which are protected or sheltered from wind are more easily protected by coils. In light winds, place coils upwind where possible to make use of associated drift. If shorter periods of use are desired, break coil into appropriate lengths and place in notched cut-out of holder. One coil will burn for approximately 6 hours.

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN. Cover exposed food. Avoid prolonged inhalation of smoke. Do not touch lit end. Do not place on flammable surfaces. After using, wash hands with soap and water.

**FIRST AID:** If contact with coil on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If ingested, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. Take container, label or product name and PCP Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**STORAGE AND DISPOSAL:** Store in a cool, dry place in original container. Do not reuse empty container. Wrap empty container and place in garbage.