

**LATER'S  
FOLPET ROSE & GARDEN FUNGICIDE**

WETTABLE POWDER

For use on Roses and listed Flowers, Vegetables and Fruits to control a number of diseases caused by fungus.

Causes Eye Irritation  
May be harmful if swallowed in inhaled

DOMESTIC

REG. NO. 15798 P.C.P. ACT

GUARANTEE: Folpet 50%

WARNING: Directions for use and cautionary statements should be carefully followed

**READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

80 g

Sure-Gro IP Inc.,  
P.O. Box 21001,  
Brantford, ON N3R 7W9  
1-800-268-2806

## **Later's Folpet Rose & Garden Fungicide**

**Use only on crops listed on this label**

**DIRECTIONS:** Later's Folpet Rose & Garden Fungicide is a wettable powder. When mixed with water it can be applied up to a day before harvest on listed fruits and vegetables except celery. Also very effective for fungus control on listed ornamentals and flowers.

1 level Tablespoon = 7 g

**HOW TO USE:** Mix 7 g of Later's Folpet with 10 L of water to make a spray solution and apply up to the day before harvest, unless otherwise indicated (1 tablespoon / 2 gals. of water).

**ASTERS, CARNATIONS, CHRYSANTHEMUMS, IRIS, MARIGOLDS, ROSES, SNAPDRAGONS:** To control Leaf Spot and Powdery Mildew, apply lightly to both leaf surfaces at weekly intervals. When heavy infestation occurs, spray twice a week.

To control Stem Rot on **AZALEA** cuttings, soak the cuttings for 15-30 minutes before planting.

**APPLES:** To control Alternaria Leaf Spot, Black Rot and Scab, apply every 7-10 days or as needed. To avoid injury for crops in Western Canada, do not use between bloom and July. Do not apply within 1 day of harvest. In Eastern Canada, treatments during the critical period (just before bloom until 30 days after petal fall) may cause russetting to sensitive varieties such as Golden Delicious, Red Delicious and Stayman Winesap.

**CELERY:** To control Early and Late Blight, apply after the plants are established in the field. Repeat at 7 day intervals. Do not apply within 7 days of harvest.

**CUCUMBERS, PUMPKINS, SQUASH and WATERMELONS:** To control Anthracnose and Powdery Mildew, make the first application when first leaves appear. Repeat at 7 day intervals until harvest. Do not apply within 1 day of harvest.

**GRAPES:** To control Blackrot and Mildew, apply just before and after bloom, and make additional applications at 2-3 week intervals and as needed. For Deadarm, apply when the new shoots are 2.5 – 5 cm long and repeat when growth is 10 - 15 cm long. Do not apply within 1 day of harvest.

**TOMATOES:** To control Anthracnose, mix 14 g of Later's Folpet with 10 L of water and begin application during first bloom and repeat at weekly intervals until harvest. (1 tablespoon / 1 gal. of water). Do not apply within 1 day of harvest.

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN. Avoid inhalation of dust or spray mist. May irritate eyes, nose, throat and skin. Avoid contact with skin, eyes and clothing. Do not contaminate feed or food. Wash hands carefully after use and before eating, drinking or smoking. May cause skin irritation or sensitization; avoid repeated skin contact. Wash contaminated clothing with soap and hot water before reuse. Keep away from heat, fire and sparks.

**ENVIRONMENTAL HAZARDS:** Do not contaminate ponds, lakes and streams.

**FIRST AID:** **If swallowed**, call a poison control centre or doctor IMMEDIATELY for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If in eyes**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If inhaled**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. **If on skin or clothing**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**STORAGE:** Keep product in original container. Store in a cool, dry place away from food.

**DISPOSAL:** Do not re-use empty container. Wrap container and discard in household garbage.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.