

€ ÆÖÖËËFFÄÛ àÄP| ÆËËJËH H

3270051567

FRONT PANEL

**Hartz® UltraGuard™
Flea & Tick Powder for Dogs**

Kills:

Fleas

Ticks

For 7 days

GUARANTEE:

Tetrachlorvinphos..... 3.3%

DOMESTIC

READ ENTIRE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

FOR USE ONLY ON DOGS

REG. NO.: 16673 PCP ACT

Net Contents: g

HARTZ CANADA INC.,
1125 Talbot Street
St. Thomas, Ontario
N5P 3W7
1-800-275-1414

BACK/SIDE PANEL

DIRECTIONS FOR USE :

Dust entire animal beginning at head and working back. Use no more than 2 grams of powder per kilogram of your pet's body weight. Avoid getting powder in pet's eyes or genital area. Make sure that the powder gets down to the skin. Take care to treat feet and legs. Dust pet's bedding and living quarters. Repeat at weekly intervals if necessary. Apply to bedding used by pets only. Do not apply to flooring, carpets, furniture, or other areas that may come into contact with humans. Use ONLY in well-ventilated areas, preferably outdoors.

Do NOT use on pets under 12 weeks of age. Consult a veterinarian before using on sick, aged, pregnant, or nursing animals or animals receiving drug or other pesticide treatment. Do not use this product on pets at the same time or within 30 days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides, or chemicals. Discontinue use if adverse effects are observed.

For effective flea and tick control, treatment of the pet should be combined with the cleanup and sanitation of any area used by the pet. Vacuum floors, carpet, furniture, etc. (discard vacuum bag after use) and wash the pets bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal treatment of the premises may be required. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

PRECAUTIONS:

HUMANS: KEEP OUT OF REACH OF CHILDREN. Avoid contact with mouth, eyes and clothing. Avoid breathing dust. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils or places where food is prepared or stored. **PETS:** Dust may irritate eyes, nose and throat. Harmful if swallowed or inhaled. Avoid contact with the dog's mouth, face, eyes and genitals. Avoid letting the pet inhale the dust.

Toxic to fish and other aquatic life. Cover fish aquaria in the vicinity of treatment area. Toxic to birds.

FIRST AID

HUMANS: IF SWALLOWED, Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **IF INHALED,** Move person to fresh air. If person is not

breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. **PETS:** Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label, or product name and P.C.P. Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: This product contains a pesticide that is a cholinesterase inhibitor (anti-cholinesterase compound). Symptoms of human poisoning may include headache, weakness, sweating, blurred vision, nausea and diarrhea. Obtain medical attention or call a poison control centre at once. Atropine is antidotal. Acute symptoms of over dosages in dogs include diarrhea, salivation, vomiting, muscular tremors and weakness. Contact a veterinarian immediately.

STORAGE AND DISPOSAL: Store in original, unopened container in a cool, dry place away from food and out of the reach of children. Do not reuse empty container. Wrap and dispose of empty container with household garbage.

For more information call our experts at 1-800-275-1414
Visit www.hartzultraguard.com

Hartz® and other trademarks are trademarks of The Hartz Mountain Corporation

RMXXXXXX

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.