

2012-03-14

2011-5357

BACKER CARD

WILSON[®]
SUPER MOUSE TREAT
with Bromadiolone

Impregnated Canary Seed for Killing Mice

Kills Mice

DOMESTIC

REG. NO. 16741 P.C.P. ACT

GUARANTEE: Bromadiolone 0.005%

WARNING



POISON

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

2 x 50 g

Sure-Gro IP Inc.,
P.O. Box 21001,
Brantford, ON
N3R 7W9
1-800-268-2806

DIRECTIONS: For House mouse (*Mus musculus*), punch out hole on face of box. Place bait in tamper-resistant bait boxes or protected areas inaccessible at the time of placement to children, pets, domestic animals and wildlife. Place 50 g boxes at intervals of 3 - 4 m . Replenish bait supply for at least 15 days or as long as there is evidence of feeding.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. Keep out of lakes, streams or ponds.

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

FIRST AID: This product contains an anticoagulant chemical which, if taken accidentally by humans, domestic animals, pets or wildlife, may reduce the clotting ability of the blood and haemorrhage may result. **If in eyes,** hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **If swallowed,** call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁, in the form of intramuscular or subcutaneous injections, or by oral ingestion, is a suggested remedial treatment for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Wrap container and discard in household garbage. Do not reuse empty container. Dispose of dead rodents in garbage or by burying.

For additional product information or in case of emergency, spills or fire, call toll-free 1-800-268-2806.

www.wilsonproducts.ca

Wilson® Registered Trademark of Spectrum Brands Canada, Inc., used under license.

CARTON

WILSON[®]
SUPER MOUSE TREAT
with Bromadiolone

Impregnated Canary Seed for Killing Mice

Kills Mice

DOMESTIC

REG. NO. 16741 P.C.P. ACT

GUARANTEE: BROMADIOLONE 0.005%

WARNING



POISON

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

50 g

Sure-Gro IP Inc.,
P.O. Box 21001,
Brantford, ON
N3R 7W9

1-800-268-2806

DIRECTIONS: For House mouse (*Mus musculus*), punch out hole on face of box. Place bait in tamper-resistant bait boxes or protected areas in accessible at the time of placement to children, pets, domestic animals and wildlife. Place 50 g boxes at intervals of 3 - 4 m . Replenish bait supply for at least 15 days or as long as there is evidence of feeding.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. Keep out of lakes, streams or ponds.

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

FIRST AID: This product contains an anticoagulant chemical which, if taken accidentally by humans, domestic animals, pets or wildlife, may reduce the clotting ability of the blood and haemorrhage may result. If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁, in the form of intramuscular or subcutaneous injections, or by oral ingestion is a suggested remedial treatment for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Wrap container and discard in household garbage. Do not reuse empty container. Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.