

2016-02-15  
2015-6668

BOTTLE

**LIQUID ROZOL RODENTICIDE**

FARM & RANCH BRAND

Concentrated Formula

**FOR CONTROL OF GROUND SQUIRRELS, NORTHERN POCKET GOPHERS,  
RATS AND MICE.**

AGRICULTURAL

This product is not to be used in and around homes or other residential areas, such as parks, school ground and playing fields.

GUARANTEE: Chlorophacinone 0.07%

REGISTRATION NO. 21160 PEST CONTROL PRODUCTS ACT

CAUTION



POISON

**KEEP OUT OF REACH OF CHILDREN  
READ THE LABEL AND LEAFLET BEFORE USING**

Only to be used by certified pest control operators, farmers and persons authorized in government-approved pest control programs.

NET CONTENTS: 250 mL

Premier Tech Brighton Ltd.  
1 avenue Premier  
Riviere-du-Loup QC G5C 6C1

**FIRST AID:** **If in eyes,** hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **If swallowed,** call a poison control centre or doctor immediately for treatment advice. Do not give **any** liquid to the person. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Vitamin K<sub>1</sub> in the form of intramuscular or subcutaneous injections or by oral ingestion is the suggested remedial treatment for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.** May be harmful or fatal if swallowed or absorbed through the skin. Chemical-resistant gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** If swallowed, this material may reduce clotting ability of the blood and cause bleeding. Keep out of lakes, streams or ponds.

**STORAGE:** Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

**DISPOSAL:** Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial regulations for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills. **Rodent Disposal:** Dispose of dead rodents in garbage or by burying.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

LEAFLET

**LIQUID ROZOL RODENTICIDE**

FARM & RANCH BRAND

Concentrated Formula

**FOR CONTROL OF GROUND SQUIRRELS, NORTHERN POCKET GOPHERS,  
RATS AND MICE.**

AGRICULTURAL

This product is not to be used in and around homes or other residential areas, such as parks, school ground and playing fields.

REGISTRATION NO. 21160 PEST CONTROL PRODUCTS ACT

GUARANTEE: Chlorophacinone 0.07%

CAUTION



POISON

**READ THE LABEL AND LEAFLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

Only to be used by certified pest control operators, farmers and persons authorized in government-approved pest control programs.

Premier Tech Brighton Ltd.  
1 avenue Premier  
Riviere-du-Loup QC G5C 6C1

### **USE LIMITATIONS – IN AND AROUND BUILDINGS:**

For indoor use only. DO NOT use in homes or other residential settings. DO NOT use in areas accessible to children. Bait MUST be placed either in tamper-resistant bait stations or in locations not accessible to pets, livestock, or non-target wildlife.

DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Bait stations (tier 2) used for the placement of rodenticide bait indoors, in locations accessible to pets, livestock or non-target wildlife must have the following characteristics: (1) be constructed of high-strength material (e.g., metal or injection moulded plastic) and resistant to destruction by non-target animals; (2) have an entrance designed so that non-target animals cannot reach the bait; (3) have an internal structure that prevents bait from being shaken loose; (4) have an access panel that fastens securely and locks (e.g., metal screw or padlock); and (5) bear the product name, active ingredient, guarantee, registration number, “WARNING POISON”, and the skull and crossbones symbol.

### **USE LIMITATIONS – GARBAGE DUMPS:**

DO NOT use in areas of the garbage dump that are or will be open to the public. DO NOT use in areas accessible to children. Bait placed outdoors, above-ground in locations accessible to pets or livestock MUST be placed in tier 1 bait stations. Tier 1 bait stations must have the following characteristics: (1) be resistant to destruction or weakening by elements of typical non-catastrophic weather (such as snow, rain, extremes of temperature and humidity, direct sunshine, etc.); (2) be constructed of high-strength material (e.g., metal or injection moulded plastic) and resistant to destruction by children and non-target animals; (3) have an entrance designed so that children and non-target animals cannot reach the bait; (4) have an internal structure that prevents bait from being shaken loose; (5) have an access panel that fastens securely and locks (e.g., metal screw or padlock); and (6) bear the product name, active ingredient, guarantee, registration number, “WARNING POISON”, and the skull and crossbones symbol.

### **USE LIMITATIONS – OTHER OUTDOOR AREAS:**

DO NOT use in residential or public areas (such as parks, playing fields), or any other sites that are or will be open to the public (for example, Pick-Your-Own sites). DO NOT use in areas accessible to children. Bait placed outdoors, above-ground in locations accessible to pets or livestock MUST be placed in tier 1 bait stations. The bait stations used in these sites MUST meet the characteristics described under **USE LIMITATIONS – GARBAGE DUMPS**.

**DIRECTIONS FOR USE:** Users should remove clothing immediately if pesticide gets inside. Then wash skin thoroughly and put on clean clothing.

**MIXING INSTRUCTIONS:** To obtain a 0.005% Chlorophacinone concentration, mix 2.75kg of clean (hull or hullless) oats or wheat with 250mL of Farm & Ranch Brand

LIQUID ROZOL. Mix thoroughly. Allow bait to absorb the product for approximately 30 minutes before using.

**FOR CONTROL OF: GROUND SQUIRRELS (RICHARDSON'S, COLUMBIAN, THIRTEEN-LINED and FRANKLIN'S):** Place 15 g of bait into each ground squirrel hole. Check every 48 hours and rebait holes only where bait has been consumed or removed.–**NORTHERN POCKET GOPHER:** Burrowing pocket gophers throw out low, fan-shaped mounds on either side of their underground tunnel. These lateral tunnels coming to the surface are on the flat side of the fan and these holes are plugged with loose soil. Treatment can be made in one or both of the following ways: 1. Carefully remove the plug on the flat side of the fan and insert 15 g of prepared bait as far down into the hole as possible. Reclose opening being careful not to cover the bait with soil. 2. Using a metal rod, probe 15 to 30 cm deep to locate the main tunnel. Place 15 g of prepared bait into the tunnel and cover the hole so light will not enter the tunnel. Make 2 or 3 treatments per burrow system. Maintain a constant supply of bait for as long as there is gopher activity. Do not apply bait on surface soil. **RATS:** Place 100 to 450 g at intervals of 4.5 to 9 m per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease. **HOUSE MICE:** Place 5 to 15 g at intervals of 2.5 to 3.5 m per placement. Larger placements may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease. **RATS/MICE:** Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**Food Processing, Food Manufacturing, Food Storage and Food Service Areas:**

For areas not directly related to food processing: Use only in non-food or non-feed area where feed, food, packaging and handling equipment are never opened or exposed. For areas where feed or food is processed, served, or stored: In meat and food processing plants (processing areas), use only when plant is not in operation. Remove or cover all food, packaging material and utensils before placing bait in baiting stations. Remove all baits and dead rodents before reuse of the plant (processing areas include storage and service).

**FIRST AID: If in eyes,** hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **If swallowed,** call a poison control centre or doctor immediately for treatment advice. Do not give **any** liquid to the person. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Vitamin K<sub>1</sub> in the form of intramuscular or subcutaneous injections or by oral ingestion is the suggested remedial treatment for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.** May be harmful or fatal if swallowed or absorbed through the skin. Chemical-resistant gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** If swallowed, this material may reduce clotting ability of the blood and cause bleeding. Keep out of lakes, streams or ponds.

**STORAGE:** Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

**DISPOSAL:** Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial regulations for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills. **Rodent Disposal:** Dispose of dead rodents in garbage or by burying.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.