

2018-3817  
2018-10-01

Page 1

## **METHOMYL TECHNICAL INSECTICIDE**

FOR MANUFACTURING, FORMULATING OR REPACKAGING

READ THE LABEL BEFORE USING

**GUARANTEE:** Methomyl 98.7%

REGISTRATION NO. 19139 PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 25 - 50 kg / 100 - 500 kg

DANGER



POISON

Production Agriscience Canada Company  
P. O. Box 730  
7398 Queen's Line  
Chatham, Ontario  
N7M 5L1  
519-352-6350

**PRECAUTIONS:**

- **PREVENT ACCESS BY UNAUTHORIZED PERSONNEL**
- **KEEP OUT OF REACH OF CHILDREN.**
- **MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.**
- Do not breathe dust or spray mist.
- Do not get in eyes, on skin, or on clothing.

**FIRST AID:**

**IF SWALLOWED:**

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:**

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:**

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:**

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**TOXICOLOGICAL INFORMATION:**

ATROPINE IS AN ANTIDOTE - SEEK MEDICAL ATTENTION AT ONCE IN ALL CASES OF SUSPECTED POISONING.

WARNING SYMPTOMS -- Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

**TREATMENT** -- Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg intravenously, every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

**DIRECTIONS FOR USE:**

To be used only in the manufacture of a pesticide which is registered under the Pest Control Products Act.

**STORAGE:**

Store product in original container only, away from other pesticides, fertilizers, food or feed. Not for use or storage in or around the home. Keep container closed.

2018-3817  
2018-10-01

Page 3

**DISPOSAL AND DECONTAMINATION:**

Canadian formulators of this technical should dispose of unwanted active and containers in accordance with municipal and provincial regulations. For additional details and cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.