

2021-2996
2021-07-29

15% PREVENTATIVE LIQUID ALGAECIDE

Controls algae in swimming pool water

ACTIVE INGREDIENT: n-Alkyl (40% C12, 50% C14, 10% C16),
Dimethyl Benzyl Ammonium Chloride....15%

DOMESTIC

REG. N° 20682.26 P.C.P. ACT

NET CONTENTS

RBF INTERNATIONAL LTÉE/LTD
780, RUE NOBEL
ST-JÉRÔME (QUÉBEC) CANADA J7Z 7A3
1 800 433-0590

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING

DIRECTIONS FOR USE

DO NOT apply when swimmers are in the immediate vicinity (a 15-minute restricted-entry interval is recommended).

1. Initial dosage for pool water having no visible algae is 135 mL of **15% PREVENTATIVE LIQUID ALGAECIDE** per 10,000 L of pool water.
2. Weekly maintenance dosage is 70 mL of **15% PREVENTATIVE LIQUID ALGAECIDE** per 10,000 L of pool water.
3. For pool water having visible algae, superchlorinate following label instructions of superchlorination product employed and wait for at least 24 hours before adding weekly maintenance dosage.
4. For visible algae growth, when superchlorination product is not employed, add 340 mL of **15% PREVENTATIVE LIQUID ALGAECIDE** per 10,000 L of pool water.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Wear rubber gloves when handling this concentrate. Harmful if swallowed. Do not mix with any other chemical. Avoid contact with skin and eyes.

DISPOSAL

Rinse container thoroughly before discarding. Do not reuse empty container. Wrap and dispose of empty container with household garbage. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer or the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.