

*Label*

*Principal Display Panel*

**BEAPHAR®  
FLEA & TICK COLLAR FOR DOGS**

Insecticidal Flea and Tick Collar for Dogs

Kills fleas and ticks on dogs.  
Kills up to 4 months.

DOMESTIC

READ THE LABEL AND THE ENCLOSED INSERT BEFORE USING. RETAIN INSERT  
AND PACKAGING INFORMATION FOR FUTURE REFERENCE

FOR USE ONLY ON DOGS

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: TETRACHLORVINPHOS 15.1 %

This product contains the toxic formulant Phenol at 0.018%  
Warning, contains the allergen, epoxidized soyabean oil.

REG. No. 21360 PCP ACT

Net Contents: 26 g WHITE COLLAR - 65 CM

Beaphar B.V.  
Postbus 7, 8100 AA  
Raalte, Holland

Sherley's Canada Inc.  
1-5765 Turner Road  
Suite 526  
Nanaimo, BC  
V9T 6M4  
1-800-668-7387

*Secondary Display Panel*

**DIRECTIONS FOR USE:** Do NOT remove the collar from its package and do not unroll collar until ready to use. This collar is intended for use as an insecticide generator dog collar. Remove the collar from the package, unroll and stretch to activate. This collar releases the active ingredient at time of use and assures full activity. Place one collar around the dog's neck, allowing a spacing of 2 fingers between the collar and neck, and buckle in place. Leave 5–8 cm on the collar for extra adjustment. Ample room should be left for growth, if being used on puppies. Cut off any excess length and dispose of it in household garbage. When collar is first worn, observe the pet's neck area every few days for irritation or adverse reaction. Do NOT use on dogs under 12 weeks of age. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. Do NOT use this product on dogs at the same time or within 30 days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides or chemicals. Replace this collar no more than once every 4 months. Replace collar every 4 months for continual pest control in situations where there is an ongoing risk of pest infestation. **RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.**

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal treatment of the premises may be required. A veterinarian should be consulted when flea infestation continues to be a problem on the pet.

## **PRECAUTIONS**

**HUMANS:** KEEP OUT OF REACH OF CHILDREN. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on the collar during storage. Dust may irritate eyes, nose and throat. Harmful if collar is sucked or chewed, or if dust is inhaled or absorbed through skin. Avoid contact with mouth, eyes, skin and clothing. Avoid breathing dust from collar. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils or places where food is prepared or stored.

**PETS:** Dust may irritate eyes, nose and throat. Harmful if collar is chewed or swallowed. Avoid contact with the dog's mouth, face, eyes and genitals. Avoid letting the pet inhale the dust from the collar.

## **FIRST AID**

### **HUMANS:**

IF SUCKED OR CHEWED, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF UNINTENTIONAL SKIN OR CLOTHING CONTACT OCCURS, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF DUST FROM COLLAR GETS IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF DUST FROM COLLAR IS INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**PETS:** Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If swallowed or chewed,

consult a veterinarian.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

**TOXICOLOGY INFORMATION:** This product contains a pesticide that is a cholinesterase inhibitor (an anti-cholinesterase compound). Symptoms of human poisoning may include headache, weakness, sweating, blurred vision, nausea and diarrhea. Obtain medical attention or call a poison control centre at once. Atropine is antidotal. Acute symptoms of overdosage in dogs include diarrhea, salivation, vomiting, muscular tremors and weakness. Contact a veterinarian immediately.

**STORAGE:** Store in original, unopened container apart from food and feed in a cool place.

**DISPOSAL:** Dispose of any excess collar, packaging and unwanted (used) collars in household garbage.

**ENVIRONMENTAL HAZARDS**

Toxic to fish and other aquatic life. Toxic to birds. Do not contaminate water by disposal of this product.

*Insert*

*Principal Display Panel*

**BEAPHAR®  
FLEA & TICK COLLAR FOR DOGS**

Insecticidal Flea and Tick Collar for Dogs

Kills fleas and ticks on dogs.  
Kills up to 4 months.

DOMESTIC

READ THE LABEL AND THE ENCLOSED INSERT BEFORE USING. RETAIN INSERT  
AND PACKAGING INFORMATION FOR FUTURE REFERENCE

FOR USE ONLY ON DOGS

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: TETRACHLORVINPHOS 15.1 %

This product contains the toxic formulant Phenol at 0.018%  
Warning, contains the allergen, epoxidized soyabean oil.

REG. No. 21360 PCP ACT

Beaphar B.V.  
Postbus 7, 8100 AA  
Raalte, Holland

Sherley's Canada Inc.  
1-5765 Turner Road  
Suite 526  
Nanaimo, BC  
V9T 6M4  
1-800-668-7387

*Secondary Display Panel*

**DIRECTIONS FOR USE:** Do NOT remove the collar from its package and do not unroll collar until ready to use. This collar is intended for use as an insecticide generator dog collar. Remove the collar from the package, unroll and stretch to activate. This collar releases the active ingredient at time of use and assures full activity. Place one collar

around the dog's neck, allowing a spacing of 2 fingers between the collar and neck, and buckle in place. Leave 5–8 cm on the collar for extra adjustment. Ample room should be left for growth, if being used on puppies. Cut off any excess length and dispose of it in household garbage. When collar is first worn, observe the pet's neck area every few days for irritation or adverse reaction. Do NOT use on dogs under 12 weeks of age. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. Do NOT use this product on dogs at the same time or within 30 days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides or chemicals. Replace this collar no more than once every 4 months. Replace collar every 4 months for continual pest control in situations where there is an ongoing risk of pest infestation. **RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.**

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal treatment of the premises may be required. A veterinarian should be consulted when flea infestation continues to be a problem on the pet.

### **PRECAUTIONS**

**HUMANS:** KEEP OUT OF REACH OF CHILDREN. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on the collar during storage. Dust may irritate eyes, nose and throat. Harmful if collar is sucked or chewed, or if dust is inhaled or absorbed through skin. Avoid contact with mouth, eyes, skin and clothing. Avoid breathing dust from collar. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils or places where food is prepared or stored.

**PETS:** Dust may irritate eyes, nose and throat. Harmful if collar is chewed or swallowed. Avoid contact with the dog's mouth, face, eyes and genitals. Avoid letting the pet inhale the dust from the collar.

### **FIRST AID**

#### **HUMANS:**

IF SUCKED OR CHEWED, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF UNINTENTIONAL SKIN OR CLOTHING CONTACT OCCURS, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF DUST FROM COLLAR GETS IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF DUST FROM COLLAR IS INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**PETS:** Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If swallowed or chewed, consult a veterinarian.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

**TOXICOLOGY INFORMATION:** This product contains a pesticide that is a cholinesterase inhibitor (an anti-cholinesterase compound). Symptoms of human poisoning may include headache, weakness, sweating, blurred vision, nausea and diarrhea. Obtain medical attention or call a poison control centre at once. Atropine is antidotal. Acute symptoms of overdosage in dogs include diarrhea, salivation, vomiting, muscular tremors and weakness. Contact a veterinarian immediately.

**DISPOSAL:** Dispose of any excess collar, packaging and unwanted (used) collars in household garbage.

**ENVIRONMENTAL HAZARDS**

Toxic to fish and other aquatic life. Toxic to birds. Do not contaminate water by disposal of this product.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.