

2023-1424
2024-01-08
Label correction 2024-08-19

This label has been updated according to the re-evaluation decision for Putrescent Whole Egg Solids RVD2023-01. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 01-13-2025 in accordance with the phase out period set out in the re-evaluation decision for Putrescent Whole Egg Solids RVD2023-01. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

2023-1424
2024-01-08
Label correction 2024-08-19

(Box/Kit Front Panel)

PROVEN PROTECTION

For Plants, Trees and Shrubs

GUARANTEED EFFECTIVE

DEER-AWAY

Big Game Repellent

Concentrate 2103

PASTE

For repelling Black-Tailed Deer (Mule Deer), White-Tailed Deer and Roosevelt Elk that browse or uproot
Conifer seedlings. Repels White-Tailed Deer from listed Ornamentals and Fruit Trees.

ONE-LITRE KIT – LIQUID FOR SPRAYING

Treats 15 – 18 Ornamental Shrubs 120 cm High

DOMESTIC

READ THE LABEL AND ENCLOSED INSTRUCTION SHEET BEFORE USING

MUST BE USED WITH FORMULA 2104

KEEP OUT OF REACH OF CHILDREN

POTENTIAL SENSITIZER

REG. NO. 17955 P.C.P. ACT

ACTIVE INGREDIENT: Putrescent Whole Egg Solids37.0%

Warning, contains the allergen eggs

Canadian Patent 1,048,408

US Patents 3,980,773; 4,065,576; 4,065,577

Concentrated: This kit makes ONE LITRE

(after mixing Concentrate 2103 and Formula 2104 with water)

NET CONTENTS OF BOX:

128 mL Concentrate 2103

120 mL of Formula 2104

Woodstream Canada Corporation

Unit 1, 25 Bramtree Court, Brampton, ON L6S 6G2

1-800-800-1819

(Side of Box)

DEER-AWAY
BIG GAME REPELLENT
Concentrate 2103

INSTRUCTIONS: See enclosed sheet for mixing and dilution instructions.

RECOMMENDED PLANTS:

- **Yew, Scotch Pine, Arborvitae, Douglas Fir, Flowering Crab, Hemlock, Honeysuckle:**
Apply to terminal to point of run-off. For large trees, apply DEER-AWAY Big Game Repellent Concentrate 2103 only to lateral branches that are within animal's reach. Repeat as necessary.
- **Ornamental Shrubs:**
DEER-AWAY Big Game Repellent Concentrate 2103 is not recommended for use during bud break or flower bloom, as foliage injury and petal drop may occur.
- **Fruit Trees:**
For use on bearing and non-bearing trees. On bearing trees use before flowering and leafing out and/or after harvest.
- **Conifer Seedlings:**
Seedlings to be protected from winter browsing and seedlings in nurseries and greenhouses must be dormant and frost hardy before treatment. Note: Browning and drop of old needles may occur on some evergreen species after using this product.
- **Established Plantations, Nurseries, Large Greenhouses:**
See enclosed instructions sheet.

NOT FOR USE ON EDIBLE CROPS. See specific directions for application to fruit trees.

FIRST AID: If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

PRECAUTIONS: May cause sensitization. Avoid contact with skin, eyes, and clothing. Avoid inhaling dust. For good hygiene practice, wear waterproof gloves when handling this product. DO NOT apply if allergic to eggs. Wash hands, arms and face with soap and water after use. Do not store near or contaminate feed or foodstuffs. If spillage of concentrate occurs, dilute with water and flush away.

2023-1424
2024-01-08
Label correction 2024-08-19

(2103 Pouch)

DEER-AWAY
Big Game Repellent

Concentrate 2103

For repelling Black-Tailed Deer (Mule Deer), White-Tailed Deer and Roosevelt Elk that browse or uproot Conifer Seedlings.

Repels White-Tailed Deer from listed Ornamentals and Fruit Trees

DOMESTIC

REG.NO. 17955 P.C.P. ACT

READ THE LABEL AND ENCLOSED INSTRUCTION SHEET BEFORE USING
POTENTIAL SENSITIZER

MUST BE USED IN CONJUNCTION WITH FORMULA 2104

ACTIVE INGREDIENT: Putrescent Whole Egg Solids.....37.0%

Warning, contains the allergen eggs

Knead thoroughly before mixing!

KEEP OUT OF REACH OF CHILDREN

Protect from freezing.

If frozen, allow to warm to room temperature and knead thoroughly before using.

NET CONTENTS: 128 mL

Makes ONE LITRE of solution (after mixing with water and Formula 2104)

DISPOSAL: Do not reuse empty container. Wrap and dispose of empty container and leftover spray in household garbage.

FIRST AID: If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

PRECAUTIONS: For good hygiene practice, wear waterproof gloves when handling this product. Wash hands, arms and face with soap and water. Do not store near or contaminate feed or foodstuffs. If spillage occurs, dilute with water and flush away. May cause sensitization. Avoid contact with skin, eyes or clothing. Avoid inhaling dust. DO NOT apply if allergic to eggs.

Woodstream Canada Corporation
Unit 1, 25 Bramtree Court, Brampton, ON L6S 6G2
1-800-800-1819

2023-1424
2024-01-08
Label correction 2024-08-19

(Formula 2104 Pouch)

DEER-AWAY
Big Game Repellent
Concentrate 2103

Formula 2104
(use ONLY with Concentrate 2103)

Knead thoroughly before mixing!

KEEP OUT OF REACH OF CHILDREN

Wash hands, arms and face with soap and water.
Do not store near or contaminate feed or foodstuffs.
If spillage of concentrate occurs, dilute with water and flush away.
May cause sensitization. Avoid contact with skin, eyes or clothing. Avoid inhaling dust. DO NOT apply if allergic to eggs.

PROTECT FROM FREEZING.

If frozen, allow to warm to room temperature and stir thoroughly before using.

READ THE LABEL AND ENCLOSED INSTRUCTION SHEET BEFORE USING.

POTENTIAL SENSITIZER

MUST BE USED IN CONJUNCTION WITH CONCENTRATE 2103.

Net Contents 120 mL

Makes ONE LITRE of solution
(after mixing with water and Concentrate 2103)

Woodstream Canada Corporation
Unit 1, 25 Bramtree Court, Brampton, ON L6S 6G2
1-800-800-1819

2023-1424
2024-01-08
Label correction 2024-08-19

(Instruction Sheet)

PROVEN PROTECTION
For Plants, Trees and Shrubs

DEER-AWAY
Big Game Repellent

Concentrate 2103

For repelling Black-Tailed Deer (Mule Deer), White-Tailed Deer and Roosevelt Elk that browse or uproot
Conifer Seedlings
Repels White-Tailed Deer from listed Ornamental and Fruit Trees

KEEP OUT OF REACH OF CHILDREN.

DOMESTIC
READ THE LABEL AND INSTRUCTION SHEET BEFORE USING
POTENTIAL SENSITIZER

REG.NO. 17955 P.C.P. ACT
ACTIVE INGREDIENT: Putrescent Whole Egg Solids..... 37.0 %
Warning, contains the allergen eggs

Canadian Patent 1,048,408
U.S. Patents 3,980,773; 4,065,576; 4,065,577

Woodstream Canada Corporation
Unit 1, 25 Bramtree Court, Brampton, ON L6S 6G2

RECOMMENDED PLANTS:

Not for use on edible crops:

See specific directions for application to fruit trees.

- **Yew, Scotch Pine, Arborvitae, Douglas Fir, Flowering Crab, Hemlock, Honeysuckle:**
Apply to terminal to point of run-off. For large trees, apply DEER-AWAY Big Game Repellent Concentrate 2103 only to lateral branches that are within animal's reach. Repeat as necessary.
- **Ornamental Shrubs:**
DEER-AWAY Big Game Repellent Concentrate 2103 is not recommended for use during bud break or flower bloom, as foliage injury and petal drop may occur.

- **Fruit Trees:**

For use on bearing and non-bearing trees. On bearing trees use before flowering and leafing out and/or after harvest.

- **Conifer Seedlings:**

Seedlings to be protected from winter browsing and seedlings in nurseries and greenhouses must be dormant and frost hardy before treatment. **Note:** Browning and drop of old needles may occur on some evergreen species.

DIRECTIONS FOR USE: Use only as directed on this label. DO NOT apply to any body of water. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

STORAGE: Store at room temperature: Protect from heat and freezing. Store this product away from food or feed.

INSTRUCTIONS: Kneading or Stirring and Dilution:

Note: Containers are not filled to capacity, but are filled by weight.

DILUTE before using: Concentrate 2103 must be mixed with Formula 2104 and water to produce a ready-to-use spray. Read all directions before mixing.

Knead or stir individual ingredients before mixing:

Before diluting, thoroughly knead or stir contents of both containers individually. If Concentrate 2103 has caked in a jar or bucket due to unavoidable settling which occurs during shipment, break cake with a stiff knife or screwdriver, stir thoroughly until free of lumps.

1-Litre Kit: To make entire one litre of ready-to-use spray:

1. Slowly and carefully knead pouch of Concentrate 2103 to break up solids that have settled during shipment. Knead gradually, starting at liquid end. Knead until lumps are completely liquified.
2. Briefly knead Formula 2104 pouch.
3. Cut off corner of pouch containing Concentrate 2103 and empty entire contents into a one-litre trigger sprayer, pressurized hand-held or back-pack garden sprayer.
4. Repeat step 3 with pouch containing Formula 2104.
5. Add 750 mL warm water to make a total volume of one litre. Shake or stir thoroughly.

2-Litre Kit: To make entire two litres of ready-to-use spray:

1. Empty entire bottle of Concentrate 2103 into a suitable mixing container. Add 1.5 litres of warm water and mix thoroughly.
2. Add entire contents of Formula 2104 jar. Shake or stir thoroughly.

To make less than entire package:

1. Knead both packets (in 1-Litre Kit) or stir both containers (in 2-Litre Kit) as explained above.
2. Mix 1 part Concentrate 2103, 1 part Formula 2104, 6 parts water. Blend until thoroughly mixed.

Use promptly after mixing: Mixed solution spoils quickly above 5° C. Do not store longer than one day; discard unused solution.

Clean sprayer: If sprayer is re-used, clean with cold water immediately after use.

Seal tightly for storage: If saving part of unmixed ingredients to use later, close containers tightly to prevent evaporation. Store this product away from food or feed.

FIRST AID: If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

PRECAUTIONS: May cause sensitization. Avoid contact with skin, eyes, and clothing. Avoid inhaling dust. For good hygiene practice, wear waterproof gloves when handling this product. DO NOT apply if allergic to eggs. Wash hands, arms and face with soap and water. Do not store near or contaminate feed or foodstuffs. If spillage occurs, dilute with water and flush away.

METHOD OF APPLICATION:

Before applying, read directions carefully.

Spray final mixed DEER-AWAY Big Game Repellent Concentrate 2103 solution in all areas of plants likely to be browsed.

Apply only in diluted form as specified. Do not apply excessively to seedlings. Mixtures or serial applications of DEER-AWAY Big Game Repellent Concentrate 2103 and TMTD (Thiram) may reduce repellency during winter.

NOT approved for use on edible crops.

See specific directions for application to fruit trees.

Timing of Treatment: To protect against winter browsing, apply repellent upon onset of browsing. If browsing extends beyond two months, seedlings may be re-treated. To protect against spring browsing of new growth, apply upon onset of browsing, but preferably after bud break and before new shoots exceed 2.5 cm in length. Do not retreat new growth. If spring browsing reaches unacceptable proportions before bud break, repellent may be applied to unopened buds, but must then be re-applied after new growth averages 10 cm in length if browsing still persists.

Weather Conditions: Apply only to dry plants and only under weather conditions favouring prompt drying of spray (lightly breezy, mild dry weather with low relative humidity). If spray is used on wet plants or in damp, cold, foggy or misty weather, or if precipitation or freezing occurs before spray can dry protection may be partially or **completely lost**.

DISPOSAL: Do not reuse empty container. Wrap and dispose of empty container and leftover spray in household garbage.

CARE OF EQUIPMENT: Promptly discard leftover spray. Rinse spraying equipment thoroughly with cold water immediately after use. Do not contaminate water by cleaning of equipment or discarding of waste.

ESTABLISHED PLANTATIONS: For use on individual seedlings, use a pressurized hand-held or backpack garden sprayer. Hold nozzle of spray wand about 25 cm vertically above tip of terminal shoot. On conifers, spray the terminal and first whorl to run-off. Two litres of diluted DEER-AWAY Big Game Repellent Concentrate 2103 mixture should be applied per 200 seedlings 60 to 120 cm tall. Treat each seedling.

NURSERIES OR LARGE GREENHOUSES: Use a pressurized tank and spray boom and assure complete coverage of seedlings to run-off by proper adjustment of tank pressure, nozzle size, number and opening, boom height above seed bed and speed of travel. This will result in about 2 mL of solution per seedling. Seedlings in nurseries and greenhouses must be sprayed as close to lifting and outplanting as possible for maximum browse protection. Re-spray before lifting if more than 6 weeks have elapsed between first application and lifting of stock.