

2015-07-31
2011-0763

**POULIN'S PEST CONTROL
GOPHER DOOM
COMMERCIAL**

**(Ready to use Bait)
KILLS GROUND SQUIRRELS,
VOLES, HOUSE MICE, NORWAY OR
ROOF RATS, AND POCKET GOPHERS**

REGISTRATION NO. 22608 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

WARNING: CONTAINS THE ALLERGEN WHEAT

**Only to be used by certified pest control operators, farmers and persons authorized
in government-approved pest control programs.**

GUARANTEE CHLOROPHACINONE 0.005%

NET CONTENTS

99.2 g, 454g, 2.27kg, 4.54kg, 11.36kg

**POULIN'S PEST CONTROL
24 Poulin Drive, Winnipeg, Manitoba R2H 0S9
1-888-768-5467**

DIRECTIONS FOR USE - GENERAL: Users should remove clothing immediately if pesticide gets inside. Then wash skin thoroughly and put on clean clothing.

USE LIMITATIONS – GENERAL: DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

DIRECTIONS FOR USE – RATS & HOUSE MICE

FOR THE CONTROL OF NORWAY OR ROOF RATS AND HOUSE MICE: Place the bait stations near living and feeding areas and water supplies. Place Poulin's Gopher Doom behind boards next to wall, in rat holes, between walls or other protected areas. Replace baits that are contaminated, spoiled or not fresh. Replenish bait supply as long as there is evidence of rat population. Protect bait against access of children, pets, livestock and wildlife.

FOR RATS: Each station should contain at least 100 g of bait. Maintain constant supply of bait at stations for at least 10 days.

FOR HOUSE MICE: Tablespoon amounts! 10-15 g of bait should be placed in 3-4 meter intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

Food Processing, Food Manufacturing, Food Storage and Food Service Areas:

For areas not directly related to food processing: Use only in non-food or non-feed area where feed, food, packaging and handling equipment are never opened or exposed. For areas where feed or food is processed, served, or stored: In meat and food processing plants (processing areas), use only when plant is not in operation. Remove or cover all food, packaging material and utensils before placing bait in baiting stations. Remove all baits and dead rodents before reuse of the plant (processing areas include storage and service).

USE LIMITATIONS – RATS & HOUSE MICE (IN AND AROUND BUILDINGS):

Use indoors and outdoors within 15 meters of buildings. Bait **MUST** be placed either in tamper-resistant bait stations or in locations not accessible to children, pets, livestock or non-target wildlife. Bait applied outdoors and above-ground **MUST** be placed in bait stations. Rodenticide bait can be placed in tamper-resistant bait stations along the fence line of properties, outside of the 15-metre limit but within a 100 meters of buildings, if the station is securely fastened (e.g., nailed down) to the fence or the ground.

To ensure safe use of product, tamper-resistant bait stations must have following characteristics:

Bait stations (**tier 3**) used for the placement of rodenticide bait indoors, in locations not accessible to pets or livestock must have the following characteristics: (1) be constructed of high-strength material (e.g., metal or injection molded plastic) and resistant to destruction by children; (2) have an entrance designed so that children cannot reach the bait; (3) have an internal structure that prevents bait from being shaken loose; (4) have an access panel that fastens securely and locks (e.g., metal screw or padlock); and (5) bear the product name, active ingredient, guarantee, registration number, "WARNING POISON", and skull and crossbones symbol.

Bait station (**tier 2**) used for the placement of rodenticide bait indoors, in locations accessible to pets or livestock must have the following characteristics, in addition to those outlined above for tier 3 bait stations; (1) resistant to destruction by non-target animals; and (2) have an entrance designed so that non-target animals cannot reach the bait.

Bait stations (**tier 1**) used outdoors, above-ground, in locations accessible to children, pets and non-target wildlife must have the following characteristic, in addition to those outlined above for tier 2 and 3 bait stations: (1) be resistant to destruction or weakening by elements of typical non-catastrophic weather (such as snow, rain, extremes of temperature and humidity, direct sunshine,, etc.).

USE LIMITATIONS – RATS & HOUSE MICE (OTHER OUTDOOR USES):

Bait placed outdoors, above-ground in residential areas, public areas or any other sites that are or will be open to the public (for example, Pick-Your-Own sites), or in locations accessible to pets or livestock, **MUST** be placed in tier 1 bait stations. The bait stations used in these sites **MUST** meet the characteristics described under **USE LIMITATIONS – RATS & HOUSE MICE (IN AND AROUND BUILDINGS)**.

DIRECTIONS FOR USE – GROUND SQUIRRELS:

FOR RICHARDSON’S, COLUMBIAN, THIRTEEN-LINED AND FRANKLIN GROUND SQUIRREL: Individual treatment of holes reduces the possibility of secondary poisoning and increases the degree of control. Locate all the holes in the area and place 15 g of Poulin’s Gopher Doom deep into each hole. Revisit the site within 48 hours and rebait all holes where bait has been consumed or removed. Do not rebait holes where bait has been undisturbed. *Remove and bury any bodies which are present above ground. A third visit may be needed for full control.

BAIT STATION CONTROL: Determine the approximate population present by visual count over a two-day period. Establish protected bait stations or tamper proof boxes every 30-60 meters in areas where the problem is present (use 1 box for every 50 animals counted). Bait stations should bear the product name, active ingredient, guarantee, registration number, “WARNING POISON”, and the skull and crossbones symbol. Use 500 g per station and check daily, and maintain an uninterrupted supply for 3 weeks or until feeding has stopped. *Remove and bury any bodies which are present above ground.

USE LIMITATIONS – GROUND SQUIRRELS: Bait placed outdoors, above-ground in residential areas, public areas, or any other sites that are or will be open to the public (for example, Pick-Your-Own sites) or in locations accessible to pets or livestock **MUST** be placed in tier 1 bait stations. The bait stations used in these sites **MUST** meet the characteristics described under **USE LIMITATIONS – RATS & MICE (IN AND AROUND BUILDINGS)**.

DIRECTIONS FOR USE – POCKET GOPHERS:

NORTHERN POCKET GOPHERS: Pocket gophers are very territorial and inspect their tunnels on a regular basis. Any disturbance will usually induce their inspection. Place bait – back from actual opening of the burrow, it will provide a good opportunity to allow the rodent to feed. If the bait is placed too close to the damaged tunnel opening, the bait usually gets pushed aside or buried.

HAND PLACEMENT: Carefully open burrow and place 15 g of bait well into the tunnel. Carefully close burrows without covering bait. A repeat treatment may be necessary where burrows have been reopened and are in active use. The unused bait can be stored in its original container and keep its killing effect. It is not affected by heat or cold.

To ensure full control, cover any hole and check the following day for openings to determine whether the hole is active. Place 15 g of Poulin’s Gopher Doom in each active hole (follow directions for hand baiting).

DIRECTIONS FOR USE – VOLES:

FOR VOLES (*Microtus* sp.): Apply after harvest, preferably before snowfall and leaf fall or lodging of grass. Apply ground broadcast at the rate of 11 kg./ha. If population is high, a second application, 1-2 months after the first treatment may be necessary (check population by apple quarter index). *Remove and bury bodies which are present.

USE LIMITATIONS – VOLES:

DO NOT use in residential areas, public areas, or in any other sites that are or will be open to the public (for example, Pick-Your-Own sites). DO NOT use in locations accessible to pets or livestock.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Keep away from humans, domestic animals, pets and non-target wildlife. May be harmful or fatal if swallowed or absorbed through the skin. If swallowed or absorbed by humans, domestic animals, pets or non-target wildlife, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.**

When handling product, or when disposing of dead rodents, unconsumed bait and empty containers, wear long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves. In addition, persons loading pellets or bait into mechanical ground equipment, or persons loading/applying with hand-pushed or hand-held equipment must wear a NIOSH-approved particulate-filter respirator and protective eyewear.

FIRST AID:

For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

Note: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

Swallowed: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control centre or a doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: Vitamin K1 in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE:

Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL:

- ◆ Dispose of stale bait and dead rodents in accordance with local requirements.
- ◆ Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- ◆ Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.
- ◆ Note: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on the label. It is an offense under the Pest Control Products Act to use this control product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.