

GROUP	4	HERBICIDE
-------	---	-----------

2,4-D Herbicide Low Volatile Ester

(A Component of EXPRESS® Pack Post-Emergent Herbicide Tank Mix)

This product must be used in combination with EXPRESS® Herbicide Dry Flowable 75%

AGRICULTURAL

READ THE LABEL AND ENCLOSED BOOKLET BEFORE USING

GUARANTEE: 2,4-D, present as 2-ethylhexyl ester.....660 g/L

REGISTRATION NUMBER 23192 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 10 L

WARNING



POISON

E. I. du Pont Canada Company
Agricultural Products
P.O. Box 2300, Streetsville,
Mississauga, Ontario
L5M 2J4
1-800-667-3925

PRECAUTIONS:

- DO NOT APPLY BY AIR.
- KEEP OUT OF THE REACH OF CHILDREN.
- Avoid contact with eyes, skin and clothing. May cause irritation.
- Do not breathe spray mist or vapours. Avoid spray or vapor drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.
- Do not spray when there is any danger of wind, or on exceptionally hot days over 27° C.
- Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles.
- Do not contaminate any body of water. Avoid contamination of foods.
- Do not use in a greenhouse.
- Keep in original container during storage.
- Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
- Keep away from fire or open flame and other sources of heat.
- Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. Do not re-use empty containers.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca.

FIRST AID:

EYE CONTACT: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

SKIN OR CLOTHING CONTACT: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INGESTED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: This product contains petroleum distillates. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS:

This tank mixture contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

PROTECTIVE CLOTHING AND EQUIPMENT:

Handling the concentrate (mixing and loading): wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eye wear (face shield or safety glasses). Rinse gloves before removal. Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5 L.

Handling the dilute spray solution (during application or repairing or cleaning equipment): wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor.

OPERATOR USE PRECAUTIONS:

1. Wear freshly laundered clothing and clean protective equipment daily.
2. Rinse gloves before removal.
3. Wash hands before eating, drinking, using tobacco or using the toilet.
4. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
5. After using this product, remove clothing and launder separately, and promptly, and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
6. After work, remove all clothing and shower using soap and water.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, 2,4-D Herbicide Low Volatile Ester is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant 2,4-D Herbicide Low Volatile Ester and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of 2,4-D Herbicide Low Volatile Ester or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local du Pont representative or the DuPont hotline at 1-800-667-3925 for further information

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Keep water-soluble bags in original outer bag away from moisture.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsing to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence *under the Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

®/MC Registered trademarks/trademarks of E. I. du Pont Company. E. I. du Pont Canada is a licensee. Member of CropLife.

*All other products mentioned are trademarks of their respective companies.

EXPRESS PACK®
POST-EMERGENT HERBICIDE TANK MIX

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION OF BRITISH COLUMBIA FOR AGRICULTURAL USE ON: WHEAT (SPRING AND DURUM), BARLEY, AND SUMMERFALLOW ONLY

COMMERCIAL

READ THE LABELS AND THIS BOOKLET BEFORE USING

WARNING



POISON

NET CONTENTS:

EXPRESS® Herbicide Dry Flowable 75% - 160 GRAMS
2,4-D Herbicide Low Volatile Ester - 10 LITRES

THIS TANK MIX USES:

EXPRESS® Herbicide Dry Flowable 75%,
REG. NO. 22335 *PEST CONTROL PRODUCTS ACT*
GUARANTEE: Tribenuron methyl 75%

GROUP	2	HERBICIDE
--------------	----------	------------------

2,4-D Herbicide Low Volatile Ester,
REG. NO. 23192 *PEST CONTROL PRODUCTS ACT*
GUARANTEE: 2,4-D, present as 2-ethylhexyl ester.....660 g/L

GROUP	4	HERBICIDE
--------------	----------	------------------

E. I. du Pont Canada Company
Agricultural Products
P.O. Box 2300, Streetsville
Mississauga, Ontario
L5M 2J4
1-800-667-3925

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- WARNING - EYE IRRITANT.
- Potential skin sensitizer.
- May be harmful if swallowed, inhaled or absorbed through skin.
- Avoid breathing spray mist or vapours. Use with adequate ventilation.
- DO NOT APPLY BY AIR.
- Do not contaminate any body of water.
- Do not contaminate food.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca.

PROTECTIVE CLOTHING AND EQUIPMENT**HANDLING THE CONCENTRATE (MIXING AND LOADING):**

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eye wear (face shield or safety glasses). Rinse gloves before removal.

Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5 L.

HANDLING THE DILUTE SPRAY SOLUTION (DURING APPLICATION OR REPAIRING OR CLEANING EQUIPMENT):

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal.

Gloves are not required during application when applicator is in an enclosed tractor.

OPERATOR USE PRECAUTIONS:

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

ENVIRONMENTAL PRECAUTIONS:

This tank mixture contains a petroleum distillate, which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

FIRST AID:**EYE CONTACT:**

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

SKIN OR CLOTHING CONTACT:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

INGESTION:

Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

INHALATION:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: This tank mixture contains petroleum distillates. Treat symptomatically.

IMPORTANT

Injury to or loss of desirable trees, shrubs, flowers, vegetables, fruits or other vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on crops.

DO NOT APPLY BY AIR.

Do not apply in high wind or on hot days (over 27 °C).

Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than spring wheat, durum or barley.

A MINIMUM RECROPPING INTERVAL OF 2 MONTHS SHOULD BE LEFT BETWEEN THE APPLICATION OF THIS PRODUCT AND SEEDING OF THE NEXT CROP. After 2 months, the following crops can be seeded: canola, flax, lentils, alfalfa.

DO NOT APPLY EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX WITH A CARRIER OTHER THAN WATER OR ANY ADDITIVE OTHER THAN THOSE INDICATED ON THIS LABEL. Other carriers may accelerate breakdown of this product and reduce its effectiveness.

PRE-HARVEST INTERVALS:

Do not permit lactating dairy animals to graze fields within 7 days after application.

Do not harvest forage or cut hay within 30 days after application.

Withdraw meat animals from treated fields at least 3 days before slaughter.

GENERAL INFORMATION

DIRECTIONS FOR USE:

DuPont EXPRESS PACK® Post-Emergent Herbicide Tank Mix provides control of certain broad-leaved weeds in spring planted wheat (including durum) and barley. ALL THE COMPONENTS IN THE EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX PACKAGE MUST BE APPLIED TOGETHER ACCORDING TO THESE RECOMMENDATIONS. This EXPRESS PACK® Post-Emergent Herbicide Tank Mix package contains 160 grams of EXPRESS® Herbicide Dry Flowable 75% and 10 litres of 2,4-D Herbicide Low Volatile Ester.

Warm, moist growing conditions promote active weed growth and enhance the activity of EXPRESS PACK® Post-Emergent Herbicide Tank Mix by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress such as cold weather, drought stress or excessive heat may not be adequately controlled or suppressed, and regrowth may occur. EXPRESS PACK® Post-Emergent Herbicide Tank Mix may only be applied using ground spray equipment. For best results, ensure thorough spray coverage of target weeds.

EXPRESS PACK® Post-Emergent Herbicide Tank Mix rapidly stops growth of susceptible weeds. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Degree of control and duration of effect depend on weed sensitivity, weed size, spray coverage and growing conditions. Activity of the herbicide mixture may be delayed by cold, dry conditions after application.

FOR WEED CONTROL IN SPRING WHEAT (INCLUDING DURUM) AND BARLEY

EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX:

Apply EXPRESS® Herbicide Dry Flowable 75% at 10 grams per hectare with 2,4-D Herbicide Low Volatile Ester at 0.625 litres per hectare. One container of EXPRESS® Herbicide Dry Flowable 75% (160 grams) with 1 container of 2,4-D Herbicide Low Volatile Ester (10 litres) will treat 16 hectares. Do not exceed 275 kPa spray pressure.

WEEDS CONTROLLED BY EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX:

Annual sunflower	Redroot pigweed
Ball mustard	Russian pigweed
Canada thistle (top growth control)	Russian thistle
Cow cockle	Shepherd's-purse (fall rosettes and spring seedlings)
Flixweed (fall rosettes and spring seedlings)	Stinkweed (fall rosettes and spring seedlings)
Hare's-ear mustard	Sweet clover
Indian mustard	Thyme-leaved spurge
Kochia (2-10 leaf)	Tumble mustard
Lamb's-quarters	Wild mustard
Narrow-leaved hawk's-beard (fall rosettes and spring seedlings)	Wild radish
Prickly lettuce	Wormseed mustard

WEEDS SUPPRESSED BY EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX[†]:

Wild buckwheat

[†] Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

WHEN TO APPLY:

Apply the tank mix from the full 3-leaf stage to just before the flag leaf (shot blade) stage of wheat (including durum) and barley, prior to head emergence. Do not apply after head has emerged.

For best results apply to young, emerged actively growing weeds that are less than 10 cm tall or across, and before the crop canopy closes. Weeds that emerge after treatment will not be controlled.

For wild buckwheat, apply to ACTIVELY GROWING wild buckwheat in the 1 to 3-leaf stage. Under dry conditions, suppression may be reduced. Plants larger than the 3-leaf stage may regrow after treatment.

NOTE:

If rain occurs soon after application control may be reduced. At least 4-6 hours of dry weather is needed to allow EXPRESS PACK® Post-Emergent Herbicide Tank Mix to be absorbed by weed foliage. Environmental conditions that slow the drying of the spray mixture on the foliage such as high relative humidity, cool air temperatures or cloud cover, may increase the time required.

EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX + ASSERT* TANK MIX

For control of WILD OATS and control or suppression of broadleaf weeds in spring (including durum) wheat and barley, apply the tank mix as follows:

TANK MIX PARTNER	RATE
EXPRESS® Herbicide Dry Flowable 75%	10 g/ha
ASSERT*	1.3 to 1.6 L/ha
2,4-D Herbicide Low Volatile Ester	0.6 L/ha

Use the low rate of ASSERT* when the majority of wild oats are in the 1 to 3-leaf stage and the higher rate if wild oats are in the 4-leaf stage. Apply when the crop is in the full 3-leaf stage.

EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX + PUMA*¹²⁰ SUPER TANK MIX

For control of GREEN FOXTAIL (WILD MILLET) and control or suppression of broadleaf weeds in spring (including durum) wheat and barley, apply the tank mix as follows:

TANK MIX PARTNER	RATE
EXPRESS® Herbicide Dry Flowable 75%	10 g/ha
PUMA* ¹²⁰ SUPER	385 mL/ha
2,4-D Herbicide Low Volatile Ester	0.6 L/ha

Apply this tank mix from the full 3-leaf stage to the 6-leaf + 3 tiller stage of wheat (including durum) and the full 3-leaf stage to the 5-leaf + 2 tiller stage of barley.

For green foxtail control, apply at the 1-6 leaf (up to emergence of the 3rd tiller) stage.

FOR WEED CONTROL IN SPRING WHEAT (EXCLUDING DURUM) AND SPRING BARLEY

EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX (EXPRESS® HERBICIDE DRY FLOWABLE 75% + 2,4-D HERBICIDE LOW VOLATILE ESTER) + BANVEL* or BANVEL* II TANK MIX

For control of sulfonylurea resistant KOCHIA and control or suppression of broadleaf weeds on the EXPRESS® Herbicide Dry Flowable 75% label in spring (excluding durum) wheat and barley, apply the tank mix as follows:

TANK MIX PARTNER	RATE
EXPRESS® Herbicide Dry Flowable 75%	10 g/ha
BANVEL* or BANVEL* II	110 mL/ha
2,4-D Herbicide Low Volatile Ester(or acid equivalent)	0.6 L/ha (or 420 g/ha acid equivalent)

Apply the tank mix from the full 3-leaf stage to the 6-leaf + 3 tiller stage of wheat (excluding durum) and barley and to kochia plants that are 2.5 – 10 cm in height. Expect superior results when kochia is small.

CONTROL OF FLIXWEED AND STINKWEED (FALL ROSETTES AND SPRING SEEDLINGS) IN SUMMERFALLOW

Apply EXPRESS® Herbicide Dry Flowable 75% at 10 grams per hectare with 2,4-D Herbicide Low Volatile Ester at 0.63 litres per hectare.

Apply the tank mix in the spring after emergence, up to the early flowering stage of the target weeds. Only weeds that have emerged at the time of application will be controlled.

MIXING INSTRUCTIONS:

1. Add one-third of the necessary volume of water to the spray tank.
2. With the agitator running, add the required amount of EXPRESS® Herbicide Dry Flowable 75%
3. When EXPRESS® Herbicide Dry Flowable 75% is in suspension, add the required amount of the appropriate tank mix partners.
4. **Specific Tank Mix Instructions:**

EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX (EXPRESS® Herbicide Dry Flowable 75% + 2,4-D Herbicide Low Volatile Ester) + Assert*: once EXPRESS® Herbicide Dry Flowable 75% is completely in suspension, add the required amount of Spray Water pH Adjuster and agitate until dissolved. Add the required amount of Assert* while maintaining agitation. Add 2,4-D Herbicide Low Volatile Ester. Addition of a surfactant is not required.

EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX (EXPRESS® Herbicide Dry Flowable 75% + 2,4-D Herbicide Low Volatile Ester) + BANVEL* or BANVEL* II: add 2,4-D Herbicide Low Volatile Ester last. A surfactant is not required with this mixture.

EXPRESS PACK® POST-EMERGENT HERBICIDE TANK MIX (EXPRESS® Herbicide Dry Flowable 75%+ 2,4-D Herbicide Low Volatile Ester) + PUMA*¹²⁰ SUPER: add the required amount of PUMA*¹²⁰ SUPER last. A surfactant is not required with this mixture.

5. Add the rest of the water. If an antifoam agent is required, add this last.
6. Use a minimum spray volume of 55 L/ha.
7. For repeat tank loads, reduce the material remaining in the tank to 10% of the original volume or less before proceeding with step 1. If this is not possible, pre-slurry EXPRESS® Herbicide Dry Flowable 75% in a small amount (5-10 L) of water before adding to the tank. The 2,4-D Herbicide Low Volatile Ester component does not need to be premixed.

Continuous agitation is required to keep the EXPRESS PACK® Post-Emergent Herbicide Tank Mix components in suspension. If the spray preparation is left standing without agitation, thoroughly agitate to resuspend the mixture before spraying. Use spray preparation of EXPRESS PACK® Post-Emergent Herbicide Tank Mix within 24 hours or product degradation may occur.

MINIMUM RECROPPING INTERVAL AND CROP ROTATION GUIDELINES:

A minimum recrop interval of 2 months should be left between the application of this product and seeding of the next crop. The following crops can be seeded two months after application of EXPRESS PACK® Post-Emergent Herbicide Tank Mix: canola, flax, lentils, and alfalfa.

EQUIPMENT-SPRAY VOLUMES:

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh filter screens or larger (metal or nylon).

Continuous agitation is required to keep EXPRESS PACK® Post-Emergent Herbicide Tank Mix in suspension. Apply uniformly and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

Reduce drift with high water volume, low pressure, coarse sprays and drop nozzles.

SPRAYER CLEANUP:

It is difficult to remove all traces of 2,4-D Herbicide Low Volatile Ester from a sprayer. A sprayer used to apply 2,4-D Herbicide Low Volatile Ester should not be used to spray crops sensitive to that herbicide. The following clean-out procedure will remove traces of EXPRESS® Herbicide Dry Flowable 75% from the sprayer.

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes to remove all visible residues.
2. Fill the tank with clean water then add 1 litre household AMMONIA (containing a minimum of 3% ammonia) per 100 litres of water. Fill boom and hoses with solution and allow sprayer to sit for 15 minutes. Drain.
3. Repeat step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of ammonia, rinse the tank, hoses and booms thoroughly with clean water.
5. Dispose of tank rinseate according to Provincial directions.

CAUTION:

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour, which can cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

USE PRECAUTIONS:

Licensed wheat and barley varieties have shown acceptable tolerance to EXPRESS® Herbicide Dry Flowable 75%. Consult DuPont Canada prior to use on unlicensed varieties for crop tolerance information.

- Do not apply to irrigated land where tail water will be used to irrigate other crop land.
- Do not contaminate irrigation water.
- Use a maximum of 10 grams of EXPRESS® herbicide Dry Flowable 75% per hectare per year.
- Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls or calcium deposits.
- EXPRESS PACK® Post-Emergent Herbicide Tank Mix applied to spring wheat, durum wheat or barley that is stressed by severe conditions such as drought, low fertility, saline soils, waterlogged soils (soils at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury, grade or yield loss.
- Do not apply EXPRESS PACK® Post-Emergent Herbicide Tank Mix to spring wheat, durum wheat or barley under sown with legumes or grasses as injury to the forage may result.
- Do not remove soil from treated fields for use on lawns, in gardens or for backfill.
- Under certain conditions such as prolonged cool weather temperature or wide fluctuations in day/night temperatures just prior to or soon after treatment, temporary yellowing and/or crop stunting may occur.

OVERSPRAY OR DRIFT TO IMPORTANT WILDLIFE HABITATS SUCH AS SHELTERBELTS, WETLANDS, SLOUGHS OR DRY SLOUGH BORDERS, OR WOODLOTS, SHOULD BE AVOIDED. LEAVE A 15 METER BUFFER ZONE BETWEEN THE LAST SPRAY SWATH AND THE EDGE OF ANY OF THESE HABITATS.

Control of weeds growing in wheel tracks may be reduced if EXPRESS PACK® Post-Emergent Herbicide Tank Mix is applied under dry, dusty conditions.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, EXPRESS PACK® Post-Emergent Herbicide Tank Mix is both a Group # 2 and Group # 4 herbicide. Any weed population may contain or develop plants naturally resistant to EXPRESS PACK® Post-Emergent Herbicide Tank Mix and other Group # 2 and/or Group # 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of EXPRESS PACK® Post-Emergent Herbicide Tank Mix or other Group # 2 and/or Group # 4 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted.
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local du Pont representative or the du Pont hotline at 1-800-667-3925 for further information.

STORAGE:

Store products in original containers only, away from fertilizer, seeds, food or feed. Not for use or storage in or around the home. Keep container closed. Keep away from fire, heat, or open flame.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsing to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

“®/™ Registered trademarks / trademarks of E. I. du Pont Company. E. I. du Pont Canada Company is a licensee. Member of CropLife.

* All other products mentioned are trademarks of their respective companies.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.