2012-5030 2013-07-31

ALGICIL-50™ Liquid Algaecide CONTROLS ALGAE IN SWIMMING POOL WATERS DOMESTIC CAUTION

POISON DOMESTIC REG. NO. 23224 P.C.P. ACT GUARANTEE: Poly [oxyethylene (dimethyliminio) ethylene (dimethyliminio) ethylene dichloride]50% 1 Litre READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN N. Jonas & Co., Inc. P.O. Box 425 4525 Adams Circle Bensalem, PA USA 19020 215-639-8071

PRECAUTIONS KEEP OUT OF REACH OF CHILDREN Harmful if swallowed. Do not mix with any other chemical. Avoid contact with skin and eyes.

DISPOSAL

Do not re-use empty container. Wrap and dispose of empty container with household garbage. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

2012-5030 2013-07-31

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL HAZARDS: Toxic to aquatic organisms.

DIRECTIONS FOR USE

1. Initial dosage for pool water having no visible algae is 80 mL of Algicil-50 per 10,000 L of pool water.

2. Weekly maintenance dosage is 30 mL of Algicil-50 per 10,000 L of pool water.

3. For pool water having visible algae, superchlorinate, following label instructions of superchlorination product employed and wait for at least 24 hours before adding weekly maintenance dosage.

4. For visible algae, when superchlorination is not employed, add 120 mL of Algicil-50 per 10,000 L of pool water.

Made and Printed in the USA