

2019-7116
2020-08-19

AIR GUARD CONTROL®
KONK®
FLEA KILLER
DOG & CAT SPRAY

PRESSURIZED
CONTACT/RESIDUAL
DOMESTIC

Kills Dog and Cat Fleas, American Dog & Brown Dog Ticks

ACTIVE INGREDIENT:

Pyrethrins 0.056%

Permethrin 0.050%

REG. NO. 23627 P.C.P. ACT

NET CONTENTS: 350g

CAUTION

EXPLOSIVE

READ THE LABEL BEFORE USING

FOR USE ONLY ON DOGS AND CATS

DO NOT USE ON PUPPIES OR KITTENS LESS THAN 8 WEEKS OLD

DO NOT USE ON ANY OTHER ANIMAL

This product may also be used indoors to kill Dog and Cat Fleas, American Dog & Brown Dog Ticks on building surfaces of the home.

KEEP OUT OF REACH OF CHILDREN

Acuity Holdings, Inc.

11627 - 178 STREET

EDMONTON AB T5S 1N6

905-669-9876

Made in Canada

2019-7116
2020-08-19

DIRECTIONS FOR USE: FOR USE ON DOGS OR PUPPIES AND CATS OR KITTENS OVER 8 WEEKS OLD:

DO NOT use on puppies or kittens under 8 weeks of age. Consult a veterinarian before using on aged or pregnant animals. Do not use on sick or nursing animals, or animals receiving drug or other pesticide treatment. This product contains permethrin. DO NOT apply another pest control product such as a spot on, collar or powder which contains permethrin to the treated animal when using KONK® FLEA KILLER DOG & CAT SPRAY. Use ONLY in a well-ventilated area, preferably outdoors. Shake well before using. Hold can in upright position 25-30 cm from dog or cat and spray entire body beginning at tail and working towards the head. Do not spray to animals face, eyes or genitals. As you spray, fluff the hair so that spray will penetrate to the skin but spray only until hair is damp - not wet. Spray ticks directly. Pay special attention to neck, hindquarters, areas under legs, around the tail and between toes. Do not spray more than once a week. . Treatment must be repeated only if the pest problem persists or reoccurs.

For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas. If pest problems persist, a premise treatment with an insecticide labelled for that use may be required. A veterinarian should be consulted when the flea infestations continue to be a problem on the pet.

FOR USE INDOORS TO KILL DOG AND CAT FLEAS, AMERICAN DOG & BROWN DOG TICKS ON BUILDING SURFACES OF THE HOME:

DO NOT APPLY AS A BROADCAST APPLICATION OR AS A SPACE SPRAY. APPLY AS AN INDOOR CRACK AND CREVICE OR SPOT TREATMENT. Spot application is localized to a surface area not more than 0.2 m². Spots are not to be adjoining. The combined area of spots is not to exceed 10% of the total surface area of a room. Crack and crevice is an application directly into narrow openings on the surface of the structure. It does not include the treatment of exposed surfaces. Narrow openings typically occur at expansion joints, utility entry points and along baseboards and mouldings.

Restrictions:

- DO NOT contaminate food/feed. DO NOT apply to surfaces that may come into contact with food/feed.
- DO NOT apply when people or pets are present.
- DO NOT allow people or pets to enter treated areas until sprays have dried.
- DO NOT allow spray to drip or allow drift onto non-target surfaces.
- DO NOT apply to people, livestock or sources of electricity.
- DO NOT apply to furniture, mattresses, linens, toys or clothing.
- DO NOT apply to overhead areas or in confined spaces (for example, attics, crawlspaces).
- Ventilate treated areas after application by opening windows and doors or using fans, where required, to aid in the circulation of air.
- DO NOT apply indoors as a broadcast application or as a space spray.

Spray indoor building surfaces, including cracks and crevices in walls and floors; along and behind baseboards; window and door frames and localized areas of floor and floor coverings. Pet bedding may be entirely treated with this product or replaced with clean, fresh bedding after

2019-7116
2020-08-19

other listed areas are treated. Treatment must only be repeated after 7 days if pest problem persists or reoccurs. For effective flea and tick control indoors, clean any area used by the pet. Cleaning the areas includes vacuuming floors, carpets, furniture, etc. (discard vacuum bag after use) and washing the pet's bedding, living quarters, and surrounding area. Only treat pets with products specifically registered for use on that type of pet.

PRECAUTIONS:

PHYSICAL AND CHEMICAL HAZARDS: Contents under pressure. Do not place in hot water or near radiators, stoves or other sources of heat or store at temperatures over 50°C. Keep away from open flames or sparks.

HUMANS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed, inhaled or absorbed through skin. Avoid skin contact by wearing rubber gloves. Avoid contact with mouth, eyes, skin and clothing.

Avoid contact with treated animal until dried. After handling or applying, wash gloved hands (or any other skin that came into contact with the product) with soap and water.

PETS: Harmful if swallowed or inhaled. May irritate eyes, nose and throat. Discontinue use if dermal or nasal irritation develops. Toxic to fish and other aquatic life. Cover fish aquaria in vicinity of treatment area. Toxic to birds.

FIRST AID:

HUMANS: IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. **IF IN EYES,** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **IF ON SKIN OR CLOTHING,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **IF SW ALLOWED,** call a poison control centre or doctor immediately for treatment advice.

Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person.

Do not give anything by mouth to an unconscious person.

PETS: Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

DISPOSAL; When container is empty, press valve to release pressure. Dispose of empty container in regular household garbage. Do Not puncture or incinerate container.