

**LATER'S
ORIGINS™
MULTI-PURPOSE YARD & GARDEN INSECT CONTROL**

For use on fruit, vegetables and ornamentals.

- Water-based
- Concentrated Insecticide
- Low odour

**READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

DOMESTIC

REG. NO. 24849 P.C.P. ACT

GUARANTEE: Permethrin 0.5%

1 L

Peel here for instructions. Press to reseal.

Sure-Gro IP Inc.
P.O. Box 21001
Brantford ON N3R7J1
1 800 268-2806

LATER'S MULTI-PURPOSE YARD & GARDEN INSECT CONTROL

DIRECTIONS: Multi-Purpose Yard & Garden Insect Control has no conspicuous odor. Water, not petroleum distillates is used as the diluent. Mix with water at the rates listed below. Spray for thorough coverage of upper and lower leaf surfaces or treated area. Apply when insects or damage first appears and then as necessary.

NOTE: Do not use when air temperature is less than 12⁰C or greater than 30⁰C. Use above 30⁰C could reduce the effectiveness of Permethrin.

5 mL = 1 teaspoon

15 mL = 1 tablespoon

HOW TO USE: SHAKE WELL BEFORE USING. *Rates shown are for 1 L water. Do not apply within 7 days of harvest, unless otherwise indicated.

APPLES: Eastern Tent Caterpillars, White Apple Leafhoppers, Plum Curculios and Eye Spotted Bud Moths: 25 mL*. Apple Maggots, Codling Moths, Lesser Appleworms, Leafrollers, Tentiform Leafminers, Tarnished Plant Bugs, Mullein Bugs: 50 mL*.

GRAPES: Do not apply within 21 days of harvest. Grape Berry Moths: 37 mL*. Grape Leafhoppers: 18.5 mL*.

PLUMS: Plum Curculios: 50 mL*.

PEACHES, NECTARINES: Oriental Fruit Moths, Plum Curculios, Plant Bugs: 50 mL*.

PEARS (Eastern Canada): Pear Psylla (nymphs & adults), Codling Moths, Green Fruitworms: 50 mL*.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS: Do not apply to cabbage, cauliflower and brussels sprouts within 3 days of harvest and to broccoli within 7 days. Imported Cabbage Worms, Cabbage Loopers, Crucifera Flea Beetles, Diamond Back Moth Larvae: 12.5 mL*.

SWEET CORN: European Corn Borers: 25 mL*. Apply before tassels show. Treat at 5 day intervals until early silk stage. Corn Earworms: 25 mL*. When 25% of ears show silk, start spraying and fully cover ears and silk. Make 3 more treatments at 4 day intervals. (3 day intervals in hot weather). Sap Beetles: 25 mL*. Apply when pink spotted beetles first appear. Do not apply with 1 day of harvest.

POTATOES, TOMATOES: Do not apply within 1 day of harvest. Colorado Potato Beetles, Potato Flea Beetles, Potato Leafhoppers, Whiteflies, Tarnished Plant Bugs: 18.5 mL*.

TOMATOES (Greenhouse): Do not apply within 1 day of harvest. Whiteflies: 18.5 mL*.

DECIDUOUS (leafy) PLANTS: Eastern and Forest Tent Caterpillars, Gypsy Moths, Open Feeding Aphids, Fall and Euonymus Webworms: 17.5 mL*. Apply at first sign of feeding. Repeat once if reinfestation occurs.

CONIFEROUS (needle) PLANTS: Spruce Budworms, Douglas Fir Tussock Moths, Whitemarked Tussock Moths, Coneworms, Yellow-headed Spruce Sawflies, Open Feeding Sawflies: 12.5 mL*. Apply at first sign of damage. Repeat once if re-infestation occurs.

Test all tree and shrub species for phototoxicity prior to application of permethrin. Spray an inconspicuous part of the tree or shrub. Wait 24 hours to see if any adverse effect occurs.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Keep children and pets away from treated areas, at least until spray has dried. Do not use as a space spray. Avoid splashing concentrate in eyes or on hands. Do not inhale spray mist. Avoid contamination of food, feed, cooking utensils or drinking water. Wash thoroughly before eating, drinking and smoking.

ENVIRONMENTAL HAZARDS: This product is very toxic to fish. Do not contaminate lakes, ponds, streams and domestic or irrigation water supplies. This product is very toxic to bees: avoid spraying when bees are foraging. Spray deposit should be dry before bees commence foraging in treated crop.

FIRST AID:

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE: Keep product in original container. Store in a cool, dry place away from food. Do not freeze.

DISPOSAL: Do not reuse empty container. Wrap container and discard in household garbage.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.