

CHECK-MARK ROC-622

Bulk

Rodenticide Pellets

Kills Rats and Mice with a Single Feeding

Kills Warfarin-Resistant Norway Rats

COMMERCIAL

GUARANTEE

Bromadiolone.... 0.005%

Warning, contains the allergens peanuts and wheat.

REGISTRATION NO. 24354 PEST CONTROL PRODUCTS ACT

**KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING**

Only to be used by certified pest control operators, farmers, and persons authorized in government-approved pest control programs.

WARNING POISON

NET CONTENTS: 10 Kg

Check-Mark

Div. of/de NCH Canada Inc. 247 Orenda Road, Brampton, Ontario L6T 1E6
Toronto * Montréal * Vancouver * Calgary * Moncton 905-457-5220

DIRECTIONS FOR USE:

Use Bromadiolone pellets for the control of Norway rats, Roof rats, and House mice indoors and outdoors within 15 metres of residential, industrial, commercial, agricultural, and public buildings. Rodenticide bait can be placed in tamper-resistant bait stations along the fence line of properties, outside of the 15-metre limit but within a 100 metres of buildings, if the station is securely fastened (e.g., nailed down) to the fence or the ground.

USE LIMITATIONS: Bait MUST be placed either in tamper-resistant bait stations or in locations not accessible to children, pets, livestock, or non-target wildlife. Bait applied outdoors and above-ground MUST be placed in bait stations. DO NOT place bait in areas where there is a possibility of

contaminating food or surfaces that may come in contact with food.

Bait stations (**tier 3**) used for the placement of rodenticide bait indoors, in locations not accessible to pets or livestock must have the following characteristics: (1) be constructed of high-strength material (e.g., metal or injection moulded plastic) and resistant to destruction by children; (2) have an entrance designed so that children cannot reach the bait; (3) have an internal structure that prevents bait from being shaken loose; (4) have an access panel that fastens securely and locks (e.g., metal screw or padlock); and (5) bear the product name, active ingredient, guarantee, registration number, "WARNING POISON", and the skull and crossbones symbol.

Bait stations (**tier 2**) used for the placement of rodenticide bait indoors, in locations accessible to pets or livestock must have the following characteristics, in addition to those outlined above for tier 3 bait stations: (1) resistant to destruction by non-target animals; and (2) have an entrance designed so that non-target animals cannot reach the bait.

Bait stations (**tier 1**) used outdoors, above-ground, in locations accessible to children, pets and non-target wildlife must have the following characteristic, in addition to those outlined above for tier 2 and 3 bait stations: (1) be resistant to destruction or weakening by elements of typical non-catastrophic weather (such as, snow, rain, extremes of temperature and humidity, direct sunshine, etc.).

Areas of use:

Food Processing, Food Manufacturing, Food Storage, and Food

Service Areas: For areas not directly related to food processing: Use only in non-food or non-feed area where feed, food, packaging, and handling equipment are never opened or exposed. For areas where feed or food is processed, served, or stored: In meat and food processing plants (processing areas), use only when plant is not in operation. Remove or cover all food, packaging material, and utensils before placing bait in baiting stations. Remove all baits and dead rodents before reuse of the plant (processing areas include storage and service). **Selection of Treatment Areas:** Place bait in locations where rats and mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in and around burrows, and in concealed locations where rats and mice feed is desirable. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly.

Application Directions:

*For Norway Rats (*Rattus norvegicus*) and Roof Rats (*Rattus rattus*):* Norway rats usually inhabit the ground floor of structures and burrow in the ground. They will range approximately 15 to 100 m from their nesting area seeking food and water.

Roof Rats, however, are excellent climbers and usually infest the upper stories of structures. Hence, special attention must be paid to this species ability to climb by placing baits in the roof rats activity area. For both species, eliminate as much of the naturally occurring food as possible and place 100 to 450 grams of bait (usually at intervals of 5-10 m) within the treatment area. Maintain an uninterrupted supply of fresh bait for at least 10 days, or until signs of rat activity cease. *For House Mice (Mus musculus):* House mice are excellent climbers and will infest all floors in a structure. Mice will only range approximately 10 m from their nesting area seeking food. Eliminate as much of the naturally occurring food as possible and place 15 to 30 grams of bait at intervals of 2-4 m within the treatment area. If a bait placement is not disturbed after several days, move the placement approximately 1 metre. Sixty (60) grams of bait per placement may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease. **Note:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent re-infestation, eliminate food, water, and harbourage as much as possible. If re-infestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as necessary. Users should remove clothing immediately if pesticide gets inside. Then wash skin thoroughly and put on clean clothing.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS, AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Chemical-resistant gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait, and empty containers. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** All handlers must wear long sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves when handling this product. In addition, persons loading pellets or bait into mechanical ground equipment, or persons loading/applying with hand-pushed or hand-held equipment, must wear a NIOSH approved dust/mist filtering respirator and protective eyewear.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds.

FIRST AID INSTRUCTIONS:

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **If swallowed,** call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a

physician or poison control centre. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL: Store bait in original, tightly sealed container. Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals. Do not contaminate water, food, or feed by storage or disposal. Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, and the clean-up of spills, contact the Provincial Regulatory Agency or the Manufacturer.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

For medical or environmental emergencies, call CHEMTREC® at 1-800-424-9300.

To reorder, call TOLL FREE: 1-800-387-1784

NOT INTENDED FOR HOUSEHOLD USE OR RESALE

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.