

2017-4707
2017-09-07



Dow AgroSciences

Prestige™ B Herbicide

(A COMPONENT OF PRESTIGE HERBICIDE)

GROUP	4	HERBICIDE
-------	---	-----------

FOR SALE FOR USE ONLY IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA

This component must only be used in the Prestige herbicide.

AGRICULTURAL

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

GUARANTEE: Clopyralid, present as acid 50 g a.e./L
MCPA, present as 2-ethylhexyl ester 280 g a.e./L
Emulsifiable concentrate

REGISTRATION NO. 25464 PEST CONTROL PRODUCTS ACT

CAUTION  **POISON**

EYE IRRITANT

NET CONTENTS: 8.0 L

Dow AgroSciences Canada Inc.
2400, 215-2nd Street S.W.
Calgary, Alberta
T2P 1M4
1-800-667-3852

®™Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

2017-4607

2017-09-07

PRECAUTIONS

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN

EYE IRRITANT

DO NOT APPLY BY AIR

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields and public buildings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

KEEP OUT OF REACH OF CHILDREN.

Do not take internally. Handle with care and mix only in a closed container. Use only in a well ventilated area.

May cause irritation of eyes and skin. Avoid inhalation of vapour or spray mist and contact with eyes, skin or clothing.

Wash thoroughly after handling and before eating, drinking, smoking or using the bathroom. Take a shower immediately after work.

If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.

Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

Do not apply where run-off or erosion is likely to occur.

Do not enter or allow workers entry into treated areas for 12 hours following application to all crops.

PROTECTIVE CLOTHING AND EQUIPMENT

All Application Equipment

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame. Do not cut container.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

2017-4607
2017-09-07

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low blood pressure and unconsciousness.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, fertilizers, drugs or clothing.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

Surface runoff

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid applying this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip (buffer zone) between the treated area and the edge of the water body.

Leaching

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

STORAGE

1. Store in original, tightly-closed container and do not allow water to be introduced into this container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool (above 5°C) dry, locked, well-ventilated area without floor drain. If product is frozen, bring to room temperature and agitate before use.
4. Herbicides should be shipped or stored separate from other pesticides to avoid cross-contamination.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

2017-4607
2017-09-07

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

112912

2017-4607
2017-09-07**Dow AgroSciences**

Prestige™

Herbicide

GROUP	4	HERBICIDE
-------	----------	-----------

FOR SALE FOR USE ONLY IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA

Prestige Herbicide provides control of a wide spectrum of broadleaved weeds including hard-to-control weeds such as cleavers, kochia, wild buckwheat, perennial sow thistle and Canada thistle, and suppression of hemp-nettle and common chickweed, in spring wheat, durum wheat and spring barley.

AGRICULTURAL

Emulsifiable concentrate

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

This co-package contains

Prestige A Herbicide

REGISTRATION NO. 25465 PEST CONTROL PRODUCTS ACT

GUARANTEE: fluroxypyr, present as the 1-methylheptyl ester 180 g/L

Prestige B Herbicide

REGISTRATION NO. 25464 PEST CONTROL PRODUCTS ACT

GUARANTEE: Clopyralid, present as acid 50 g a.e./L
MCPA, present as 2-ethylhexyl ester 280 g a.e./L

DANGER  **POISON**

WARNING - EYE IRRITANT

CAUTION - SKIN IRRITANT

NET CONTENTS: Prestige A Herbicide 6.4 L
Prestige B Herbicide 2 x 8.0 L

Dow AgroSciences Canada Inc.

2400, 215-2nd Street S.W.

Calgary, Alberta

T2P 1M4

1-800-667-3852

®™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

2017-4607
2017-09-07

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DANGER - CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. DO NOT GET IN EYES.

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN

CAUSES SKIN IRRITATION

DO NOT APPLY BY AIR

Prestige B Herbicide contains an ingredient that is rapidly absorbed through the skin and respiratory tract.

DO NOT get in eyes. Avoid contact with skin and clothing. Avoid breathing spray mist or vapour.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields and public buildings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

PROTECTIVE CLOTHING AND EQUIPMENT

All Application Equipment

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.

Do not enter or allow workers entry into treated areas for 12 hours following application to all crops.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame. Use only in a well ventilated area.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

2017-4607
2017-09-07

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low blood pressure and unconsciousness.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, fertilizers, drugs or clothing.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

Surface runoff

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid applying this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip (buffer zone) between the treated area and the edge of the water body.

Leaching

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

STORAGE

Store in original tightly closed containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate food, feedstuffs or domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

2017-4607
2017-09-07

GENERAL INFORMATION

Prestige Herbicide is a selective postemergence herbicide for the control of broadleaved weeds in spring wheat, durum wheat, and spring barley (not underseeded to legumes). Prestige Herbicide provides control of hard-to-kill broadleaved weeds such as cleavers, kochia, wild buckwheat, Canada thistle and perennial sow thistle, and suppression of common chickweed and hemp-nettle. The components of Prestige Herbicide move within the plant to control exposed and underground plant tissues. It mimics naturally occurring plant hormones which control weeds by disrupting normal plant growth patterns. Symptoms of effect include epinasty (twisting of the stems) and swollen nodes.

DIRECTIONS FOR USE

GENERAL USE PRECAUTIONS

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE.
DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.

Do not use in greenhouses.

Apply only when the potential for drift to areas of human habitation or areas of human activity (such as houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment sprayer settings.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Extreme growing conditions such as

- Drought or near freezing temperatures prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.
 - Wet foliage at the time of application may result in reduced weed control.
 - Over-application may result in crop injury.
 - Make only 1 application per year.
 - Do not apply through any type of irrigation system.
- **DO NOT APPLY BY AIR. GROUND EQUIPMENT ONLY.**

Sprayer Clean-up

Sprayers used to apply Prestige Herbicide should not be used to apply other pesticides to sensitive crops without a thorough cleaning. Do not contaminate irrigation or drinking water, lakes, streams or ponds when cleaning the sprayer. Spray equipment should be cleaned after use by the following procedure:

1. Drain the sprayer tank.
2. Hose down the interior surfaces of the tank. Flush tank, hoses, boom and nozzles with clean water for 10 minutes. Fill the tank with water and recirculate for 15 minutes. Spray part of the mixture through the hoses, boom and nozzles and drain the tank.
3. Remove the nozzles and screens and clean them separately.
4. If the spray equipment will be used on other crops other than cereal crops, repeat 1 and 2 again and thoroughly wash the spray mixture from the outside of the spray tank and the boom.

2017-4607
2017-09-07

SENSITIVE PLANTS

Do not apply Prestige Herbicide directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers, fruit, and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

ROTATIONAL CROP RESTRICTIONS

Fields previously treated with Prestige herbicide can be seeded to wheat, barley, oats and rye (not underseeded to forage legumes, clover or alfalfa), canola, field peas, flax, mustard, or can be summer-fallowed. Seed only those crops listed above in the year following treatment. Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application, delay seeding field peas an additional 12 months (total 22 months following application). Contact your local Dow AgroSciences Canada representative or retailer for more information before seeding field peas following drought conditions in the previous year.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Dow AgroSciences Canada Inc at 1-800-667-3852 or www.dowagro.ca for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Dow AgroSciences Canada Inc.

DIRECTIONS FOR USE

Prestige Herbicide Alone

Prestige Herbicide will provide wide spectrum broadleaved weed control in spring wheat, durum wheat, and spring barley.

Grazing Restriction

- Do not permit any grazing within 7 days after application.
- Do not harvest forage or cut hay within 7 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest the treated crop within 60 days after application.

Restricted Entry Interval (REI)

- All crops – 12-hour REI

DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.

2017-4607
2017-09-07

I. Low Rate: To control the following weeds apply a uniform spray containing 0.6 L/ha of Prestige A Herbicide and 1.5 L/ha of Prestige B Herbicide in a minimum of 100 L of water to thoroughly cover the weeds.

Weeds Controlled:

cleavers (1-4 whorls)	lamb's-quarters
Canada thistle (low infestations)	mustard, wild
flax, volunteer (1-12 cm)	shepherd's purse
flixweed (spring seedling 2-4 lf)	stinkweed
kochia ***	sunflower, volunteer

Weeds Suppressed:

stork's-bill (1-8 leaf)

II. High Rate: To control the above weeds plus the following weeds apply a uniform spray containing 0.8 L/ha of Prestige A Herbicide and 2.0 L/ha of Prestige B Herbicide in a minimum of 100 L of water to thoroughly cover the weeds.

Weeds Controlled:

buckwheat, tartary	pigweed, red-root
buckwheat, wild (1-4 leaf)	pigweed, Russian
Canada thistle * (medium to high infestations)	scentless chamomile
canola, volunteer	smartweed
dandelion **	sowthistle, annual
groundsel, common	sowthistle, perennial *
mallow, round-leaved (1-6 leaf)	

Weeds Suppressed:

common chickweed ***
hemp-nettle (2-6 leaf stage)

*Season long control, with some regrowth in the fall (top growth control).

**Spring rosettes only.

***Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

The active ingredients in this tank mixture will only control chickweed that is emerged at time of application. Chickweed plants which emerge after application will not be controlled. To improve the reduction in chickweed population at the end of season, delay the timing of application as late as possible to when the majority of chickweed plants have emerged. The reduction in chickweed population will also be improved in crops which are more competitive and which allow limited light penetration.

NOTE: Prestige Herbicide activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

Timing of Application

Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, and spring barley are in the 3-leaf to just before the flag leaf emergence stage. Do not apply later than the flag-leaf emergence stage. A maximum of 1 application per year.

2017-4607
2017-09-07

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required number of jugs of Prestige A Herbicide.
4. Add the required number of jugs of Prestige B Herbicide.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g., canola and legumes).
7. Follow sprayer clean-up directions as listed.

Spray Application

Apply Prestige Herbicide in sufficient spray volume to provide uniform coverage using properly calibrated ground equipment. A total spray volume of 100 L/ha using low pressure (200 to 275 kPa) application is recommended. Maintain sufficient agitation in the spray tank during mixing and spraying to ensure a uniform spray mixture.

TANK MIXES WITH PRESTIGE HERBICIDE FOR ANNUAL GRASS CONTROL

Tank mixtures of Prestige Herbicide with other herbicides will provide control of additional weeds. Read all precautions, minimal interval to harvest and directions for use on the Prestige Herbicide and tank-mix partner labels, and follow the more stringent of the two.

Herbicide Tank-mix Partner	Crops Registered	Rate of Herbicide Tank-mix Partner	Adjuvant Rate	Additional Weeds
Achieve Liquid	spring wheat, durum wheat, spring barley	0.5 L/ha	Turbocharge 0.5 L/100 L of spray volume (0.5% v/v)	green foxtail wild oats
Horizon Herbicide Tank Mix	spring wheat, durum wheat	230 mL/ha	Score 0.8 L/ha (0.8% v/v)	green foxtail wild oats
Assert 300 SC	spring wheat, durum wheat, spring barley	1.3 L/ha 1.6 L/ha	Refer to Assert 300 SC label	wild oats (1-3 leaf) wild oats (4 leaf)
Everest Solupak 70 DF	spring wheat, durum wheat	43 g/ha	Refer to Everest Solupak 70 DF label	green foxtail wild oats
Puma 120 Super	spring wheat, durum wheat, spring barley	385 mL/ha 770 mL/ha	None required	green foxtail only green foxtail, wild oats & barnyard grass

Tank mixes with ACHIEVE herbicides may cause temporary injury if applied before the 4-leaf stage, however, yield will not normally be affected.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. Boom height must be 60 cm or less above the crop or ground. Do not spray in dead calm near sensitive plants. The "cloud" of suspended droplets may drift onto sensitive plants when the wind comes up. Spray only when wind is blowing away from a sensitive crop, shelterbelt or garden. Spray when wind is less than 8 km/h.

2017-4607
 2017-09-07
 DO NOT apply by air.

Buffer Zones

Use of the following spray methods or equipment DO NOT require a buffer zone: handheld or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind hedge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Spring wheat, durum wheat, winter wheat, spring barley, seedling and established tall fescue grown for seed, seedling and established forage grasses grown for seed, canaryseed and foxtail millet	1	0	1	0	1

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) [Fine or Medium or Coarse] classification.' To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS: The DIRECTIONS FOR USE for this product for the use(s) described below were developed by persons other than Dow AgroSciences Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Dow AgroSciences Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop(s) listed below.

Accordingly, the Buyer and User assumes all risks related to performance and crop tolerance arising, and agrees to hold Dow AgroSciences Canada Inc. harmless from any claims based on efficacy and/or phytotoxicity in connection with the use(s) described below.

DIRECTIONS FOR USE

2017-4607
2017-09-07

SEEDLING AND ESTABLISHED TALL FESCUE GROWN FOR SEED: For control of labelled weeds on seedling (4 leaf to flag leaf) or established tall fescue grown for seed. Apply one post-emergent application when weeds are at the 2-4 leaf stage. Ground application only. Do not cut treated fields for hay/forage. Do not graze treated fields.

Application rates (Apply as a tank-mix only): Apply Prestige A Herbicide at a rate of 0.8 L/ha. Apply Prestige B Herbicide at a rate of 2 L/ha.

Apply in a minimum spray volume of 100 L water/ha.

FOR CONTROL OF LABELLED WEEDS IN THE FOLLOWING SEEDLING AND ESTABLISHED FORAGE GRASSES GROWN FOR SEED IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY:

Creeping Red Fescue
Intermediate Wheat Grass
Crested Wheat Grass
Meadow Brome Grass
Smooth Brome Grass
Timothy

Apply one application per year in a spray volume of 100-200 L/ha to control labelled weeds in seedling (4 leaf to flag leaf) or established forage grasses. Application should be made postemergence when weeds are in the 2-4 leaf stage. Apply in a broadcast spray using a hydraulic boom ground sprayer.

Application rates (Apply as a tank-mix only): Prestige A Herbicide: 0.8 L/ha; Prestige B Herbicide: 2.0 L/ha.

- Do not cut treated fields for hay/forage. Do not graze treated fields. Do not harvest the treated crop within 60 days of application.

LABELLED ANNUAL AND PERENNIAL BROADLEAVED WEEDS IN CANARYSEED (*Phalaris canariensis*)

For the control of labelled annual and perennial broadleaved weeds in canaryseed, apply Prestige A Herbicide at a rate of 0.8 L/ha and Prestige B Herbicide at a rate of 2.0 L/ha in a minimum of 100 L of water to thoroughly cover the weeds with a spray pressure of 200 to 275 kPa. Apply when weeds are in the seedling stage (2-4 leaf) and when canaryseed is in the 3-leaf stage up to just before the flag leaf emergence stage. Apply using a ground boom sprayer, and apply only one post-emergent application per season.

DO NOT apply within 60 days of harvest. For use in the Prairie Provinces and the Peace River regions ONLY of British Columbia.

A maximum of 2 applications are permitted per season with a minimum retreatment interval of 21 days.

FOXTAIL MILLET:

For the control of broadleaf weeds and Canada thistle, apply Prestige A Herbicide at the rate of 0.6 to 0.8 litres per hectare plus Prestige B Herbicide at the rate of 1.5 to 2.0 litres per hectare. Use the higher rate on moderate to heavy populations of Canada thistle and to expand the number of broadleaf weeds controlled (see main label for list of weeds). Apply when Canada thistle is 10 to 15 centimetres high and when foxtail millet is in the 2- to 5-leaf stage. Apply in 100 litres of water per hectare.

Do not permit lactating dairy animals to graze fields within 7 days after application.

Do not harvest forage or cut hay within 60 days after application.

Withdraw meat animals from treated field at least 3 days before slaughter.

Refer to the main Prestige Herbicide label for additional details and instructions before using.

2017-4607

2017-09-07

READ THE ROTATIONAL CROPPING RESTRICTIONS ON FULL LABEL BEFORE USING THIS PRODUCT

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Prestige Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Prestige Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Prestige Herbicide or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Dow AgroSciences Canada Inc. at 1-800-667-3852 or at www.dowagro.ca.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

All products listed are registered trademarks of their respective companies.

112912