

2018-6400
2019-01-17

{ booklet label }

GROUP

3

FUNGICIDE

JADE™ Fungicide

COMMERCIAL

Emulsifiable Concentrate for disease control in Peaches, Nectarines, Plums, Sweet Cherries, Sour Cherries, Apricots, Highbush and Lowbush Blueberries, Saskatoon Berries, Cranberries, Caneberries, Rutabagas, Asparagus, Western Red Cedar, Kentucky Bluegrass Grown for Seed and Strawberries.

GUARANTEE:

Propiconazole

250 g/L

READ THE LABEL AND PAMPHLET BEFORE USING

REGISTRATION NO.: 24030
PEST CONTROL PRODUCTS ACT



WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

Net Contents: 2L

Belchim Crop Protection Canada Inc.
104 Cooper Drive, Unit 3
Guelph ON N1C 0A4
1-866-613-3336

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the **PEST CONTROL PRODUCTS ACT** to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

In case of contact with eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. In case of contact with skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

There is no specific antidote for this product. Apply symptomatic therapy. This product contains petroleum distillates.

STORAGE

To prevent contamination store this product away from food or feed.

PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin-sensitization reactions in certain individuals.

Wear overalls over long pants, a long sleeve shirt, shoes and socks and chemical-resistant gloves during mixing/loading, application, clean-up and repair activities.

Wear protective goggles or faceshield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used.

Do not eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes. After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse.

Do not contaminate food or feed.

NOTE: Do not graze livestock on treated green crops.

This pesticide is toxic to fish. Do not spray any body of water by direct application, drift or by cleaning and rinsing spray equipment.

This product contains a petroleum distillate that is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

DO NOT allow entry into treated area for 12 hours following application. See the **DIRECTIONS FOR USE** section for crop specific restricted entry intervals.

CONTAINER DISPOSAL:

For recyclable container for commercial use

DO NOT REUSE THIS CONTAINER FOR ANY PURPOSE. This is a recyclable container and is to be recycled at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Dispose of the rinsings in accordance with provincial requirements
2. Make the empty container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, or in the case of a spill or spill clean-up, contact the manufacturer or the provincial regulatory agency.

For returnable containers commercial use

DO NOT reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

For information on disposal of unused, unwanted product, or in the case of a spill or spill clean-up, contact the manufacturer or provincial regulatory agency

ENVIRONMENTAL HAZARDS:

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

Toxic to aquatic organisms and non-target terrestrial plants

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application sites such as hedgerows and woodland.

***IN CASE OF EMERGENCY INVOLVING A POISONING OR MAJOR SPILL,
CALL 1-866-336-2983***

RESISTANCE MANAGEMENT :

For resistance management, please note that JADE FUNGICIDE contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to JADE FUNGICIDE and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed. To delay fungicide resistance:

2018-6400
2019-01-17

- Where possible, rotate the use of JADE FUNGICIDE or other Group 3 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group when such use is permitted.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact you local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact Belchim Crop Protection Canada Inc. at 1-866-613-3336 or at www.belchimcanada.com.

GENERAL INFORMATION

JADE FUNGICIDE is a broad-spectrum systemic fungicide for control of a wide range of diseases on certain crops and will protect the crop from yield and quality losses due to disease. JADE FUNGICIDE may be used in conjunction with higher seeding rates, higher fertilizer inputs, plant growth regulators and other fungicides as required. (See "NOTE" under Precautions)

2018-6400
2019-01-17

Diseases Controlled:

Cranberries:	Cottonball (<i>Monilinia oxycocci</i>)
Kentucky Bluegrass Grown for Seed:	Powdery Mildew
Lowbush Blueberries:	Monilinia Blight (Mummyberry)
Highbush Blueberries:	Mummyberry (<i>Monilinia vaccinii – corymbosi</i>)
Peaches, Nectarines, Plums, Apricots:	Brown Rot Blossom Blight, Fruit Brown Rot.
Sweet and Sour Cherries:	Brown Rot Blossom Blight, Fruit Brown Rot, Cherry Leaf Spot (<i>Blumeriella jaapii</i>)
Plums and Sour Cherries:	Black Knot (<i>Apiosporina morbosa</i>) [suppression only]
Caneberries (Crop group 13A; cultivars and/or varieties of red and black raspberry, logan berry and blackberry):	Yellow Rust
Rutabagas:	Powdery Mildew
Asparagus:	Rust (<i>Puccinia asparagi</i>)
Saskatoon Berry	Entomosporium Leaf and Berry Spot, Saskatoon Juniper Rust
Western Red Cedar	Keithia Foliar Blight
Strawberries	Leaf Spot (<i>Mycosphaerella fragariae</i>)

Factors Affecting JADE FUNGICIDE Performance:

JADE FUNGICIDE should be applied as a preventative disease control measure. Established diseases are more difficult to control and may have already reduced crop vigour.

If rainfall occurs within one hour of application, reapplication is necessary.

DIRECTIONS FOR USE:

DO NOT use in greenhouses

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

FIELD SPRAYER APPLICATION

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

AIRBLAST APPLICATION

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

AERIAL APPLICATION

Apply only by fixed-wing or rotary aircraft equipment that has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial applications on this label. If directions for aerial application do not appear for a specific use, then aerial equipment cannot be used to apply this product for that purpose.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precaution

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide*

Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 m/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor span.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals in a closed system is permitted.

It is desirable that the pilot has communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls over long pants and long sleeve shirt, shoes and socks and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have any questions, call the manufacturer at 1-866-613-3336 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

1. Apply the recommended rate in a minimum spray volume of 40 litres per hectare.
2. Wind speed during application should be less than 16 km/h to avoid drift.
3. High humidity and low temperatures (10-20°C) allow for a better deposition of spray droplets.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				Terrestrial habitat
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		
			Less than 1 m	Greater than 1m	Less than 1 m	Greater than 1 m	
Field sprayer*	Rutabagas, asparagus, Kentucky bluegrass (seed prod.), Western cedar		1	0	1	1	1
Airblast	Cherries	Early growth stage	5	0	10	3	10

		Late growth stage	2	0	4	2	4
	Blueberries, apricots, nectarines, peaches, plums, Saskatoon berries, blackberries, loganberries, raspberries, other berries	Early growth stage	4	0	5	2	5
		Late growth stage	2	0	3	1	3
Aerial	Blueberries, Kentucky bluegrass (seed prod.)	Fixed wing	1	0	3	1	20
		Rotary wing	1	0	1	1	20

* For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

SPRAYER AND APPLICATION INFORMATION

The performance of this product depends on correct application. Follow the guidelines given below for optimal application of JADE FUNGICIDE.

	GROUND APPLICATION	AERIAL APPLICATION
- Spray Volume:	Minimum 200 litres of water per hectare	40-50 litres of water per hectare
- Spray Pressure:	200-300 kPa	100-200 kPa
- Nozzle Type:	110° Flat Fan (XR11004, 4110-20)	Flat Fan 6510-6515 or Hollow Cone (D8-45)
- Droplet Size:	Medium Spray (300-400 microns VMD)	Medium Spray (350-400 microns VMD)
- Ground Speed:	10 km/h	
- Nozzle Angle:	90° (straight down)	90° (straight down)
- Boom Height:	40-50 cm above the crop canopy	2-3 m above the crop canopy

Ground Application - Mixing and Spraying Instructions

1. Spray equipment should be thoroughly flushed with clean water before mixing JADE FUNGICIDE.
2. Fill spray tank 1/2 full with clean water. Engage gentle agitation.
3. Add the required amount of JADE FUNGICIDE and agitate thoroughly.
4. Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tankmix partner.
5. Complete filling the spray tank with water, maintaining agitation during mixing and spraying operations.
6. Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Aerial Application - Mixing and Spraying Instructions

1. Spray equipment should be thoroughly flushed with clean water before mixing JADE FUNGICIDE.
2. Fill premix tank 1/2 full with clean water. Engage gentle agitation.
3. Add the required amount of JADE FUNGICIDE and agitate thoroughly.
4. Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tankmix partner.
5. Complete filling the premix tank with water.
6. Maintain gentle agitation during mixing.
7. Transfer the premix contents into the aircraft spray tank.
8. Maintain sufficient agitation during the mixing and spraying operation to ensure JADE FUNGICIDE remains in suspension.

9. Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Fertilizer - Mixing and Spraying Instructions

If desired, small amounts of nitrogen may be applied with JADE FUNGICIDE. The appropriate amount of urea can be dissolved in water and added to the spray tank before adding JADE FUNGICIDE. The rate of actual nitrogen must not exceed 10 kg/ha.

CAUTION: Excessive nitrogen concentrations may injure the crop.

NOTE: DO NOT add nitrogen when tank-mixing JADE FUNGICIDE with an herbicide.

NOTE: Do not graze livestock on treated crop.

TANKMIXES

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

PEACHES, NECTARINES, PLUMS, AND APRICOTS - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Brown Rot Blossom Blight.	500 mL	Apply 500 mL of JADE FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1 st application at early bloom with a 2 nd application at 50% - 75% bloom. If disease conditions persist, make a 3 rd application at petal fall.
Fruit Brown Rot	500 mL	Apply no more than 2 applications in the 3 weeks prior to harvest. Apply 500 mL of product in a minimum of 500 L of water per hectare by ground.
Suppression of Black Knot (Plums only)	500 mL	Apply 500 mL of JADE FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1 st application at early bloom with a 2 nd application at 50% - 75% bloom. If disease conditions persist, make a 3 rd application at petal fall.
DO NOT APPLY WITHIN 3 DAYS OF HARVEST. DO NOT REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION. IF REQUIRED, INDIVIDUALS MAY REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION FOR SHORT TERM TASKS WHICH DO NOT REQUIRE HAND LABOUR IF AT LEAST 4 HOURS HAVE PASSED SINCE APPLICATION AND PROVIDED LONG PANTS, LONG-SLEEVED SHIRTS, AND CHEMICAL-RESISTANT GLOVES ARE WORN.		

SWEET AND SOUR CHERRIES - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Brown Rot Blossom Blight.	500 mL	Apply 500 mL of JADE FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1 st application at early bloom with a 2 nd application at 50% - 75% bloom. If disease conditions persist, make a 3 rd application at petal fall.
Fruit Brown Rot	500 mL	Apply no more than 2 applications in the 3 weeks prior to harvest. Apply 500 mL of product in a minimum of 500 L of water per hectare by ground.
Cherry Leaf Spot (<i>Blumeriella jaapii</i>)	500 mL	Apply a maximum of 3 applications per season for control of Cherry Leaf Spot. Make the 1 st application at petal fall. In the 3 weeks prior to harvest make a 2 nd and 3 rd application at a 7-10 day interval. Do not apply within 3 days of harvest. Apply 500 mL of product in a minimum of 500 L of water per hectare by ground.
Suppression of Black Knot (sour cherries only)	500 mL	Apply 500 mL of JADE FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1 st application at early bloom with a 2 nd application at 50% - 75% bloom. If disease conditions persist, make a 3 rd application at petal fall.
It is recommended that no more than 2 consecutive applications of JADE FUNGICIDE be made before switching to another fungicide with a different mode of action according to disease management practices.		
APPLY A MAXIMUM OF 5 APPLICATIONS PER SEASON OF JADE FUNGICIDE TO SWEET AND SOUR		

CHERRIES. DO NOT APPLY WITHIN 3 DAYS OF HARVEST. DO NOT REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION. IF REQUIRED, INDIVIDUALS MAY REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION FOR SHORT TERM TASKS WHICH DO NOT REQUIRE HAND LABOUR IF AT LEAST 4 HOURS HAS PASSED SINCE APPLICATION AND PROVIDED LONG PANTS, LONG-SLEEVED SHIRTS, AND CHEMICAL-RESISTANT GLOVES ARE WORN.

Highbush Blueberry - Spray Schedule		
DISEASE	RATE/HA	REMARKS
Mummyberry (<i>Monilinia vaccinii-corymbosi</i>)	500 mL	Apply first application at or near flower bud swelling; make a second application at leaf bud swelling. Apply by ground only, making no more than two applications per year. In BC only, apply by ground, a third application at pink bloom and a fourth application 7-10 days later at early bloom, making no more than 4 applications per year. Use a minimum of 200 L of water per hectare.
<p>LAST APPLICATION MUST BE MADE PRIOR TO 60 DAYS BEFORE HARVEST. A restricted entry interval of 5 days is required for workers hand pruning highbush blueberries.</p>		

Lowbush Blueberry - Spray Schedule		
DISEASE	RATE/HA	REMARKS
Monilinia Blight	500 mL	Apply first application when flower bud scales first appear and make a second application 10 days later. Use ground application or aerial application equipment, making no more than two applications per year. Use a minimum of 200 L of water per hectare if applying by ground equipment; use 40-50 L of water per hectare if applying by air.
<p>LAST APPLICATION MUST BE MADE PRIOR TO 60 DAYS BEFORE HARVEST.</p>		

Saskatoon Berry - Spray Schedule		
DISEASE	RATE/HA	REMARKS
<i>Entomosporium</i> Leaf and Berry Spot; Saskatoon Juniper Rust.	500 mL	As a foliar spray, apply up to 3 applications per season. The first application to occur at “white tip”, the second application at “petal fall”, and the third application at “green fruit”. Apply 500 mL in a minimum of 200 L of water per hectare by ground, applying to runoff.
<p>LAST APPLICATION MUST BE MADE PRIOR TO 38 DAYS BEFORE HARVEST.</p>		

Cranberry - Spray Schedule		
DISEASE	RATE/HA	REMARKS
Cottonball (<i>Monilinia oxycocci</i>)	500 mL	Apply the first application at leaf bud break. Make a second application 10-14 days later, a third application at early bloom and a fourth application 10-14 days after the third application. Make no more than 4 applications per year. Apply product by ground.
<p>LAST APPLICATION MUST BE MADE PRIOR TO 45 DAYS BEFORE HARVEST.</p>		

CANEBERRIES (CROP GROUP 13A; CULTIVARS AND/OR VARIETIES OF RED AND BLACK RASPBERRY, LOGAN BERRY AND BLACKBERRY) - SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Yellow rust	500 mL	Apply 500 mL of JADE FUNGICIDE in a minimum of 500 L of water per hectare by ground application at first detection of disease in the field and a second application 14 days later.
It is recommended that no more than 2 consecutive applications of JADE FUNGICIDE be made before switching to another fungicide with a different mode of action according to disease management practices.		
APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF JADE FUNGICIDE TO CANEBERRIES. DO NOT APPLY WITHIN 30 DAYS OF HARVEST. DO NOT REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION. IF REQUIRED, INDIVIDUALS MAY REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION FOR SHORT TERM TASKS WHICH DO NOT REQUIRE HAND LABOUR IF AT LEAST 4 HOURS HAS PASSED SINCE APPLICATION AND PROVIDED LONG PANTS, LONG SLEEVED SHIRTS, AND CHEMICAL RESISTANT GLOVES ARE WORN.		

RUTABAGAS - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Powdery Mildew	400 mL	Make two applications per season with the first application at 50 days after planting and the second application 20 days later. Apply to vegetative foliage. Apply 400 mL in a minimum 200 L of water per hectare by ground.
LAST APPLICATION MUST BE MADE PRIOR TO 21 DAYS BEFORE HARVEST.		

ASPARAGUS- SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Rust (<i>Puccinia asparagi</i>)	250 mL	Apply JADE FUNGICIDE to asparagus ferns in Ontario and Quebec only. Once harvest is complete, make the first application of JADE FUNGICIDE as soon as fern growth begins, followed by applications at 14 to 21 day intervals. For new, non-harvested plantings, apply JADE FUNGICIDE when first sign of rust is visible, followed by applications at 14 to 21 day intervals. Apply by ground only, making no more than three applications per year. Use a minimum of 370 L of water per hectare.
LAST APPLICATION MUST BE MADE PRIOR TO 8 MONTHS BEFORE HARVEST.		

WESTERN RED CEDAR - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Keithia Foliar Blight	500 mL	Apply using ground application equipment every four weeks. Make a maximum of 6 applications per year. Apply 500 mL of JADE FUNGICIDE in a volume of 1000 L of water per hectare by ground.

KENTUCKY BLUEGRASS FOR SEED PRODUCTION - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Powdery Mildew	500 mL	Apply as a foliar spray. Make no more than 2 applications per crop year with the first application at pre-heading and the second at 50% - 100% heading. Apply in 200-300 L/ha of water by ground or 40-50 L/ha of water by air.

NOTICE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Belchim Crop Protection Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion Program. Belchim Crop Protection Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Belchim Crop Protection Canada Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

STRAWBERRIES - SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Leaf Spot (<i>Mycosphaerella fragariae</i>)	500 mL	Apply 500 mL of JADE FUNGICIDE per hectare by ground in enough water to ensure thorough coverage. Make 1 st application when disease levels are no more than 5%. Apply JADE FUNGICIDE at 10 day intervals for control of leaf spot.

It is recommended that no more than 2 consecutive applications of JADE FUNGICIDE be made before switching to another fungicide with a different mode of action according to disease management practices.

APPLY A MAXIMUM OF 4 APPLICATIONS PER SEASON OF JADE FUNGICIDE TO STRAWBERRIES. DO NOT APPLY WITHIN 1 DAY OF HARVEST.

TM JADE is a trademark of Belchim Crop Protection Canada Inc.

2018-6400
2019-01-17

{ principal display panel }

GROUP	3	FUNGICIDE
-------	----------	-----------

JADE™ Fungicide

COMMERCIAL

Emulsifiable Concentrate for disease control in Peaches, Nectarines, Plums, Sweet Cherries, Sour Cherries, Apricots, Highbush and Lowbush Blueberries, Saskatoon Berries, Cranberries, Caneberries, Rutabagas, Asparagus, Western Red Cedar, Kentucky Bluegrass Grown for Seed and Strawberries.

GUARANTEE:	
Propiconazole	250 g/L

READ THE LABEL AND PAMPHLET BEFORE USING

REGISTRATION NO.: 24030
PEST CONTROL PRODUCTS ACT



WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

Net Contents: 2L

Belchim Crop Protection Canada Inc.
104 Cooper Drive, Unit 3
Guelph ON N1C 0A4
1-866-613-3336

2018-6400
2019-01-17

{ base container label }

GROUP

3

FUNGICIDE

JADE™ Fungicide

COMMERCIAL

GUARANTEE: propiconazole 250 g/L

READ THE LABEL AND PAMPHLET BEFORE USING

REGISTRATION NO.: 24030

PEST CONTROL PRODUCTS ACT



WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

**IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-866-336-2983**

Net Contents: 6 x 2L

Belchim Crop Protection Canada Inc.
104 Cooper Drive, Unit 3
Guelph ON N1C 0A4
1-866-613-3336

™ JADE is a trademark of Belchim Crop Protection Canada Inc.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the **PEST CONTROL PRODUCTS ACT** to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

In case of contact with eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. In case of contact with skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

There is no specific antidote for this product. Apply symptomatic therapy. This product contains petroleum distillates.

STORAGE

To prevent contamination store this product away from food or feed.

PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin-sensitization reactions in certain individuals.

Wear overalls over long pants, a long sleeve shirt, shoes and socks and chemical-resistant gloves during mixing/loading, application, clean-up and repair activities.

Wear protective goggles or faceshield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used.

Do not eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes. After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse.

Do not contaminate food or feed.

NOTE: Do not graze livestock on treated green crops.

This pesticide is toxic to fish. Do not spray any body of water by direct application, drift or by cleaning and rinsing spray equipment.

This product contains a petroleum distillate that is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

DO NOT allow entry into treated area for 12 hours following application. See the **DIRECTIONS FOR USE** section for crop specific restricted entry intervals.

CONTAINER DISPOSAL:

For recyclable container for commercial use

DO NOT REUSE THIS CONTAINER FOR ANY PURPOSE. This is a recyclable container and is to be recycled at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Dispose of the rinsings in accordance with provincial requirements
2. Make the empty container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, or in the case of a spill or spill clean-up, contact the manufacturer or the provincial regulatory agency.

For returnable containers commercial use

DO NOT reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

For information on disposal of unused, unwanted product, or in the case of a spill or spill clean-up, contact the manufacturer or provincial regulatory agency

IN CASE OF EMERGENCY INVOLVING A POISONING OR MAJOR SPILL, CALL 1-866-336-2983