

2014-02-07
2014-0314

Front Panel

**Hartz® UltraGuard®
Flea & Tick Collar for Dogs
Slow-Release Generator**

Kills:

Fleas for 5 Months

Ticks for 4 Months

GUARANTEE :

Tetrachlorvinphos....14.55%

Warning, contains the allergen soy.

DOMESTIC

READ ENTIRE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

FOR USE ONLY ON DOGS

REG. NO.: 25620 PCP ACT

Net Contents: g

HARTZ CANADA INC.,
1125 Talbot Street
St. Thomas, Ontario
N5P 3W7
1-800-275-1414

Back Panel

DIRECTIONS FOR USE:

Do NOT remove the collar from its package until ready to use. Remove the collar from the package, unroll and STRETCH to activate. This releases the active ingredient at time of use and assures full activity. Place the flea collar around the dog's neck, allowing a spacing of 2 fingers between collar and neck and buckle in place. Leave 5 to 8 cm on the collar for extra adjustment. Cut off any excess length, wrap and dispose of it in the garbage. Wash hands with soap and water after handling collar.

When collar is first worn, observe the pet's neck area every few days for irritation or adverse reaction. Do NOT use on dogs under 12 weeks of age. Consult a veterinarian before using on sick, aged, pregnant, or nursing animals or animals receiving drug or other pesticide treatment. Remove collar when bathing pet with soap or detergent. For continuous protection in situations where there is an ongoing risk of pest infestation, replace the collar every five months for fleas, every four months for ticks. Under conditions where dogs are exposed to severe flea or tick infestations it may be necessary to replace the collar more frequently. Replace this collar no more than once every 4 months. **RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.**

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

PRECAUTIONS:

HUMANS: Do not allow children to handle the collar. Do not unroll collar until ready to use. This collar is intended for use only as an insecticide generator dog collar. Dust will form on the collar during storage. Dust may irritate eyes, nose and throat. Harmful if collar is chewed, or if dust is inhaled or absorbed through skin. Avoid contact with mouth, eyes, skin and clothing. Avoid breathing dust from collar. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils, or places where food is prepared or stored.

PETS: Dust may irritate eyes, nose and throat. Harmful if collar is chewed or swallowed. Avoid contact with the dog's mouth, face, eyes and genitals. Avoid letting the pet inhale the dust from the collar.

Do not use this product on dogs at the same time or within 30 days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides, or chemicals.

FIRST AID

HUMANS: IF CHEWED, Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce

vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING**, Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **IF DUST FROM COLLAR GETS IN EYES**, Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **IF INHALED**, Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

PETS: Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If swallowed or chewed, consult a veterinarian.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: This product contains a pesticide that is a cholinesterase inhibitor (anti-cholinesterase compound). Symptoms of human poisoning may include headache, weakness, sweating, blurred vision, nausea and diarrhea. Obtain medical attention or call a poison control centre at once. Atropine is antidotal. Acute symptoms of over dosages in dogs include diarrhea, salivation, vomiting, muscular tremors and weakness. Contact a veterinarian immediately.

STORAGE AND DISPOSAL: Wrap and discard excess collar, unwanted (used) collars and packaging in household garbage. Do not reuse empty pouch.

For more information call our experts at 1-800-275-1414
Visit www.hartzultraguard.com

Hartz® and other trademarks are trademarks of The Hartz Mountain Corporation

RMXXXXXX