

2016-5616
2017-02-27

This product has been cancelled effective as of 24 July 2023.
Manufacturing and importing of the product are no longer permitted. The existing stocks in Canada are authorized under paragraph 21(5)(a) of the Pest Control Products Act for depletion for a limited time according to the phase-out schedule listed in the Pesticide Product Information Database (<https://pesticide-registry.canada.ca>).

AIR GUARD®

**KONKER
COMMERCIAL AND FARM
INSECT KILLER**

**PRESSURIZED SPRAY
COMMERCIAL**

For use on cattle, in animal quarters, milk rooms, dairy barns, horse barns, piggeries, shelter sheds, food processing plants, restaurants, hotels, hospitals, and other commercial and industrial buildings.

Kills and Repels: stable flies, horse flies, face flies, deer flies, house flies, horn flies, mosquitoes, biting midges, wasps, flying moths, and crawling insects such as cockroaches, spiders, crickets, ants, carpet beetles, centipedes, silverfish, dog & cat fleas, brown dog and American dog ticks and bed bugs.

GUARANTEE:

Pyrethrins..... 0.30%
Piperonyl Butoxide.. 2.40%

REGISTRATION NO. 24435
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

DANGER EXTREMELY FLAMMABLE
EXPLOSIVE

NET CONTENTS: 425 g

Acuity Holdings, Inc.
11627 - 178 STREET
EDMONTON AB T5S 1N6
L4K 3X5
1-905-669-9876

DIRECTIONS FOR USE:

SPACE SPRAY: Food processing plants, restaurants, hotels, hospitals, and other commercial and industrial buildings. Remove all personnel, patients or animals before starting to treat area. Direct spray upwards to all parts of the room, especially windows and other light sources. Apply at a rate of 6 to 10 seconds per 30 cubic metres. Keep spray at least 1 metre from interior walls, fabrics and furniture. For best results, close all doors and windows during treatment and keep room closed for 15 minutes before ventilating. Do not remain in treated areas. Ventilate treated areas thoroughly for at least 30 minutes before re-entry. In meat packing and food processing plants and food service establishments, apply only when plant is not in operation. Remove all food before treatment. Cover all food processing surfaces and clean thoroughly with potable water before resuming operations. In hospitals do not use in nurseries or rooms where infants, ill or aged people are confined.

LIVESTOCK SPRAY: To treat beef and dairy cattle (lactating and nonlactating) against flies. Spray animals once daily with fine mist. Apply at a rate of 6 to 8 seconds per animal. Spray should be enough to wet ends of hair only, do not wet the skin. Take care not to spray in animal's eyes or mouth. Treat dairy cattle at least 20 minutes before milking. Wash udders before milking.

MILK PARLORS AND MILKROOM SPACE SPRAY: When used in milk houses, and milk parlours use only when milk house is not in operation. Remove any animals from treatment area before spraying. Cover or remove all milk and milking implements before spraying. After spraying, wash all surfaces that may contact milk and rinse thoroughly with potable water before resuming operations.

CRAWLING INSECTS: Apply to insect hiding places and breeding areas such as baseboards, moist areas, openings around sinks, drains, and pipes, behind cabinets and storage areas. If possible, hit insects directly to kill. Spray ant trails, around door sills, window frames or in ant nest. **BED BUGS:** Spray mattress, particularly around seams and tufts. Take bed apart, spray joints. Repeat as required. **FLEAS:** Spray infested areas such as pet's bed and resting quarters, nearby cracks and crevices, along and behind baseboards, windows and door frames, and localized areas and floor covering where pests may be present. Remove old bedding and replace with fresh clean bedding after treatment. Insects must be contacted to be killed. Do not treat animals with this product.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not remain in treated areas. Ensure proper ventilation after areas have been treated. In hospitals, do not use in nurseries or rooms where infants, ill or aged persons are confined. Cover milk and milking utensils to prevent contamination from spray and dead insects. (Any treated surface that may contact food or feed products should be washed thoroughly with potable water after spraying and before re-use). Do not contaminate feed, food or water troughs. Avoid inhalation of spray. Avoid contact with skin and eyes. Wash after using. **CONTENTS UNDER PRESSURE.** Do not place in hot water or near radiators, stove or other sources of heat. Do not puncture or incinerate can or store at temperatures over 50°C. Do not spray into or near any open flame, spark, or while smoking.

FIRST AID: If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. Take container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.