

IFC and Blister Card

Front Panel

**Hartz® UltraGuard Plus® Flea & Tick Drops
for Dogs and Puppies Weighing over 14KG (30lbs)**

SOLUTION

Kills:

Fleas

Ticks

Plus Kills and Prevents:

Flea Eggs for 30 Days

ACTIVE INGREDIENTS:

Permethrin45%

(S) – Methoprene.....2.9%

DOMESTIC

READ ENTIRE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

FOR USE ONLY ON DOGS

DO NOT USE ON CATS



Toxic to cats

REG. NO. 26490 PCP ACT

Net Contents: 3 Tubes x 2 mL each

Hartz Canada, Inc.
1125 Talbot Street
St. Thomas, Ontario
N5P 3W7
1-800-275-1414

Back Panel

DIRECTIONS FOR USE:

This product contains permethrin and (S)-methoprene. Do not apply another pest control product such as a shampoo, collar, or powder that contains permethrin or (S)-methoprene to the treated animal after applying Hartz® UltraGuard Plus® Flea & Tick Drops for Dogs and Puppies Weighing over 14KG (30lbs).

DO NOT USE ON CATS

1. Remove one applicator tube from package and hold in an upright position.
2. Pull cap off. Turn the cap around and place on top of tube.
3. Press cap down to break the seal of tube.
4. Remove cap carefully. Position the tip of the tube on the dog's back between the shoulder blades and squeeze out half the contents of the tube.
5. Invert the applicator tube upright and go to the dog's tail. Position the tip of the tube at the base of the tail and apply the remaining contents of the tube to the top of the dog's rump near the base of the dog's tail.
6. Repeat every month if flea or tick problems persist.

Do not use this product more often than once every 3 weeks. After application, through its natural movement, your pet's skin and hair oils carry this product over the pet's body for protection against fleas, ticks and flea eggs. Reapply every month for continual pest control in situations where there is an ongoing risk of pest infestation. Do not use on animals under 12 weeks of age. Do NOT administer by mouth. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. Discontinue use if adverse effects are observed. If condition persists, consult your veterinarian. **RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.**

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

PRECAUTIONS:

HUMANS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. May cause eye irritation. Avoid contact with mouth, eyes and clothing. Use rubber gloves when applying product. Avoid contact with pet until dry. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils or places where food is prepared or stored. **PETS:** Toxic to cats. Cats that actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harm, including death. Avoid contact with the dog's mouth, face, eyes and genitals. Sensitivities may occur after using any pesticide for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. Harmful if swallowed.

Side Effects: Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching, changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur, consult your veterinarian or Hartz Canada, Inc. at 1-800-275-1414.

FIRST AID

HUMANS: IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. No specific antidote.

STORAGE AND DISPOSAL: Store in a cool, dry place away from food and out of the reach of children. Do not reuse empty container. Wrap and dispose of empty container with household garbage.

For more information call our experts at 1-800-275-1414
Visit www.hartzultraguard.com

Hartz® and other trademarks are trademarks of The Hartz Mountain Corporation

2020-5496
2020-12-18

TUBE COPY

Hartz Ultraguard Plus Flea & Tick Drops for Dogs and Puppies Over 14 Kg (30 lbs.)

DOMESTIC

Permethrin.....45%
(S)-Methoprene.....2.9%

REG. NO.: 26490 PCP ACT

READ LABEL BEFORE USING
DO NOT USE ON CATS

Net Contents: 2 mL

.
Hartz Canada, Inc.
1125 Talbot Street
St. Thomas, Ontario
N5P 3W7
1-800-275-1414