

2020-08-04  
Sub.No.2020-1550

[FRONT PANEL - ENGLISH]

**FAST-ACTING FORMULA  
RESULTS IN 24 HOURS!**

With FastAct® foam ... see exactly where you've sprayed

**ROUNDUP® READY-TO-USE  
GRASS AND WEED CONTROL  
With FastAct® Foam**

Non-Selective Herbicide

SOLUTION  
WATER-BASED  
[CONTROLS ENTIRE PLANT, INCLUDING THE ROOT]  
KILLS THE ROOTS!  
RAIN-PROOF IN TWO HOURS!  
DOMESTIC

|                   |    |                   |
|-------------------|----|-------------------|
| NET CONTENTS: 1 L | TO | NET CONTENTS: 6 L |
|-------------------|----|-------------------|

GUARANTEE:

Glyphosate ..... 7 g/L  
(present as isopropylamine salt)

REG. NO.: 24299 P.C.P. ACT

Contains 5-Chloro-2-methyl-4-isothiazolin-3-one at 0.0006% and  
2-Methyl-4-isothiazolin-3-one at 0.0002% as preservatives.


or

Contains 1,2-Benzisothiazolin-3-one at 0.0193% as a preservative.

KEEP OUT OF REACH OF CHILDREN  
READ THE LABEL BEFORE USING

Bayer CropScience Inc.  
Suite 200, 160 Quarry Park Blvd. SE  
Calgary, AB  
T2C 3G3  
1-866-476-7645  
Made and printed in U.S.A.

[BACK PANEL - ENGLISH]



|  |
|--|
| <b>FOR USE AND PRECAUTIONS<br/>BEFORE USING.</b> |
|--|

With QUICK CONNECT® sprayer.

**USE AND STORAGE DIRECTIONS FOR THE QUICK CONNECT® SPRAYER:**

1. Pull trigger sprayer from bottle.
2. Pull hose connector and coil from trigger sprayer handle.
3. Insert hose connector into [spout/knob/opening on cap] until it "clicks" in place.
4. [Flip up spout./Flip up spout until fully upright./Twist [spout/knob] to "ON"/Pull spout up.]
5. Adjust spray nozzle to STREAM for cracks and crevices or FOAM, which marks weeds for up to 5 minutes and lets you easily see where you've sprayed.
6. Aim sprayer nozzle at centre of weed and spray to wet.
7. Storage: After use, [close spout on cap./flip spout down to the "flat" position to close./twist [spout/knob] to "OFF"/push spout down.] No need to disconnect trigger sprayer. Close nozzle on trigger sprayer. Snap trigger sprayer back into place on bottle.

**NO RESIDUAL ACTIVITY IN SOIL**

ROUNDUP®, a non-selective vegetation control herbicide, controls most annual and perennial grasses, including lawn grasses, broadleaf weeds and brush (i.e. virtually anything that is green and growing). A systemic herbicide, it is absorbed by the leaves and moves throughout stem and roots to control the entire plant. Visible symptoms of control, such as browning or yellowing of leaves, will occur within 1-2 days. For best results, to ensure that Roundup® gets down to the roots, avoid tillage or disturbing the treated area for 7-10 days (or as otherwise specified on the label). This will ensure that the entire plant, including the root, is eliminated. Rain or watering two hours after application will not wash away effectiveness.

**NOTE:** Roundup® has no soil activity and will not leach or run off to affect nearby vegetation. Plants must be present to be controlled. Repeat applications will be required as new seedlings and vegetation emerge.

**USES:** Lawn edging/trimming, patio weed control, lawn/garden renovation, hard to mow areas.

Product Use Information: 1-866-476-7645  
Website: [www.roundupcanada.ca](http://www.roundupcanada.ca)

**NOTE: AVOID SPRAY CONTACT ON DESIRABLE PLANTS AS ANY PLANT MAY BE KILLED BY CONTACT.**

**DIRECTIONS FOR USE:**

Glyphosate is not to be applied using hand-wicking or hand-daubing methods.

| USE  | APPLICATION INSTRUCTIONS  | CONTROLS   | COVERAGE  |
|--|---|--|---|
| Undesirable vegetation control (i.e. hard to mow areas, around buildings). Lawn trimming/edging, patio weed control, vacant lots, storage & recreational areas, driveways. | Apply to vigorously growing plants. Apply to annual weeds anytime after emergence. For perennial weeds apply after seed heads, flowers or berries appear. | Most annual and perennial grasses and weeds, such as quackgrass, chickweed, ragweed, knotweed, poison ivy, Canada thistle, milkweed, bindweed. | Ensure uniform and complete spray coverage but not to the point of run-off. |
| Around Trees/Shrubs/Ornamentals  | Apply around base in a neat fashion. Avoid spray or drift contact with foliage, stems or green bark of desirable plants.                                  |  |   |
| Lawn/Garden renovation   | For best results on lawns, skip one mowing prior to treatment. Allow at least 3 days after treatment before disturbing the soil and/or replanting.        |  |   |
| Brush control  | Apply after leaves are fully expanded. Treat trees and brush up to 1.5 m in height only; for taller plants cut down and spray regrowth.                   | Most brush such as Poplar, Alder, Maple and Raspberry  |   |

**CAUTION:**

2020-08-04  
Sub.No.2020-1550

Avoid spray contact on desirable plants as any plant contacted may be killed. Do not spray in windy conditions. Do not spray to the point of run-off. Rain or watering two hours after application will not wash away effectiveness. However, heavy rainfall immediately after application may wash the chemical off the foliage and repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

**PRECAUTIONS:**

**CAUTION: EYE IRRITANT. HARMFUL IF SWALLOWED.** Avoid contact with eyes or prolonged contact with skin. Do not allow others such as children and pets on treatment area during application or to re-enter treated areas until spray has dried. Wash immediately after using this product. Do not contaminate water, foodstuffs, seed or feed by storage or disposal.

**FIRST AID:**

**If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If ingested,** call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If in eyes,** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:**

Treat symptomatically.

**DISPOSAL:**

Do not reuse empty container. Wrap empty container and discard in household garbage.

Roundup® is a registered trademark of Bayer Group. Used under license ©2020 Bayer Group. All rights reserved.

Quick Connect is a trademark of OMS Investments, Inc.

Not for reformulation or repackaging.  
Canada Patent No. 2,178,002.

Product Use Information: 1-866-476-7645  
Website: [www.roundupcanada.ca](http://www.roundupcanada.ca)