

2012-10-03  
2012-2567

## **CALYPSO 600**

LIQUID ALGAECIDE CONCENTRATE

CONTROLS ALGAE in Swimming Pool Waters

DOMESTIC

REG. NO. 25541 P.C.P. ACT

GUARANTEE: Poly [(oxyethylene dimethyliminio) ethylene- (dimethyliminio) ethylene dichloride] ..... 60%

CAUTION



POISON

Net Contents {Format}

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

Calypso, 42, rue de l'Artisan Victoriaville, (Québec) G6P 7E3  
1-866-470-7264

### **DIRECTIONS FOR USE**

1. Initial dosage for pool water having no visible algae is 70 mL of CALYPSO 600 per 10,000 L of pool water.
2. Weekly maintenance dosage is 25 mL of CALYPSO 600 per 10,000 L of pool water.
3. For pool water having visible algae, superchlorinate following label instructions of superchlorination product employed and wait at least 24 hours before adding weekly maintenance dosage.
4. For visible algae, when superchlorination is not employed, add 100 mL of CALYPSO 600 per 10,000 L of pool water.

**ENVIRONMENTAL HAZARDS** Toxic to aquatic organisms

**PRECAUTIONS** KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Do not mix with any other chemical. Avoid contact with skin and eyes.

**DISPOSAL** Rinse container thoroughly before discarding. Do not reuse empty container. Offer for recycling if available, or dispose in accordance with local and provincial regulations. . For information on disposal of unused, unwanted product, contact the manufacturer or the Provincial Regulatory Agency. Contact the manufacturer and the Provincial Regulatory Agency in case of a spill and for clean-up of spills.

**FIRST AID** **If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If swallowed**, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION** Treat symptomatically.