

Label Update 08-JUN-2010

For Dogs Weighing More Than 15 Kg.

BIO SPOT®
FOR LARGE DOGS

READ THE LABEL BEFORE USING
FOR USE ONLY ON DOGS
KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: Three 3.0 mL Applicator

CAUTION POISON

Toxic to cats

GUARANTEE:

Permethrin:45.0%

Pyriproxyfen:5.0%

- Controls mosquitoes for 3 weeks***
- Controls fleas for 3 weeks***
- Inhibits flea egg development for 3 months***
- Controls ticks, including the blacklegged (deer) tick, which may carry the organism which causes Lyme Disease, for 3 weeks***

DOMESTIC

REG. No 26921 P.C.P. ACT • U.S. Patent No. 5,942,525

Manufactured for:

Farnam Pet Products

A Division of Farnam Companies, Inc.
301 W. Osborn Road • Phoenix, AZ 85013
Kills fleas, ticks and mosquitoes

DO NOT USE ON CATS

DIRECTIONS FOR USE:

Do NOT administer by mouth. Use ONLY on dogs and puppies over 12 weeks of age and weighing between 15 and 30 kg.

NOT USE ON CATS OR OTHER ANIMALS. NOT TO BE USED ON DOGS OVER 30 KG.

Do not get this product in your dog's eyes or mouth. Repeat applications may be made if necessary (i.e., if the flea infestation persists), but do not apply more often than once every 3 weeks. IF ONLY TICKS OR MOSQUITOES ARE A PROBLEM, IT IS NOT NECESSARY TO USE A PRODUCT WHICH CONTAINS PYRIPROXYFEN. Efficacy of the product may be affected by washing the pet. A veterinarian should be consulted when pest infestation continues to be a problem on the pet. For Dogs Weighing between 15 and 30kg: (One Applicator): Apply one tube (3.0 mL) of **Bio Spot® for Large Dogs** solution to the dog's back at skin level between the shoulder blades. Remove product tube from packaging by tearing along perforation. Holding tube with notched end pointed up and away from face and body, cut off notched end of tube. Position the tip of the tube on the dog's back between the shoulder blades and squeeze out half the contents of the tube. Invert the applicator tube upright and go to the dog's tail. Position the tip of the tube at the base of the tail and apply the remaining contents of the tube to the top of the dog's rump near the base of the dog's tail. Wrap tube and put in garbage. Consult a veterinarian before

using on sick, aged or pregnant animals or animals under drug or other pesticide treatment. Do not use on nursing animals or animals under 12 weeks of age.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. HUMANS: Keep children and toddlers from touching the dog after application until the product has completely dried. Not for use on humans. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. May cause eye irritation. Wear gloves when applying. Wash thoroughly with soap and water after handling. For external use on dogs only. **PETS:** Discontinue use if adverse effects are observed. If condition persists, consult your veterinarian. **DO NOT USE ON CATS** or animals other than dogs. Cats that actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harm, including death. Sensitivities may occur after using any pesticide for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully. **Adverse Reactions:** Some animals may be sensitive to ingredients in this product. Reactions in dogs may include skin sensitivity. Dogs may show lethargy, increased pruritus (itchiness), erythema (redness), rash and hair discoloration or hair loss at the application site. Observe the dog following treatment. If sensitivity to the product occurs, discontinue use and bathe the dog with a mild, non-insecticidal shampoo. Seek veterinary advice.

□□□□□□□□ □□□□

FIRST AID: HUMANS: If Swallowed: Call a physician or Poison Control Centre IMMEDIATELY. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or, if available, by administration of syrup of ipecac. **DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. If In Eyes:** Flush eyes with plenty of water. Get medical attention if irritation persists. **If On Skin:** Wash affected area with soap and water. If irritation persists, get medical attention. Take container, label or product name and PCP Registration Number with you when seeking medical attention. **PETS:** If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE AND DISPOSAL:

Store in a cool, dry place. Do not reuse empty container. Wrap empty container and discard with household garbage.

Bio Spot for Large Dogs is an effective and easy to use product. **Bio Spot for Large Dogs** inhibits flea eggs laid on treated dogs from developing into biting adults for 3 months. **Bio Spot for Large Dogs** controls fleas, brown dog ticks, and American dog ticks for 3 weeks. **Bio Spot for Large Dogs** kills blacklegged (deer) ticks (which may carry the organism which causes Lyme Disease) for 3 weeks. Ticks present on the animal at the time of treatment may take several days to die. For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, furniture, etc. discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problem persists, an insecticidal premise treatment may be required.

Visit our Website at www.biospot.ca

In case of emergency or for product information, call (800) 548-2828,
Monday - Friday, 7:00 AM - 3:45 PM MST

Nylar® is a registered trademark of McLaughlin Gormley King Company.

©2001 Farnam Companies, Inc. • 02-1862

Resident Canadian Agent:

Farnam (Canada) Marketing, Inc.

#700, 10104 - 103 Avenue

Edmonton, AB T5J 0H8

602-285-1660

Farnam 02-1861

APPLICATOR LABEL

Side One

BIO SPOT LARGE DOG

GUARANTEE : Pyriproxyfen – 5.0%; Permethrin – 45.0%

REG NO. 26921 P.C.P. ACT

Side Two

READ BLISTER CARD LABEL BEFORE USING. 3.0 ml

CAUTION POISON DOMESTIC

Toxic to cats

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.