

2022-2295  
2022-07-22

## **SePRO**

### **A-Rest Solution**

- **For Professional Use**
- **Horticultural**

ACTIVE INGREDIENT: Ancymidol..... 264 mg/L

Contains 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride at 0.13% as a preservative.

REGISTRATION NO. 16393 PEST CONTROL PRODUCTS ACT

COMMERCIAL

NET CONTENTS: 113.6 L

READ THE LABEL BEFORE USING

#### **CAUTION:**

KEEP OUT OF REACH OF CHILDREN. MAY BE HARMFUL IF SWALLOWED. AVOID CONTACT WITH SKIN, EYES, OR CLOTHING. AVOID CONTAMINATION OF FOODSTUFFS, FEEDS AND FISH PONDS.

**SePRO Corporation**  
**Suite 600 – 11550 North Meridian Street**  
**Carmel, Indiana**  
**46032-4565**  
**(317) 580-8281**

### **FIRST AID**

Take container, label or product name and Pest Control Product Registration Number with you, when seeking medical attention.

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

### **PRECAUTIONS:**

Wear long pants, a long-sleeved shirt, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair.

Do not re-enter or allow worker entry into treated areas until 12 hours after application.

### **ENVIRONMENTAL PRECAUTIONS:**

DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters. TOXIC to aquatic plants.

## **GENERAL INFORMATION**

Read all directions carefully before applying.

A-REST for Height Control of Container-Grown Lilies, Poinsettias, and Chrysanthemum A-REST is a plant growth regulator, which effectively reduces internode elongation resulting in more desirable compact plants. When used as directed, A-REST produces no phytotoxic effects.

A-REST is not a substitute for any cultural practices (lighting, watering and temperature control). Use optimum cultural practices for growing plants listed on this label.

## **APPLICATION DIRECTIONS**

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Mix A-REST with water during spray tank filling operation. Specific rates will vary with plant species, variety, timing and type of treatment, and growing conditions. The degree of response desired often varies with location and individual cultural practices. The rate of application within the recommended range depends upon the degree of growth inhibition desired by the grower and should be established by using the product on a limited basis at varying rates.

The amount of growth inhibition from A-REST treatment depends upon the total quantity of active ingredients applied per plant. Treatment techniques and equipment should ensure uniform distribution of A-REST within the treated area.

## SOIL DRENCH APPLICATIONS

### Chrysanthemums

Dilute the A-REST solution as indicated in Table I, and apply as a drench treatment to moisten soil. Treat plants 5 - 15 cm in height (approximately two weeks following pinch).

### Lilies

Dilute the A-REST solution as indicated in Table I, and apply as a drench treatment to moisten soil. Treat plants from emergence to 30 cm in height. For optimum results, treatments should be made to plants 5 to 15 cm tall.

### Poinsettias

Dilute the A-REST solution as indicated in Table I, and apply as a drench treatment to moisten soil. Treat plants at pinch to 4 weeks after pinch, or 8 to 12 weeks before finishing.

TABLE I

No. of 15-cm Pots to be Treated	Drench Volume (mL)	Dilution Ratio A-REST to Water	Rate (mg/15-cm Pot)	Final Drench Concentration (mg/L)
250	120	250 mL 29.75 L	0.25	2
250	60	250 mL 14.75 L	0.25	4
250	60	500 mL 14.5 L	0.50	8
250	120	500 mL 29.5 L	0.50	4

2022-2295  
2022-07-22

## **WARNING**

Do not add any additional wetting agents to A-REST solutions. Plants treated with A-REST may require less water than untreated plants. Over-application or uneven application of A-REST will result in excessive or irregular growth control. A slight delay (two to five days) in flower development may occur at the higher recommended treatment levels.

## **DO NOT FREEZE**

### **DISPOSAL**

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.