

Label Update 08-JUN-2010

ACTIVE-3

DOMESTIC

SOLD THROUGH YOUR VETERINARIAN

Kills Fleas, Ticks and Mosquitoes

1 APPLICATOR - 1 DOSE BOX FOR SMALL DOGS
[2 APPLICATORS - 1 DOSE BOX FOR LARGE DOGS or 2 DOSE BOX FOR SMALL DOGS]
[3 APPLICATORS - 3 DOSE BOX FOR SMALL DOGS].

- Controls fleas for 3 weeks
- Inhibits flea egg development for 3 months.
- Controls ticks, including the blacklegged (deer) tick, which may carry the organism which causes Lyme Disease, for 3 weeks.
- Controls mosquitoes for 3 weeks.

GUARANTEE:

Permethrin45.0%
Pyriproxyfen.....5.0%

REG. No. 26620 P.C.P. ACT

**READ THE LABEL BEFORE USING
FOR USE ONLY ON DOGS
KEEP OUT OF REACH OF CHILDREN**

CAUTION



POISON

Toxic to cats

Net Contents: One 1.5 mL Applicator *[Two 1.5 mL Applicator][Three 1.5 mL Applicators]*



**2000 Chemin Georges, Lavaltrie, Québec J5T 3S5
1-800-363-1700**

Note: ITALICS = Alternate Text

Active-3 is an effective and easy to use product. **Active-3** inhibits flea eggs laid on treated dogs from developing into biting adults for 3 months **Active-3** controls fleas, brown dog ticks, and American dog ticks for 3 weeks **Active-3** kills blacklegged (deer) ticks (which may carry the organism which causes Lyme Disease) for 3 weeks. Ticks present on the animal at the time of treatment may take several days to die.

For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, furniture, etc. (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas. If pest problems persists, an insecticidal premise treatment may be required.

DIRECTIONS FOR USE:

RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE. Use ONLY on dogs and puppies 12 weeks of age or older. **DO NOT USE ON CATS OR OTHER ANIMALS.** NOT TO BE USED ON DOGS OVER 30 KG. Do NOT administer by mouth. Do not get this product in your dog's eyes or mouth. Repeat applications may be made if necessary (i.e., if the flea infestation persists), but do not apply more often than once every 3 weeks. A veterinarian should be consulted when pest infestation continues to be a problem on the pet. IF ONLY TICKS OR MOSQUITOES ARE A PROBLEM, IT IS NOT NECESSARY TO USE A PRODUCT WHICH CONTAINS PYRIPROXYFEN. Efficacy of the product may be affected by washing the pet. Consult a veterinarian before using on sick, aged or pregnant animals or animals receiving drug or other pesticide treatment.

For Dogs Weighing Less Than 15 kg (One Applicator): Apply one tube (1.5 mL) of Active-3 solution to the dog's back at skin level between the shoulder blades.

For Dogs Weighing 15 to 30 kg (Two Applicators): Apply one tube (1.5 mL) of Active-3 solution to the dog's back at skin level between the shoulder blades and apply a second tube (1.5 mL) to the dog's back at skin level directly in front of the base of the tail.

HOW TO APPLY:

Remove product tube from packaging by tearing along perforation. Holding tube with notched end pointed up and away from face and body, cut off notched end of tube. Position the tip of the tube on the dog's back between the shoulder blades and squeeze out half the contents of the tube. Invert the applicator tube upright and go to the dog's tail. Position the tip of the tube at the base of the tail and apply the remaining contents of the tube to the top of the dog's rump near the base of the dog's tail. Wrap tube and put in garbage.

PRECAUTIONS - HUMANS:

KEEP OUT OF REACH OF CHILDREN. Keep children and toddlers from touching the dog after application until the product has completely dried. Not for use on humans. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. May cause eye irritation. Wear gloves when applying. Wash thoroughly with soap and water after handling.

PRECAUTIONS - PETS:

For external use on dogs only. Do not use on nursing animals or animals under 12 weeks of age. Discontinue use if adverse effects are observed. If condition persists, consult your veterinarian.

DO NOT USE ON CATS or animals other than dogs. Cats that actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harm, including death. Sensitivities may occur after using any pesticide for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully.

Adverse Reactions : Some animals may be sensitive to ingredients in this product. Reactions in dogs may include skin sensitivity. Dogs may show lethargy, increased pruritis (itchiness), erythema (redness), rash and hair discolouration or hair loss at the application site. Observe the dog following treatment. If sensitivity to the product occurs, discontinue use and bathe the dog with a mild, non-insecticidal shampoo. Seek veterinary advice. Cats that actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harm, including death. Sensitivities may occur after using any pesticide for pets. Smaller animals are more likely to experience an adverse reaction. Observe carefully.

FIRST AID - HUMANS:

If Swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If In Eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If On Skin Or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

STORAGE AND DISPOSAL:

Store in a cool, dry place. Do not reuse empty container. Wrap empty container and discard with household garbage.

ACTIVE-3
DOMESTIC

GUARANTEE :

Permethrin45.0%
Pyriproxyfen.....5.0%

REG. No. 26620 P.C.P. ACT

READ BLISTER CARD LABEL BEFORE USING

CAUTION



POISON

Toxic to cats

1.5 ml

 **Vétoquinol**

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.