

# **OXAMYL TECHNICAL**

FOR MANUFACTURING, FORMULATING OR REPACKAGING

**DANGER** 



**POISON** 

GUARANTEE: Oxamyl 24%

REGISTRATION NO. 24949 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

NET CONTENTS: \_\_\_\_\_

Corteva Agriscience Canada Company Suite 240, 115 Quarry Park Rd. SE Calgary, AB T2C 5G9 1-800-667-3852

<sup>™</sup>Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN AND PREVENT ACCESS BY `

UNAUTHORIZED PERSONNEL. Avoid breathing spray mist or vapours. Avoid contact with skin, eyes and clothing.

Do not contaminate any body of water. Do not contaminate food. Wear chemical resistant gloves during mixing, loading and cleanup. Wash hands and face before eating, drinking, smoking and using the toilet.

Take a shower immediately after use.

Wash protective clothing in detergent and hot water, separately from household

laundry, before reuse.

Fatal or poisonous if swallowed.

Harmful or fatal if absorbed through the skin.

May be fatal if inhaled.

#### FIRST AID:

IN CASE OF POISONING, CONTACT A PHYSICIAN OR A POISON CONTROL CENTRE IMMEDIATELY.

EYE CONTACT: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove

contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a

poison control centre or doctor for treatment advice.

SKIN CONTACT: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20

minutes. Call a poison control centre or doctor for treatment advice.

INGESTION: Call a poison control centre or doctor immediately for treatment advice. Have

person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an

unconscious person.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Oxamyl is a carbamate which is a cholinesterase inhibitor. Typical symptoms of overexposure to cholinesterase inhibitors include malaise, muscle weakness, dizziness and sweating. Headache, salivation, nausea, vomiting, abdominal pain and diarrhea are often prominent. A life-threatening poisoning is signified by loss of consciousness, incontinence, convulsions and respiratory depression with a secondary cardiovascular component. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). However, if a blood sample is taken several hours after exposure, it is unlikely that blood cholinesterase activities will be depressed, due to rapid reactivation of cholinesterase. Atropine, only by injection, is the preferable antidote. Do not use pralidoxime. In cases of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration. With oral exposure, the decision of whether to induce vomiting or not should be made by an attending physician.

### **DIRECTIONS FOR USE:**

To be used only in the manufacture of a pesticide which is registered under the *PEST CONTROL PRODUCTS ACT*.

## STORAGE:

Store product in original container only. Store product in cool, dry, well ventilated place, away from other pesticides, fertilizer, food or feed. Do not contaminate water, food or feed by storage. Keep container closed.

# **DECONTAMINATION AND DISPOSAL:**

Canadian formulators using this product should dispose of unwanted active ingredient and containers in accordance with municipal or provincial regulations. For additional details and information on clean-up of spills, contact the provincial regulatory agency or the manufacturer.

### NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Purchase of this material does not confer any rights under patents of countries outside Canada.

™Trademark of Corteva Agriscience and its affiliated companies.