

2021-6463
2022-01-14

BARQUAT MB-40

Swimming Pool Algaecide

Controls Algae in Swimming Pool Waters

DOMESTIC

ACTIVE INGREDIENT :

Alkyl (40% C₁₂, 50% C₁₄, 10% C₁₆) dimethyl benzyl ammonium chloride 40%

KEEP OUT OF REACH OF CHILDREN

DANGER



POISON



CORROSIVE

READ THE LABEL BEFORE USING

REGISTRATION NO. 27342 PEST CONTROL PRODUCTS ACT

Net Contents _____

ARXADA, LLC
412 Mount Kemble Avenue, Suite 200S
Morristown NJ 07960
201-316-9200

DIRECTIONS FOR USE:

Ensure all pool equipment is working properly. Backwash the filter system following manufacturer’s directions. Adjust sanitizer, pH, total alkalinity, and calcium hardness to the recommended levels identified on the label of the sanitizer product being used. In outdoor chlorine treated pools, sanitizer can be added at levels between 30-100 ppm to reduce the degradative effect of sunlight upon chlorine residual. Presence of metal may require the addition of a stain and scale inhibitor to prevent staining of pool surface. Consult your pool professional.

Wear rubber gloves when handling this concentrate.

DO NOT apply when swimmers are in the immediate vicinity (a 15 minute restricted-entry interval is recommended.)

1. **INITIAL APPLICATION** for pool water having no visible algae: Pour directly into pool at a rate of 50 mL of this product per 10,000L of water. More rapid distribution will be achieved by pouring a portion of the specified amount into several areas of the pool until all of the required quantity has been added. Recirculation accomplished by running the filter will assure rapid dispersion of product.
2. **WEEKLY MAINTENANCE:** Pour directly into pool at a rate of 25 mL per 10,000L of water. Algicide additions should be made even when the pool is not in use for periods of time since a heavy accumulation of algae growth is difficult to remove. More rapid distribution will be achieved by pouring a portion of the specified amount into several areas of the pool until the required quantity has been added. Recirculation accomplished by running the filter will assure rapid dispersion of product.
3. For pool water having visible algae, superchlorinate following label instructions of superchlorinate product employed and wait at least 24 hours before adding weekly maintenance dosage.
4. For visible algae, when superchlorination is not employed, add 125mL of Barquat MB-40 per 10,000 L of pool water.
5. Re-entry interval: swimmers are allowed to re-enter the pool 15 minutes after application of the product.
6. Ensure that water quality conditions are maintained for the successful use of the product.

Swimming Pool Capacity Litres of Water	Initial Treatment	Weekly Maintenance
10,000	100 mL	25 mL
50,000	250 mL	100 mL
80,000	400 mL	200 mL
100,000	500 mL	250 mL
150,000	750 mL	375 mL
180,000	900 mL	450 mL
200,000	1 L	500 mL

FOR WINTERIZING POOLS BEFORE FREEZING:

1. Maintain recommended sanitizer, pH, total alkalinity, and calcium hardness levels identified on the label of the swimming pool sanitizer product being used.
2. For pool water having visible algae, superchlorinate following instructions of superchlorinate product employed and wait for at least 24 hours before adding Barquat MB-40.
3. For covered pools, apply a Barquat MB-40 dosage of 250mL per 10,000L. Add directly into pool water, walking around the pool, so that it is evenly distributed. Allow water to circulate for 2 to 4 hours.

For uncovered pools, apply a Barquat MB-40 dosage of 250mL per 10,000L. Add directly into pool water, walking around the pool, so that it is evenly distributed. Allow water to circulate for 2 to 4 hours. Uncovered pools may require a weekly maintenance dosage of 25mL per 10,000L until the pool water has frozen.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. CORROSIVE. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Harmful if inhaled. Avoid breathing vapour or spray mist. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID:

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE:

Do not store on side. Avoid creasing or impacting of side walls. Store in original container. Do not reuse empty container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use. Keep from freezing.

2021-6463
2022-01-14

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills. Rinse the emptied container thoroughly and add the rinsings to the treatment site.