

2022-1663
2022-11-21

This label has been updated according to the special review decision of Isoxaflutole RVD2022-04. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 04/07/2024 in accordance with the phase out period set out in the special review decision of Isoxaflutole RVD2022-04. This previously approved label will be provided upon request by emailing hc.pma.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

2022-1663
2022-11-21

****Container Label****



GROUP	27	HERBICIDE
-------	-----------	-----------

**CONVERGE 75 WDG HERBICIDE
FOR SALE FOR USE IN EASTERN CANADA AND BRITISH COLUMBIA ONLY**

For use in field corn

AGRICULTURAL

ACTIVE INGREDIENT:
ISOXAFLUTOLE.....75%

REGISTRATION NUMBER: 26142 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION EYE IRRITANT

Product Information: 1-888-283-6847

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. SE
Calgary, Alberta
T2C 3G3

In case of spills, poisoning or fire, telephone emergency response number
1-800-334-7577 (24 hours per day)

NET CONTENTS: 0.140 Kg - Bulk

2022-1663
2022-11-21

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes and clothing. Wear coveralls over long-sleeved shirt and long pants, and rubber boots during all activities. Wear chemical resistant gloves and protective eyewear during mixing/loading, clean-up and repair. Wash from skin or eyes immediately. Avoid breathing dust or spray mist. After use, wash hands and other exposed skin.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings. If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth if possible. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: No specific antidote. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS: TOXIC to aquatic organisms and terrestrial plants. Avoid spray drift onto crops other than those recommended. Observe buffer zones specified under DIRECTIONS FOR USE.

STORAGE: Do not contaminate water, food or feed by storage or disposal. To prevent contamination, store this product away from food or feed. Store the container tightly closed, away from seeds, fertilizer, plants and foodstuffs. Store in a cool, dry secured storage area. Do not store this product under wet conditions.

DISPOSAL:

RECYCLABLE CONTAINER: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

2022-1663
2022-11-21

RETURNABLE CONTAINER: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

DISPOSAL OF UNUSED, UNWANTED PRODUCT: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

220425

2022-1663
2022-11-21

****Booklet Label****



GROUP	27	HERBICIDE
-------	-----------	-----------

CONVERGE 75 WDG HERBICIDE

FOR SALE FOR USE IN EASTERN CANADA AND BRITISH COLUMBIA ONLY

For use in field corn

AGRICULTURAL

ACTIVE INGREDIENT:
ISOXAFLUTOLE.....75%

REGISTRATION NUMBER: 26142 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION EYE IRRITANT

Product Information: 1-888-283-6847

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. SE
Calgary, Alberta
T2C 3G3

In case of spills, poisoning or fire, telephone emergency response number
1-800-334-7577 (24 hours per day)

NET CONTENTS: 0.140 Kg - Bulk

2022-1663
2022-11-21

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes and clothing. Wear coveralls over long-sleeved shirt and long pants, and rubber boots during all activities. Wear chemical resistant gloves and protective eyewear during mixing/loading, clean-up and repair. Wash from skin or eyes immediately. Avoid breathing dust or spray mist. After use, wash hands and other exposed skin.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth if possible. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: No specific antidote. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS: TOXIC to aquatic organisms and terrestrial plants. Avoid spray drift onto crops other than those recommended. Observe buffer zones specified under DIRECTIONS FOR USE.

STORAGE: Do not contaminate water, food or feed by storage or disposal. To prevent contamination, store this product away from food or feed. Store the container tightly closed, away from seeds, fertilizer, plants and foodstuffs. Store in a cool, dry secured storage area. Do not store this product under wet conditions.

DISPOSAL:

RECYCLABLE CONTAINER: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

2022-1663
2022-11-21

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
 2. Make the empty, rinsed container unsuitable for further use.
- If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL APPLICATION INFORMATION:

CONVERGE 75 WDG Herbicide is a water dispersible granule formulation of isoxaflutole at a concentration of 75% active ingredient. CONVERGE 75 WDG Herbicide is a selective herbicide for season long control of important broadleaf and grass weeds infesting field corn.

Surface Preplant: CONVERGE 75 WDG Herbicide can be applied to the soil surface up to 14 days prior to planting. CONVERGE 75 WDG Herbicide must be tankmixed with atrazine when applied surface preplant. If weed growth is present at the time of application, Roundup Transorb, Roundup Weathermax, Roundup Ultra, or Vantage Plus Max can be added to the CONVERGE 75 WDG Herbicide + atrazine treatment for burndown control of these weeds. Do not incorporate CONVERGE 75 WDG Herbicide.

Preemergence: CONVERGE 75 WDG Herbicide can also be applied after planting to just prior to crop emergence. Atrazine and/or Roundup Transorb, Roundup Weathermax, Roundup Ultra, or Vantage Plus Max can be tankmixed with preemergent applications of CONVERGE 75 WDG Herbicide.

When activated by rainfall, CONVERGE 75 WDG Herbicide will control susceptible weed seedlings that are up to 5 cm in height. Shallow tillage may be necessary to control weeds beyond this stage.

SPECIAL PRECAUTIONS:

Under certain conditions, temporary yellowing of some corn leaves may occur when the plant is in the seedling stage. These conditions include cold weather, excessive moisture and/or compacted soils. The symptoms are most visible where excessive rates have been applied, such as sprayer overlaps. The crop typically recovers quickly.

Plant corn at least 4 cm deep. Failure to close the seed furrow may allow herbicide spray to directly contact the seed which can cause injury. If drought conditions persist after either a pre-

2022-1663
2022-11-21

plant surface or pre-emergent application, weed control, especially annual grass control, may be reduced.

DO NOT apply CONVERGE 75 WDG Herbicide to field corn grown on loamy sands or sands and/or soils with less than 2 % O.M.

DO NOT use on sweet or seed corn.

DO NOT graze or feed the treated immature corn to livestock prior to 60 days after application of CONVERGE 75 WDG Herbicide.

Do not harvest corn grain within 136 days of application.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

DIRECTIONS FOR USE:

ADDITIONAL ENVIRONMENTAL PRECAUTIONS:

- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.
- DO NOT apply by aerial application equipment.

Spray Buffer Zones:

A spray buffer zone is not required for:

- uses with hand-held application equipment permitted on this label,

- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage.

The spray buffer zones specified in the table below are for isoxaflutole. These spray buffer zones are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat:
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Field corn	1	1	1	1	15

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

MIXING INSTRUCTIONS:

Fill spray tank with ¼ to ½ of the required volume of water prior to the addition of CONVERGE 75 WDG Herbicide. Add CONVERGE 75 WDG Herbicide slowly to the tank and agitate. If CONVERGE 75 WDG Herbicide is applied in tank-mixtures with other pesticides, add CONVERGE 75 WDG Herbicide to the spray tank first and ensure that it is thoroughly dispersed before continuing to fill the spray tank and adding other pesticides. Continue to fill the tank with water to the desired volume while agitating. Continue agitation when applying to ensure a uniform spray mixture.

Application with water or nitrogen solution as a diluent carrier: Nitrogen solution (28-0-0) may replace all or part of the water as a carrier. Check for compatibility by combining all ingredients in a small container in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear 5-15 minutes after mixing.

APPLICATION PROCEDURES:

Apply CONVERGE 75 WDG Herbicide in conventional, minimum tillage, or zero tillage crop management systems up to 14 days prior to planting, or after planting but before weeds or crop emerge. DO NOT apply after the corn emerges or crop injury may occur.

2022-1663
2022-11-21

Apply CONVERGE 75 WDG Herbicide alone or in tank-mixtures, by ground equipment only, in a minimum of 200 litres of water per hectare at a spray pressure of 138-275 kPa. For best results use flat fan nozzles or comparable nozzles to achieve uniform spray distribution. DO NOT apply with hollow cone nozzles or with other application equipment which does not provide uniform coverage.

WEEDS CONTROLLED: Apply CONVERGE 75 WDG Herbicide at 105 g/ha to provide season long control of the following weeds:

<i>common lamb's quarters*</i>	<i>redroot pigweed*</i>
<i>common ragweed*</i>	<i>tall waterhemp*</i>
<i>eastern black nightshade</i>	<i>plantain, seedling</i>
<i>dandelion, seedling</i>	<i>witchgrass</i>
<i>crabgrass, smooth and large</i>	<i>mustard, wild and wormseed</i>
<i>velvetleaf</i>	

* Includes triazine and ALS resistant biotypes

Apply CONVERGE 75 WDG Herbicide at 140 g/ha to provide season long control of the above listed weeds plus:

<i>barnyard grass</i>	<i>green foxtail</i>
-----------------------	----------------------

TANK-MIXTURES:

CONVERGE 75 WDG Herbicide + atrazine: Apply CONVERGE 75 WDG Herbicide at 105 g product per hectare plus atrazine at 800 g ai/ha for control of the weeds listed above plus:

<i>wild buckwheat</i>	<i>yellow foxtail</i>
<i>lady's thumb</i>	

Apply CONVERGE 75 WDG Herbicide at 140 g product per hectare plus atrazine at 1063 g ai/ha for control of the weeds listed above plus:

fall panicum
proso millet (suppression only)

Observe all directions for use, precautions, and restrictions on the Atrazine label.

CONVERGE 75 WDG Herbicide + Roundup Transorb, Roundup Weathermax, Roundup Ultra, Vantage Plus Max: Apply CONVERGE 75 WDG Herbicide at 105 – 140 g per hectare tankmixed with Roundup Transorb at 2.5 L/ha, or with Vantage Plus Max at 1.88 L/ha, or with Roundup Weathermax or Roundup Ultra at 1.67 L/ha, for burndown control of emerged weeds in all tillage management systems and improved control of established dandelion in zero-tillage management systems. A three-way tankmix of CONVERGE 75 WDG Herbicide + atrazine + Roundup Transorb, Roundup Weathermax, Roundup Ultra, or Vantage Plus Max can be used to

2022-1663
2022-11-21

provide residual control of the weeds listed in the CONVERGE 75 WDG Herbicide + atrazine section.

ROTATIONAL CROP INFORMATION:

CONVERGE 75 WDG Herbicide treated field corn fields may be planted to field corn, soybean, alfalfa, potato, tomato, timothy, processing peas, spring canola, barley (spring and winter), spring oats and spring wheat the year after treatment. Caution should be used when planting tomato 12 months following an application of CONVERGE 75 WDG Herbicide + atrazine if conditions were exceptionally dry during the season of application. CONVERGE 75 WDG Herbicide treated field corn fields may be planted to all classes of dry common beans in the second year after treatment. Treated corn fields may be planted to winter wheat in the same year of a spring CONVERGE 75 WDG Herbicide application provided that 4 months have elapsed since the CONVERGE 75 WDG Herbicide application. A field bioassay must be conducted the year prior to growing any other crop of interest to confirm crop safety.

FIELD BIOASSAY:

Select a representative area or areas of the field previously treated with CONVERGE 75 WDG Herbicide to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results. Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones. Use standard tillage and seeding equipment to plant the bioassay. Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also, plant into an adjacent area not treated with CONVERGE 75 WDG Herbicide to use as a comparison. As the crop(s) emerges and grows, examine these key points in CONVERGE 75 WDG Herbicide-treated and non-treated areas:

crop stand	plant colour and vigour
root development	yield
rate of growth	

Allow the bioassay crop(s) to grow to maturity while making your observations. Do not overspray the test strips with herbicides that may damage the bioassay crop(s). If the bioassay indicates that CONVERGE 75 WDG Herbicide residues are still present, continue cropping only to those crops listed on the label, and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally. **DO NOT ROTATE TO OTHER CROPS UNTIL BIOASSAY INDICATES NORMAL GROWTH WITH NO YIELD REDUCTIONS.**

Failure to follow the recropping guidelines could result in injury to seeded crop(s).

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that CONVERGE 75 WDG Herbicide is a Group 27 herbicide. Any weed population may contain or develop plants naturally resistant to CONVERGE 75 WDG Herbicide and other Group 27 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of CONVERGE 75 WDG Herbicide or other Group 27 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by using an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Bayer representative, or contact Bayer CropScience Inc. toll-free at 1-888-283-6847 or visit www.cropscience.bayer.ca.