AGRICULTURAL

Norac Concepts Inc. P.O. Box 31097 Guelph, ON N1H 8K1 (519) 821-3633

Net Contents: 10 kg - 25 kg

GROUP M FUNGICIDE

CONDOR MZ

POTATO SEED PIECE TREATMENT To Control Fusarium Seed Piece Decay Contains Alder Bark

READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN DUST GUARANTEE: mancozeb 16%

REGISTRATION NO. 26158 PEST CONTROL PRODUCTS ACT

PRECAUTIONS

Store product in a cool, dry, ventilated place. Do not allow product to become wet or overheated during storage as chemical changes may occur which impair fungicidal effectiveness and may also generate flammable vapours. May irritate eyes, nose, throat and skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating, drinking or smoking. When treating or handling treated seed, work in a well-ventilated area, and wear a suitable dust mask, goggles and gloves. Do not contaminate food or feed. Do not store near food or feed. Do not contaminate any body of water. Treated seed should be labelled "POISONOUS TO MAN AND ANIMALS. This seed has been treated with mancozeb for the control of fusarium decay. Do not use for food or feed purposes".

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca

FIRST AID

If swallowed, Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

EMERGENCY TELEPHONE NUMBER

All hours, (613) 787-5620 ONLY for health and environmental information.

TOXICOLOGICAL INFORMATION

Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

- 2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal
- 3. Make the empty container unsuitable for further use.
- 4. Dispose of container in accordance with provincial requirements.

5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE

Apply at the rate of 500 g per 100 kg of seed. Thoroughly coat the surface of whole or cut seed pieces with dust. If treated whole seed is cut, make a second application to protect the cut surfaces. Plant as soon as possible after treating. However, if planting of cut seed is delayed beyond 2 days after treating, seed should be stored in open crates to allow air drying until dry before bagging or loose piling.

RESISTANCE MANAGEMENT

For resistance management, please note that CONDOR MZ contains a Group M fungicide. Any fungal population may contain individuals naturally resistant to CONDOR MZ and other Group M fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

• Where possible, rotate the use of CONDOR MZ or other Group M fungicides with different groups that control the same pathogens.

• Use tank mixtures with fungicide from a different group when such use is permitted.

• Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.

• Monitor treated fungal populations for resistance development.

• If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-

management and/or IPM recommendations for specific crops and pathogens.

• For further information and to report suspected resistance, contact Norac Concepts at (613) 841-2907 or at <u>www.noracconcepts.com</u>.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.