

**EVEREST® SOLUPAK® 70 DF**  
**70% WATER DISPERSIBLE GRANULAR HERBICIDE**  
**IN WATER-SOLUBLE PACKETS**

<b>GROUP</b>	<b>2</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**FOR POSTEMERGENCE CONTROL OF WILD OAT AND GREEN FOXTAIL  
IN ALL TYPES OF SPRING WHEAT AND DURUM WHEAT**

**FOR SALE AND FOR USE IN MANITOBA, SASKATCHEWAN, ALBERTA AND  
PEACE RIVER REGION OF BRITISH COLUMBIA ONLY**

**AGRICULTURAL**

**ACTIVE INGREDIENT: Flucarbazone.....66%**  
**(present as flucarbazone-sodium)**

**Warning, contains the allergen milk.**

**REGISTRATION NO. 26448 PEST CONTROL PRODUCTS ACT**

**READ THE LABEL AND BOOKLET BEFORE USING**  
**KEEP OUT OF REACH OF CHILDREN**  
**(See back panel for additional precautions)**

**DO NOT SELL WATER-SOLUBLE PACKETS SEPARATELY.**  
**KEEP WATER-SOLUBLE PACKETS IN THIS CONTAINER AND STORE IN A COOL, DRY**  
**PLACE.**  
**DO NOT ALLOW PACKETS TO BECOME WET PRIOR TO ADDING TO THE SPRAY**  
**TANK.**

**CONTAINS FOUR 174 GRAM WATER-SOLUBLE PACKETS**  
**TOTAL NET CONTENTS: 696 GRAMS**

**For Product Information call: 1-800-438-6071**

**UPL AgroSolutions Canada Inc.**  
**630 Freedom Business Center, Suite 402**  
**King of Prussia, PA 19406**

**FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.**

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

### **PRODUCT INFORMATION**

EVEREST SOLUPAK 70 DF Herbicide is a selective postemergence herbicide for use in spring wheat (hard red spring, Canada Prairie spring, soft white spring, and extra strong or utility) and durum wheat.

**EVEREST SOLUPAK 70 DF Herbicide must be tank mixed with a surfactant listed on this label. For broader spectrum activity, EVEREST SOLUPAK 70 DF Herbicide may be tank mixed with a broad-leaf herbicide listed on this label. See section on Tank Mixes for recommended products.**

EVEREST SOLUPAK 70 DF Herbicide is absorbed by foliage of wild oat and green foxtail. Wild oat and green foxtail cease growth soon after EVEREST SOLUPAK 70 DF Herbicide application. Complete weed control may not be seen for one to two weeks. However, weeds will have stopped growing and the competitive effects removed.

EVEREST SOLUPAK 70 DF Herbicide will control wild oat biotypes, which have developed resistance to ACCase (Group 1) and triallate (Group 8) herbicides. It will also control green foxtail biotypes resistant to ACCase (Group 1) and dinitroaniline (Group 3) herbicides.

One pouch of 4 x 174 g water-soluble packets contains enough EVEREST SOLUPAK 70 DF Herbicide to treat 16 - 32 hectares (40 – 80 acres) depending on the target grassy weed to be controlled.

### **DIRECTIONS FOR USE**

Read the entire DIRECTIONS FOR USE before using EVEREST SOLUPAK 70 DF Herbicide.

- Ground application only. **DO NOT APPLY BY AIR.**
- Do not mix, load or clean spray equipment within 10 metres of well-heads or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc.
- Buffer Zones: Overspray or drift to sensitive habitats should be avoided. A buffer zone of 20 m is required between the downwind edge of the boom and the closest edge of sensitive terrestrial habitats including forested areas, shelter belts, woodlots, hedgerows, and shrublands. A buffer zone of 35 m is required between the downwind edge of the boom and the closest edge of sensitive aquatic habitats including sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, wetlands, and wildlife habitat at the edge of these bodies of water. Do not contaminate these habitats when cleaning and rinsing spray equipment or containers.
- Consult the label of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.
- Do not apply during periods of dead calm, when winds are gusty or when wind speed is greater than 15 km/hr at 2 m high above ground at the site of application.

- **Do not apply if it is raining or if rainfall is expected within one (1) hour after application.**
- Do not feed or graze treated fields. Wheat grain or straw from treated fields may be fed to livestock.
- Do not allow this chemical to drift on to other crops, especially canola, tame oats, or other non target crops.
- Observe minimum interval to harvest of 80 days after treatment.
- Do not treat wheat underseeded to legumes.

### TIMING OF APPLICATION

Do not apply EVEREST SOLUPAK 70 DF Herbicide before crop emergence is complete. Research has demonstrated that optimum wheat yield is obtained by early removal of grassy weeds before tillering. Apply EVEREST SOLUPAK 70 DF Herbicide to wheat that has a minimum of one leaf to a maximum of four leaves on the main stem, plus two tillers. Application beyond this stage may result in yield losses due to weed competition.

### RECOMMENDED APPLICATIONS

APPLICATION TIMING		
CROP/WEEDS	GROWTH STAGE	REMARKS
Spring Wheat Durum Wheat	1 leaf to 6 total leaves	1 leaf to 4 leaves on main stem, plus 2 tillers
Wild oat ( <i>Avena fatua</i> ), Green foxtail ( <i>Setaria viridis</i> ), Volunteer tame oat ( <i>Avena sativa</i> )	1 leaf to 6 total leaves	1 leaf to 4 leaves on main stem, plus 2 tillers
Shepherd's Purse ( <i>Capsella bursa-pastoris</i> )	2 to 6 leaf stage	
Redroot Pigweed ( <i>Amaranthus retroflexus</i> )	2 to 6 leaf stage	
Volunteer Canola ( <i>Brassica napus</i> )	2 to 6 leaf stage	Will not control imidazolinone-tolerant (CLEARFIELD®) canola varieties.
Green Smartweed ( <i>Polygonum scabrum moench</i> )	2 to 6 leaf stage	
Wild Mustard ( <i>Brassica kaber</i> )	2 to 6 leaf stage	
Stinkweed ( <i>Thlaspi arvense</i> )	2 to 9 leaf stage	

GRASS WEED CONTROL IN SPRING AND DURUM WHEAT			
CROP	WEED	RATE	REMARKS
Spring Wheat  <b>Including:</b> hard red spring, Canada Prairie spring, soft white spring and extra strong (utility) wheat	Wild oat, green foxtail (wild millet)	28.5 g/ha (each packet treats 6 hectares or 15 acres)	For control of wild oat and green foxtail, apply EVEREST SOLUPAK 70 DF Herbicide plus a surfactant when the majority of wheat has a minimum of one leaf to a maximum of four leaves on the main stem, plus two tillers. EVEREST SOLUPAK 70 DF may be tank mixed with a broadleaf herbicide listed on the label.
	Green foxtail (wild millet) ONLY	21.5 g/ha (each packet treats 8 hectares or 20 acres)	Wild oat should be no larger than 4-leaf, 2 tillers.  Green foxtail should be no larger than 4-leaf, 2 tillers.
	Durum Wheat  Volunteer tame oat	28.5 g/ha (each packet treats 6 hectares or 15 acres)	For control of volunteer tame oat, apply EVEREST SOLUPAK 70DF Herbicide plus a surfactant, when the majority of wheat has a minimum of one leaf to a maximum of four leaves on the main stem, plus two tillers.  Volunteer tame oat should be no larger than 4-leaf, 2 tillers.

BROADLEAF WEED CONTROL IN SPRING AND DURUM WHEAT		
WEED	RATE	REMARKS
Shepherd's Purse ( <i>Capsella bursa-pastoris</i> )	28.5 g/ha  (each packet treats 6 hectares or 15 acres)	For control of shepherd's purse, apply EVEREST SOLUPAK 70 DF Herbicide to weeds at the 2 to 6 leaf stage.
Redroot Pigweed ( <i>Amaranthus retroflexus</i> )		For control of redroot pigweed, apply EVEREST SOLUPAK 70 DF Herbicide to weeds at the 2 to 6 leaf stage.
Wild Mustard ( <i>Brassica kaber</i> )		For control of wild mustard, apply EVEREST SOLUPAK 70 DF Herbicide to weeds up to 5 cm in height.
Stinkweed ( <i>Thiaspi arvense</i> )		For control of stinkweed, apply EVEREST SOLUPAK 70 DF Herbicide to weeds up to 5 cm in height.
Volunteer Canola ( <i>Brassica napus</i> )		For control of volunteer canola, apply EVEREST SOLUPAK 70 DF Herbicide to weeds up to 5 cm in height.
Green Smartweed ( <i>Polygonum scabrum moench</i> )		For control of green smartweed, apply EVEREST SOLUPAK 70 DF Herbicide to weeds up to 5 cm in height.

Wheat exposed to water-logged or saturated soils, or temperature extremes such as heat or freezing weather, or drought, low fertility or plant disease at application time could show unacceptable injury symptoms. Weed control also may be reduced by these same conditions.

### TANK MIXES

For broader spectrum control of both annual grasses and broadleaf weeds, EVEREST SOLUPAK 70 DF Herbicide may be mixed with one of the broadleaf herbicides plus a surfactant listed in the following table. With all tank-mix partners, read and follow the use directions, rates, precautions, timing, recropping restrictions, grazing interval restrictions and recommendations on the broadleaf herbicide and surfactant labels.

APPROVED TANK MIX PARTNERS IN SPRING WHEAT					
BROADLEAF HERBICIDE TANK MIX PARTNER		BROADLEAF HERBICIDE PRODUCT RATES per hectare	EVEREST SOLUPAK 70 DF Herbicide Rate		Surfactants (0.25% v/v or 0.25L/100L)
			Wild Oat Green Foxtail	Green Foxtail ONLY***	
1	2,4-D Amine	Up to 560 g a.e./ha	43 g/ha	21.5 g/ha	Surf 92
					Super Spreader LI 700® Penetrating Surfactant Agral® 90 Ag-Surf
2	2,4-D Ester	Up to 560 g a.e./ha	43 g/ha	21.5 g/ha	Surf 92
					Super Spreader LI 700® Penetrating Surfactant

APPROVED TANK MIX PARTNERS IN SPRING WHEAT					
BROADLEAF HERBICIDE TANK MIX PARTNER		BROADLEAF HERBICIDE PRODUCT RATES per hectare	EVEREST SOLUPAK 70 DF Herbicide Rate		Surfactants (0.25% v/v or 0.25L/100L)
			Wild Oat Green Foxtail	Green Foxtail ONLY***	
					Agral® 90 Ag-Surf®
3	**Ally® + 2,4-D Amine	5 or 7.5 g/ha Ally + up to 560 g ae/ha 2,4-D Amine	***43 g/ha	No	Agral® 90 Ag-Surf®
4	**Ally® + 2,4-D Ester	5 or 7.5 g/ha Ally + up to 560 g ae/ha 2,4-D Ester	***43 g/ha	No	Agral® 90 Ag-Surf®
5	Attain® Herbicide Tank Mix	0.6 L/ha Attain A + 1 L/ha Attain B	***43 g/ha	No	Agral® 90 Ag-Surf®
6	Buctril® M	1 L/ha	43 g/ha	21.5 g/ha	Surf 92 Super Spreader LI 700® Penetrating Surfactant Agral® 90 Ag-Surf®
7	Curtail® M	1.5 – 2.0 L/ha	***43 g/ha	No	Agral® 90 Ag-Surf®
8	*DyVel®	1.25 L/ha	***43 g/ha	No	Agral® 90 Ag-Surf®
9	*DyVel® DS	850 mL/ha	***43 g/ha	No	Agral® 90 Ag-Surf®
10	Estaprop® Dichlorprop-D	1.75 L/ha	43 g/ha	21.5 g/ha	Surf 92 Super Spreader LI 700® Penetrating Surfactant Agral® 90 Ag-Surf®
11	Express Pack®	10 grams/ha Express + 0.625 L/ha 2,4-D	43 g/ha	No	Agral® 90 Ag-Surf®
12	Frontline™ 2,4-D Herbicide Tank Mix	100 ml/ha Frontline 2,4-D A + 1.0 L/ha Frontline 2,4-DB	43 g/ha	No	Agral® 90 Ag-Surf®
13	Frontline™ Herbicide Tank Mix	100 ml/ha Frontline A + 0.84 L/ha Frontline B	43 g/ha	No	Agral® 90 Ag-Surf®
14	MCPA Amine	up to 560 g ae/ha	43 g/ha	No	Agral® 90 Ag-Surf®
15	MCPA Ester	up to 560 g ae/ha	43 g/ha	No	Agral® 90 Ag-Surf®
16	Pardner®	1 L/ha	43 g/ha	No	Agral® 90 Ag-Surf®
17	Prestige® Herbicide Tank Mix	0.8 L/ha Prestige A + 2.0 L/ha Prestige B	***43 g/ha	No	Agral® 90 Ag-Surf®
18	**Refine Extra® + 2,4-D Amine	20 g/ha Refine Extra + up to 560 g ae/ha 2,4-D Amine	43 g/ha	21.5 g/ha	Surf 92 Super Spreader LI 700® Penetrating Surfactant Agral® 90 Ag-Surf®
19	**Refine Extra® + 2,4-D Ester	20 grams/ha Refine Extra + up to 560 g ae/ha 2,4-D Ester	43 g/ha	No	Agral® 90 Ag-Surf®
20	*Target®	1 to 1.5 L/ha	***43 g/ha	No	Agral® 90 Ag-Surf®
21	Spectrum™ Herbicide Tank Mix	100 ml/ha Spectrum A + 1.5 L/ha Spectrum B	43 g/ha	No	Agral® 90 Ag-Surf®
22	Thumper®	1 L/ha	43 g/ha	No	Surf 92

APPROVED TANK MIX PARTNERS IN SPRING WHEAT					
BROADLEAF HERBICIDE TANK MIX PARTNER		BROADLEAF HERBICIDE PRODUCT RATES per hectare	EVEREST SOLUPAK 70 DF Herbicide Rate		Surfactants (0.25% v/v or 0.25L/100L)
			Wild Oat Green Foxtail	Green Foxtail ONLY***	
					Super Spreader <del>LI 700® Penetrating Surfactant</del> Agral® 90 Ag-Suri®
23	**Unity® Herbicide Tank Mix	0.5 L/ha Unity EC + 10.6 grams/ha Unity 75 WG	***43 g/ha	No	Agral® 90 Ag-Suri®

APPROVED TANK MIX PARTNERS IN DURUM WHEAT					
BROADLEAF HERBICIDE TANK MIX PARTNER		BROADLEAF HERBICIDE PRODUCT RATES per hectare	EVEREST SOLUPAK 70 DF Herbicide Rate***		Surfactants (0.25% v/v or 0.25L/100L)
			Wild Oat Green Foxtail	Green Foxtail ONLY	
1	2,4-D Amine	420 g ae/ha	43 g/ha	21.5 g/ha	Agral® 90 Ag-Suri®
2	2,4-D Ester	420 g ae/ha	43 g/ha	21.5 g/ha	Agral® 90 Ag-Suri®
3	Frontline™ Herbicide Tank Mix	100 ml/ha Frontline A + 0.84 L/ha Frontline B	43 g/ha	No	Agral® 90 Ag-Suri®
4	Spectrum™ Herbicide Tank Mix	100 ml/ha Spectrum A + 1.5 L/ha Spectrum B	43 g/ha	No	Agral® 90 Ag-Suri®

\* Reduction in wild oat control may be observed with this tank-mix partner.

\*\* Addition of a second surfactant is not required for EVEREST SOLUPAK 70 DF Herbicide tank mixes with the herbicides Ally, Refine Extra and Unity Herbicide Tank Mix. Use only one of the recommended surfactants from the EVEREST SOLUPAK 70 DF Herbicide label at an application rate of 0.25% v/v (0.25 L per 100 L total spray solution).

\*\*\* Apply specified dosage in at least 100 L of water per hectare (45 L/acre or 10 gallons/acre).

## MIXING INSTRUCTIONS

The enclosed packets containing EVEREST SOLUPAK 70 DF Herbicide are **water-soluble** and will dissolve completely in water. After opening the outer bag, drop the required number of unopened inner packets into the spray tank as directed. Reseal outer bag to protect remaining packets. Do not excessively handle water-soluble packets or expose to moisture since this may cause breakage. Do not allow packets to become wet prior to mixing spray solution. Do not handle with wet hands.

## CAUTION

Do not use packets in liquid fertilizer. Do not use packets in a tank-mix with products that contain boron or release free chlorine. The resultant reaction of water-soluble packets and boron or free chlorine is a plastic which is not soluble in water or solvents such as diesel oils, kerosene, gasoline or alcohol.

Ensure the spray tank is clean. In-line strainers and nozzle screens should be clean and 50-mesh or coarser.

1. Fill the spray tank 1/3 to 1/2 full with clean water and begin agitation or bypass.
2. Add the required number of unopened packets of EVEREST SOLUPAK 70 DF Herbicide, as determined under "Recommended Rates", directly to the spray tank. Vigorous agitation is required to dissolve the EVEREST SOLUPAK 70 DF Herbicide. Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within approximately 5 to 10 minutes from the time they were added to the water. Check bottom of tank to ensure the EVEREST SOLUPAK 70 DF Herbicide is completely dissolved. Maintain sufficient agitation during both mixing and application.
3. Add the broadleaf weed herbicide.
4. Add the surfactant, then complete filling tank with balance of water needed.

### SPRAYING INSTRUCTIONS

Apply in a spray volume of 50 - 100 L/ha (22.5 - 45 L/acre or 5 - 10 gallons/acre) - unless otherwise specified in tank-mix partner section of this label - at 207-345 kPa (30-50 PSI) pressure to ensure proper weed coverage. Flat fan nozzles of 80° or 110° are recommended for optimum coverage. Do not use floodjet or control droplet application equipment. Nozzles may be oriented 45° forward to enhance crop penetration and to give better weed coverage.

### ROTATIONAL CROPS

The following crops may be planted 11 months after an application of EVEREST SOLUPAK 70 DF Herbicide.

<b>Soil Zones and Rotational Crops</b>			
<b>Gray-Wooded</b>	<b>Black</b>	<b>Dark Brown</b>	<b>Brown</b>
Spring wheat Barley Canola Field peas*	Spring wheat Barley Canola Field peas* Field bean Smart Canola (also called imitolerant or Clearfield canola) Flax Durum wheat	Spring wheat Barley Canola Field peas* Flax Durum wheat	Spring wheat

\*Field peas may be grown the year following EVEREST SOLUPAK 70 DF Herbicide application in fields where precipitation has been normal or above normal (10 year average) during the growing season, and where organic matter content is above 4% and pH below 7.5.

Or, fields can be summer fallowed.

Until further rotational studies are complete, do not plant crops other than those listed the year following application of EVEREST SOLUPAK 70 DF Herbicide on wheat.

### SPRAYER CLEAN-UP



1. Drain the tank and thoroughly rinse spray tank, boom and hoses with clean water. Pay particular attention to flushing out any visible deposits.
2. Fill the tank with clean water and 1% v/v (1 L/100L) household ammonia. Flush the hoses, boom and nozzles with the cleaning solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more, then drain the tank;
3. Clean nozzles and screens in a separate container using the ammonia and water;
4. Repeat #2;
5. Rinse tank, boom and hoses with clean water.
  - Do not clean sprayer near desirable vegetation, wells, or other water sources;
  - Dispose of all rinsings in accordance with provincial regulations;
  - Check tank-mix partner label for any additional clean-up procedures.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, EVEREST SOLUPAK 70 DF Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to EVEREST SOLUPAK 70 DF Herbicide and other Group 2 wild oat herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of EVEREST SOLUPAK 70 DF Herbicide or other Group 2 wild oat herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact UPL AgroSolutions Canada Inc at 1-800-438-6071.

## **PRECAUTIONS**

### **KEEP OUT OF REACH OF CHILDREN**

#### **PROTECTIVE CLOTHING**

When handling (mixing and loading) EVEREST SOLUPAK 70 DF Herbicide, wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.

Read and follow handling precautions and protective clothing recommendations on broadleaf herbicide and surfactant tank-mix partner labels. Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep PPE apart from other laundry and wash separately.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside, then wash thoroughly and put on clean clothing. Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change in to clean clothing.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

#### **FIRST AID**

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

#### **TOXICOLOGICAL INFORMATION**

No specific antidote is available. Treat the patient symptomatically.

#### **STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original container and

out of reach of children, preferably in a locked storage area. EVEREST SOLUPAK 70 DF Herbicide is not affected by freezing; however, the water-soluble packet becomes brittle and may break if handled roughly in a frozen condition. Avoid physical damage to EVEREST SOLUPAK 70 DF Herbicide water-soluble packets during storage and prior to mixing in spray tank.

**SPILL CLEAN-UP:** Follow safety precautions as directed for handling the product. Sweep spilled product with a broom and shovel into tightly closable container. If on a floor or hard surface, wash the surface or floor with detergent and water, then rinse. If on soil, collect surface soil contaminated with the product. If the product is clean, it may be used; otherwise, follow the DISPOSAL instructions.

**DISPOSAL:** Do not contaminate water, food or feed by storage or disposal.

1. After the outer bag is empty, make it unsuitable for further use.
2. Dispose of the outer bag in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

<p><b>FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.</b></p>
--

UPL, the UPL logo and EVEREST are registered trademarks of a UPL Corporation Limited Group Company.

©2021 UPL Corporation Limited Group Company

SOLUPAK is a registered trademark of Bayer CropScience Inc. used under license by Arysta LifeScience North America LLC.

®All other products are registered trademarks of their respective companies.

***LABELLING FOR WATER-SOLUBLE PACKET***

---

**EVEREST<sup>®</sup> SOLUPAK<sup>®</sup> 70 DF**  
**70% Water Dispersible Granular Herbicide**  
**In Water-Soluble Packets**

**ACTIVE INGREDIENT: Flucarbazone at  
66%(present as flucarbazone-sodium)**

**Warning, contains the allergen milk**

**REGISTRATION NO. 26448 PEST CONTROL PRODUCTS ACT**

**KEEP OUT OF REACH OF CHILDREN**

**Net Contents: 174 Grams**

---

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.