

09-MAY-2007

(# 26459 FRONT PANNEL)

SUPERIOR control products

Paraffinized pellets

For indoor and outdoor use

DOMESTIC

For control of:

House mice, Norway rats,

Meadow voles, Ground squirrels

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC PETS

RATOL

REGISTRATION NUMBER 26459

PEST CONTROL PRODUCT ACT

GUARANTEE: Chlorophacinone 0.005%

WARNING, contains the allergens wheat, peanuts and soya

NET CONENTS:

454g to 2Kg

SUPERIOR control products inc.

3715 La Fayette est, Boisbriand, Québec, Canada, J7H 1S4, (888) 689-2171

## **(#26459)RATOL side panel**

**USE LIMITATIONS :** Bait **MUST** either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

### **DIRECTIONS FOR USE:**

**FOR CONTROL OF RATS AND MICE:** Place baits stations near nest feeding areas and water supplies. Place RATOL paraffinized pellets bait behind boards next to wall, in rat holes, between walls or other protected areas. Replace baits that are contaminated, spoiled or not fresh. Replenish baits supply as long as there is evidence of rat population. Protect bait against access of children, pets, livestock and wildlife. **FOR RATS:** Each station should contain at least 100g of bait: maintain constant supply of bait at stations for at least 10 days. **FOR HOUSE MICE:**Place 10-15g bait (one tablespoon) at distance of 3 to 4 meters intervals an uninterrupted supply of bait should be maintained for at least 15 days. **FOR MEADOW VOLES (Microtus SP):** FALL PERIOD: Supply with RATOL paraffinized pellets preferably before snowfall and leaves fall or lodging of grass. To protect your fruit trees and ornaments scatter a handful of pellets (50-100g) around tree (do not leave in piles). Using bait brush piles away from trees can help reduce over wintering population of meadow voles. SPRING PERIOD: Fill up the holes with RATOL and check daily in order to keep a constant feeding (do not replace uneaten baits). **FOR GROUND SQUIRREL:** Individual treatment of ground squirrel holes reduce the possibility of secondary poisoning and increases the degree of control. Locate all the ground squirrel holes and place 6 RATOL paraffinized pellets into each hole. Revisit the site within 48 hours and re-bait all holes where bait has been consumed or removed. Do not re-bait holes where bait has been undisturbed. A third visit may be needed for full control.

**BAIT STATION CONTROL:** Determine the approximate population present by visual count over a two day period. Establish protected bait stations or tamper resistant boxes every 30 to 60 meter in areas where the ground squirrel problem is present (use 1 box for every group of 50 animals counted). Use 500g per station and check daily and maintain an uninterrupted supply for 3 weeks or until feeding has stopped. To ensure full control, cover any ground squirrel hole and check the following day for opening to determine whether the hole is active. Place 6 RATOL paraffinized pellets in each active hole.

### **PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.**

**DO NOT MANIPULATE WITH YOUR BARE HANDS.** May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** Keep out of lakes, streams or ponds.

**FIRST AID:** If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give 1 to 2 glasses of water to patient and induce vomiting by placing finger down throat. Repeat until vomit is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** If swallowed or absorbed, this material may reduce the clotting ability of blood and cause bleeding. Vitamin K1 in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**DISPOSAL:** Do not reuse container. Dispose of empty container with household garbage.

**RODENT DISPOSAL:** Dispose of dead rodents in garbage or by burying.

**STORAGE:** Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals. Keep away from fire, open flame or other source of heat.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.