

GROUP 2 HERBICIDE

Thimax™

HERBICIDE

DRY FLOWABLE

FOR SALE FOR USE ON SOYBEANS AND FIELD TOMATOES ONLY

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Thifensulfuron methyl 75%

REGISTRATION NO. 29627 PEST CONTROL PRODUCTS ACT

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

NET CONTENTS: 64 GRAMS
(8 x 8 grams water-soluble bags)

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- AVOID BREATHING SPRAY MIST.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
- KEEP WATER SOLUBLE BAGS IN ORIGINAL OUTER BAG AND AWAY FROM WATER.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY.
- DO NOT APPLY BY AIR
- WEAR LONG-SLEEVED SHIRT, LONG PANTS, SOCKS, AND CHEMICAL-RESISTANT GLOVES AND FOOTWEAR DURING MIXING, LOADING,

APPLICATION, CLEAN-UP AND REPAIR. CHEMICAL-RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS.

FIRST AID:

EYE CONTACT:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice

SKIN OR CLOTHING CONTACT:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

INGESTION:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

INHALATION:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call **1-800-331-3148** (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL HAZARDS:

- TOXIC to plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).
- Avoid application of this product when heavy rain is forecast.

- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE:

Store product in original container in a secure, dry area away from pesticides, fertilizers, food or feed. Not for use or storage in or around the home. Keep container tightly closed and dry. Keep water-soluble bags in original outer bag away from moisture.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the empty container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use this product in a way that is inconsistent with the directions on the label.

®™ Thimax is a Registered trademarks/trademarks of FMC Corporation or an affiliate.

<DFU Booklet>

GROUP 2 HERBICIDE

Thimax™

HERBICIDE

DRY FLOWABLE

FOR SALE FOR USE ON SOYBEANS AND FIELD TOMATOES ONLY

AGRICULTURAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

ACTIVE INGREDIENT: Thifensulfuron methyl 75%

REGISTRATION NO. 29627 PEST CONTROL PRODUCTS ACT

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

NET CONTENTS: 64 GRAMS
(8 x 8 grams water-soluble bags)

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- AVOID BREATHING SPRAY MIST.

- AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
- KEEP WATER SOLUBLE BAGS IN ORIGINAL OUTER BAG AND AWAY FROM WATER.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY.
- DO NOT APPLY BY AIR
- WEAR LONG-SLEEVED SHIRT, LONG PANTS, SOCKS, AND CHEMICAL-RESISTANT GLOVES AND FOOTWEAR DURING MIXING, LOADING, APPLICATION, CLEAN-UP AND REPAIR. CHEMICAL-RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS.

FIRST AID:

EYE CONTACT:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

SKIN OR CLOTHING CONTACT:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

INGESTION:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

INHALATION:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

APPLICATION PRECAUTIONS:

DO NOT APPLY BY AIR.

Do not re-enter treated fields until 12 hours after application.

Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts or similar areas.
- Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops.
- Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than soybeans.
- Do not handle pesticides with bare hands. Chemical-resistant gloves significantly reduce hand exposure. Wear chemical-resistant gloves for mixing/loading/clean-up operations and when making sprayer and nozzle repairs and adjustments. Do not use leather or cloth gloves.

PRODUCT DESCRIPTION:

Thimax™ is recommended for selective postemergence control of broadleaf weeds in soybeans and field tomatoes. Thimax™ is in the form of a dry flowable granule to be mixed in water and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile, and does not freeze.

Thimax™ must be applied with a recommended surfactant early postemergence to the main flush of actively growing weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of Thimax™ by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress may not be adequately controlled or suppressed and regrowth may occur. Thimax™ may only be applied using ground equipment. For best results ensure thorough spray coverage of target weeds. Refer to "RATES AND TIMING" section of the label for complete use details.

Thimax™ rapidly stops growth of susceptible weeds. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Favourable growing conditions following treatment promote the activity of Thimax™ while cold, dry conditions delay the activity.

Degree of control and duration of effect depend on weed sensitivity, weed size, crop competition, growing conditions and spray coverage.

Thimax™ may be applied on all soil types.

ENVIRONMENTAL HAZARDS:

- **TOXIC** to plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

DIRECTIONS FOR USE

- Thimax™ is formulated as a wettable granule or a soluble granule sealed within a water-soluble package (WSP). Drop intact WSP directly into spray tank. The WSP and pesticide will dissolve readily in water.
- Do not allow the WSP to become wet prior to use. Do not handle individual WSP with wet hands or wet gloves as this may cause breakage. Do not remove WSP from overwrap container except for immediate use.
- DO NOT open WSP or attempt to remove the contents from them. The contents are not designed to be re-measured or subdivided
- Do not use opened or punctured WSP.
- If broken WSP is found when the over-wrap bag is opened, avoid contact with, and inhalation of the product. Handle with chemical-resistant gloves and dispose of broken WSP according to the DISPOSAL section.

Field sprayer application:

- DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty.
- DO NOT apply with spray droplets smaller than the ASAE medium classification.

Buffer zones:

Use of the following spray methods or equipment DO NOT require a buffer zone: hand-held or backpack sprayer, inter-row hooded sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands, and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Buffer zone (metres) required for the protection of:		
	Terrestrial habitat	Aquatic habitats of water depths:	
		<1m	>1m
Field sprayer*	15	1	0

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American

Society of Agricultural Engineers (ASAE) [fine medium or coarse] classification.' To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

Thimax™ USED ALONE

SPRAY ADJUVANTS:

Thimax™ must be applied with a recommended non-ionic surfactant, such as AG-SURF*, AGRAL* 90, or CITOWETT* PLUS at 1 litre per 1000 litres water (0.1% v/v) OR a crop oil concentrate (if applicable), such as SURE-MIX* at 0.5% v/v (0.5 litres per 100 litres of spray solution) or ASSIST* at 1-2 litres per hectare.

RATES AND TIMING:

Thimax™ may be applied anytime after the soybeans have the first trifoliolate leaf fully expanded and before soybeans have initiated flowering. Thimax™ can be applied up to 60 days before soybean harvest.

Use 5.5 or 8 grams Thimax™ per hectare. For best results apply to young actively growing weeds before the soybean canopy closes. Weeds should be less than 10 cm tall or across. Weeds that emerge after treatment will not be controlled.

NOTE:

If rain occurs after application, control may be reduced. Several hours of dry weather are needed to allow Thimax™ to be absorbed by weed foliage.

Apply 5.5 or 8 grams Thimax™ per hectare as a broadcast spray with a recommended surfactant.

WEEDS CONTROLLED AT 5.5 GRAMS PER HECTARE:

- Redroot pigweed
- Wild mustard
- Lady's thumb

One 64 gram container of Thimax™ treats 11.5 ha (28.5 acres) at 5.5 grams/ha.

WEEDS CONTROLLED AT 8.0 GRAMS PER HECTARE:

- Weeds controlled at 5.5 grams per hectare plus:
- Lamb's-quarters

Velvetleaf †

†The addition of a nitrogen-containing fertilizer such as 28% N, at 4 litres per 100 litres of spray solution, or an equivalent amount of nitrogen as urea (46-0-0) per ha may improve control of velvetleaf.

One 64 gram container of Thimax™ treats 8 ha (20 acres) at 8 grams/ha.

Thimax™ TANK MIXES

Thimax™ + ASSURE* II

For the control of annual grassy and specific broadleaf weeds, apply a tank mixture of the following:

Thimax™	5.5 or 8.0 g/ha
plus	
ASSURE* II	0.5 L/ha
plus	
SURE-MIX*	0.5% v/v (0.5 litres per 100 litres of spray solution)

Consult the complete ASSURE* II Directions for Use pamphlet for grassy weeds controlled. For the correct use rate for the broadleaf weeds present, refer to the Weeds Controlled section in this label.

Consult the timing section of this label for the correct timing of application to grassy weeds. For optimum leaf stage timing of broadleaf weeds, consult the General Information section of this label. If leaf stages of the grass and broadleaf weeds do not coincide, a sequential application of the grass and broadleaf herbicides is required to ensure satisfactory control. Do not apply before the crop has 1-2 full trifoliate leaves.

Velvetleaf control may be reduced with a single tank mix application. For optimum control, make separate applications of Thimax™ and ASSURE* II Herbicides.

Consult the ASSURE* II label for precautions, weeds controlled and additional restrictions on the use of ASSURE* II.

Thimax™ + ASSURE* II + BASAGRAN* FORTE

For the control of annual grassy and broadleaf weeds, apply a tank mixture of the following:

Thimax™	5.5 or 8.0 g/ha
plus	
ASSURE* II	0.63 L/ha
plus	
BASAGRAN* FORTE	1.75 or 2.25 L/ha
plus	
SURE-MIX*	0.5% v/v (0.5 litres per 100 litres of spray solution)

Consult the BASAGRAN* FORTE label and the above table for the correct use rate for the broadleaf weeds present. Weeds controlled by 0.63 L/ha of ASSURE* II in this tank mix are listed on the ASSURE* II label at 0.5 L/ha (when ASSURE* II is used alone). The increase in the application rate of ASSURE* II in this tank mix is required for consistent grassy weed control.

Consult the ASSURE* II label for the correct timing of application to grassy weeds. For optimum leaf stage timing of broadleaf weeds, consult the BASAGRAN* FORTE label and as well as the General Information and Directions for Use sections in this label. If leaf stages of the grass and broadleaf weeds do not coincide, a sequential application of the grass and broadleaf herbicides is required to ensure satisfactory control. Do not apply before the crop has 1-2 full trifoliolate leaves.

Consult the ASSURE* II and BASAGRAN* FORTE labels for precautions, weeds controlled and additional restrictions.

Thimax™ plus BASAGRAN* or BASAGRAN* FORTE tank mix

Thimax™ may be tank-mixed with BASAGRAN* or BASAGRAN* FORTE to control additional weeds not controlled by Thimax™ alone. If BASAGRAN* FORTE is used, an additional adjuvant is not required. This tank mix includes:

Thimax™	8.0 g/ha
plus	
BASAGRAN* FORTE	1.75 or 1.25 L/ha

Thimax™	8.0 g/ha
plus	
BASAGRAN*	1.75 or 2.25 L/ha

plus	
ASSIST*	1.0 – 2.0 L/ha

The tank mixture will control the weeds listed for control by Thimax™ alone (at 8 g/ha), plus the weeds listed on the BASAGRAN* or BASAGRAN* FORTE label. Please refer to the BASAGRAN* or BASAGRAN* FORTE label for weeds controlled at the appropriate use rate, as well as additional application instructions and use precautions.

MIXING INSTRUCTIONS:

1. Make sure that spray tank is thoroughly cleaned before mixing.
2. Fill tank half full with water. KEEP AGITATOR RUNNING.
3. Add the required amount of Thimax™ to the tank with the agitator running. Continuous agitation is required to keep the product in suspension. Avoid overfilling of spray tank.
4. After Thimax™ has been thoroughly mixed and is in suspension, add the proper amount (if applicable) of ASSURE* or ASSURE* II, BASAGRAN* or BASAGRAN* FORTE to the spray tank with the agitator running.
5. After Thimax™ (and tank mix herbicide if applicable) has been well mixed and is in suspension, add a recommended non-ionic surfactant, such as AG-SURF*, AGRAL* 90, or CITOWETT* PLUS at 1 litre per 1000 litres water (0.1% v/v) OR a crop oil concentrate (if applicable), such as SURE-MIX* at 0.5% v/v (0.5 litres per 100 litres of spray solution) or ASSIST* at 1-2 litres per hectare.
6. The addition of a nitrogen containing fertilizer such as 28% N, at 4 litres per 100 litres of spray solution, or an equivalent amount of nitrogen as urea (46-0-0) per ha may improve control of velvetleaf and should be added after the surfactant.
7. Add the rest of the required water to the tank. Mix well before applying to the crop.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of volume about to be mixed.

Agitation is required for uniform mixing and application. Apply Thimax™ in a minimum of 100 litres of water per hectare. Use spray preparation of Thimax™ within 24 hours or product degradation may occur resulting in a

loss of weed control. Use vigorous agitation to thoroughly disperse spray mixtures that have been allowed to stand in the tank.

Calibrate sprayer only with clean water away from well site. Make scheduled checks of spray equipment.

MINOR USES

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than FMC and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. FMC itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) of this product when used on the crops listed below. Accordingly, the User assumes all risk related to performance and crop tolerance arising, and agrees to hold FMC harmless from any claims based on efficacy or phytotoxicity in connection with the use described below.

FOR SALE FOR USE ON FIELD TOMATOES ONLY

For control of lady's thumb, redroot pigweed and wild mustard apply 5.5 grams Thimax™ per hectare with a recommended non-ionic surfactant (AGRAL* 90 only) at 2.0 litres per 1,000 litres of spray solution (0.2% v/v).

For control of the weeds listed above, together with lamb's-quarters and velvetleaf apply 8.0 grams Thimax™ per hectare with a recommended non-ionic surfactant (AGRAL* 90 only) at 2.0 litres per 1,000 litres of spray solution (0.2% v/v).

Apply Thimax™ to field tomatoes three (3) weeks after transplanting in the field. Weeds should be less than 10 cm tall or across. Weeds that emerge after treatment will not be controlled.

FOR SALE FOR USE ON PROCESSING FIELD TOMATOES ONLY

Thimax™ (5.5 g/ha) + PRISM* Herbicide 25% Dry Flowable (60 g/ha)

This tank mixture controls the following weeds:

Red Root Pigweed	Wild Mustard	Lady's Thumb
Green Foxtail		Fall Panicum
Hairy Nightshade	Lamb's-quarters*	Barnyard grass

*Suppression only

Thimax™ (8 g/ha) + PRISM* Herbicide 25% Dry Flowable (60 g/ha)

**This tank mixture controls the following weeds:
Weeds listed above plus Lamb's-quarters and Velvetleaf.**

Tank mix PRISM* Herbicide 25% Dry Flowable (60 g/ha) and Thimax™ (5.5 g/ha or 8 g/ha) and apply as a single post-emergent spray using ground application only. Apply to weeds up to 10 cm or 21 days after transplanting. Use with a non-ionic surfactant, Agral*90 at 0.2% v/v. Use a Spray Volume of 200L/ha.

Do not harvest crop within 45 days of treatment. Refer to the appropriate sections of the PRISM* Herbicide 25% Dry Flowable and Thimax™ labels for recropping information, and for additional application instructions, specific weed recommendations and/or use precautions.

NOTE:

If rain occurs after application, control may be reduced. Several hours of dry weather are needed to allow Thimax™ to be absorbed by weed foliage.

Thimax™ can be applied 45 days before harvest of field tomatoes.

Please refer to the appropriate sections of the Thimax™ label for additional use precautions, application and/or mixing instructions.

SPRAYER CLEANUP:

CLEANING PRECAUTIONS:

CAUTION: DO NOT use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour, which may cause eye, nose, throat and lung irritation. DO NOT clean equipment in an enclosed area.

Extreme care should be taken to avoid damage by drift onto plants outside the target area or onto ponds, waterways or ditches or land intended for cropping. When spraying near adjacent sensitive crop or plants, do everything possible to reduce spray drift.

CLEANING INSTRUCTIONS:

To avoid subsequent injury to crops other than soybeans and field tomatoes, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of Thimax™ from mixing and spray equipment as follows:

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to assure removal of all visible residues of Thimax™. If necessary, repeat step 1. **DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.**
2. Fill tank with clean water while adding 1 litre household ammonia (containing a minimum 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

USE PRECAUTIONS:

- Do not apply to soybeans or field tomatoes that are stressed by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage as crop injury may result.
- Do not make more than one application per growing season.
- Do not apply to irrigated land where tail water will be used to irrigate crop land. Do not contaminate irrigation water.
- Under certain conditions (such as heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures), temporary

lightening in crop colour and occasionally a slight reduction in crop height may occur.

- Because crop varieties differ in their tolerance to herbicides, limit first use of Thimax™ to a small area of each variety prior to adoption as a field practice.
- Do not mix, load, or apply within 15 meters of all wells, bodies of water and wetland areas.

CAUTION:

Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, Thimax™ is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Thimax™ and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of Thimax™ or other Group 2 herbicides with different groups that control the same weed in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted.
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local FMC representative or the FMC hotline at 1-833-362-7722 for further information.

STORAGE:

Store product in original container in a secure, dry area away from pesticides, fertilizers, food or feed. Not for use or storage in or around the home. Keep container tightly closed and dry. Keep water soluble bags in original outer bag away from moisture.

DISPOSAL:

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.

4. Dispose of the empty container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact FMC at 1-833-362-7722 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by FMC

®/™ Thimax is a Registered trademarks/trademarks of FMC Corporation or an affiliate.

* All other products mentioned are trademarks of their respective companies.

<<Water Soluble Bag>>

Thimax™

KEEP OUT OF REACH OF CHILDREN

SEE DIRECTIONS FOR USE BROCHURE FOR COMPLETE PRECAUTIONARY
STATEMENTS

ACTIVE INGREDIENT: Thifensulfuron Methyl 75%

DRY FLOWABLE

NET CONTENTS: 8 GRAMS