

** BASE CONTAINER LABEL **

GROUP	4	INSECTICIDE
-------	---	-------------

NIPSIT INSIDE® 600 INSECTICIDE
SUSPENSION

NIPSIT INSIDE 600 INSECTICIDE is a systemic insecticide seed treatment for use on labeled crops for the control of listed insect pests

COMMERCIAL

ACTIVE INGREDIENT: Clothianidin 600 g/L

Contains 1,2-benzisothiazolin-3-one at 0.04%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.001% and 2-methyl-4-isothiazolin-3-one at 0.0004% as preservatives.

OR

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0006% and 2-methyl-4-isothiazolin-3-one at 0.00019% as preservatives OR Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0017% and 2-methyl-4-isothiazolin-3-one at 0.0006% as preservatives.

OR

Contains 1,2-benzisothiazolin-3-one as a preservative at 0.0483%

REGISTRATION NUMBER 28975 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

WARNING

<<Skull and Cross Bones in Diamond>>

POISON

NET CONTENTS: 20 - 1200 L

This product contains no colourant. An appropriate colourant must be added when this product is applied. Regulations pertaining to the "Seeds Act" must be strictly adhered to when using this product. A baby blue colourant must be added when this product is applied to canola/rapeseed.

HEALTH EMERGENCY OR SPILL (24 hr):

(800) 682-5368

TRANSPORTATION (24 hr.): CHEMTREC

(800) 424-9300 or (202) 483-7616

GROWER INFORMATION:

Valent Canada, Inc.

201-230 Hanlon Creek Blvd., Guelph, Ontario, N1C 0A1 Contact Phone Number: (519) 767-9262

For canola, rapeseed, carinata and corn: For use with commercial seed treaters (facilities and mobile treaters) with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. No open transfer of NIPSIT INSIDE 600 INSECTICIDE.

For wheat: For use in commercial seed treatment facilities (with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only) and for use on-farm (open transfer including open mixing, loading calibrating, and open treatment equipment is allowed).

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if swallowed. Harmful if absorbed through skin. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Seed Treatment Workers: Workers mixing, loading, treating or maintaining and repairing seed treatment equipment, bagging, sewing or stacking treated seed must wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, head gear, and respiratory protection (i.e., NIOSH - approved dust-filtering respirator or fresh air hood) and socks and shoes. While cleaning treating equipment workers must wear chemical resistant coveralls over long-sleeved shirt, long pants,

chemical-resistant gloves, head gear, and respiratory protection (i.e., NIOSH - approved dust-filtering respirator or fresh air hood) and socks and shoes.

Planters must use a closed cab planter when planting treated wheat seeds.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.
- Follow manufacturer's instructions for cleaning/maintaining. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the US, contact Valent Canada, Inc.

ENVIRONMENTAL PRECAUTIONS:

Use restrictions (wheat; all other CG15 cereal seeds excluding corn): Additionally, wheat and all treated CG 15 cereal seed (excluding corn) for sale or use in Canada must be labeled with the following information: Clothianidin is toxic to bees. Dust generated during planting of treated seed may be harmful to bees and other pollinators. To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators

When using a seed flow lubricant with this treated seed, only a dust reducing fluency agent is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for corn seed treated with these insecticides. Carefully follow use directions for this seed flow lubricant.

Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds.

When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.

Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatments. When used according to label directions minimal exposure or risk is expected.

Toxic to aquatic organisms. Treated seed is toxic to birds and small wild mammals.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact Valent Canada, Inc.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE: Store in a cool place, away from food and feed. Do not store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

RECYCLABLE CONTAINER:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER:

1. After use, return the container to the point of purchase or designated locations.

2. This container must only be refilled with NIPSIT INSIDE 600 INSECTICIDE at designated locations.
3. Do not reuse container for other purposes.
4. Prior to refilling inspect thoroughly for damage.
5. Check for leaks after refilling and before transportation.
6. Do not refill or transport damaged or leaking containers.
7. If the container is not being refilled, return to the point of purchase.

NON-RECYCLABLE/RETURNABLE/REFILLABLE CONTAINER:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

NIPSIT INSIDE® is a registered trademark of Valent U.S.A. LLC

** BOOKLET**

GROUP	4	INSECTICIDE
-------	---	-------------

NIPSIT INSIDE™ 600 INSECTICIDE
SUSPENSION

NIPSIT INSIDE 600 INSECTICIDE is a systemic insecticide seed treatment for use on canola, rapeseed, carinata, corn, and wheat for the control of listed insect pests. As a potato seed-piece treatment, NipsIt INSIDE 600 INSECTICIDE controls listed insect pests and provides suppression of wireworms.

COMMERCIAL

ACTIVE INGREDIENT: Clothianidin 600 g/L

Contains 1,2-benzisothiazolin-3-one at 0.04%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.001% and 2-methyl-4-isothiazolin-3-one at 0.0004% as preservatives.

OR

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0006% and 2-methyl-4-isothiazolin-3-one at 0.00019% as preservatives OR Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0017% and 2-methyl-4-isothiazolin-3-one at 0.0006% as preservatives.

OR

Contains 1,2-benzisothiazolin-3-one as a preservative at 0.0483%

REGISTRATION NUMBER 28975 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

WARNING

<<Skull and Cross Bones in Diamond>>

POISON

NET CONTENTS: 20 - 1200 L

This product contains no colourant. An appropriate colourant must be added when this product is applied. Regulations pertaining to the "Seeds Act" must be strictly adhered to when using this product. A baby blue colourant must be added when this product is applied to canola/rapeseed.

HEALTH EMERGENCY OR SPILL (24 hr):

(800) 682-5368

TRANSPORTATION (24 hr.): CHEMTREC

(800) 424-9300 or (202) 483-7616

GROWER INFORMATION:

Valent Canada, Inc.

201-230 Hanlon Creek Blvd., Guelph, Ontario, N1C 0A1 Contact Phone Number: (519) 767-9262

For canola, rapeseed, carinata and corn: For use with commercial seed treaters (facilities and mobile treaters) with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. No open transfer of NIPSIT INSIDE 600 INSECTICIDE.

For wheat: For use in commercial seed treatment facilities (with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only) and for use on-farm (open transfer including open mixing, loading calibrating, and open treatment equipment is allowed).

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if swallowed. Harmful if absorbed through skin. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Seed Treatment Workers: Workers mixing, loading, treating or maintaining and repairing seed treatment equipment, bagging, sewing or stacking treated seed must wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, head gear, and respiratory protection (i.e., NIOSH - approved dust-filtering respirator or fresh air hood) and socks and shoes. While cleaning treating equipment workers must wear chemical resistant coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, head gear, and respiratory protection (i.e., NIOSH - approved dust-filtering respirator or fresh air hood) and socks and shoes.

Planters must use a closed cab planter when planting treated wheat seeds.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.
- Follow manufacturer's instructions for cleaning/maintaining. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., [contact](#) Valent Canada, Inc.

ENVIRONMENTAL PRECAUTIONS:

Clothianidin is toxic to bees. Dust generated during planting of treated seed may be harmful to bees and other pollinators.

To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators

When using a seed flow lubricant with this treated seed, only a dust reducing fluency agent is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for corn seed treated with these insecticides. Carefully follow use directions for this seed flow lubricant.

Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds.

When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.

Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatments. When used according to label directions minimal exposure or risk is expected.

Toxic to aquatic organisms. Treated seed is toxic to birds and small wild mammals.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact Valent Canada, Inc.

USE RESTRICTIONS:

- Do not use treated seed for food, feed, or oil processing.
- Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed.
- Rape greens and seed grown or harvested from NIPSIT INSIDE 600 INSECTICIDE-treated seed must not be used for feed or human consumption.
- Rapeseed and carinata grown and harvested from NIPSIT INSIDE 600 INSECTICIDE-treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.

LABELLING OF TREATED SEED:

All bags containing treated seed must be labeled or tagged as follows:

- This seed was treated with a product containing the active ingredient clothianidin.
- Do not use for feed, food or oil processing.
- Rape greens and seed grown or harvested from NIPSIT INSIDE 600 INSECTICIDE-treated seed must not be used for feed or human consumption.
- Store away from feeds and other foodstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- This product is toxic to aquatic invertebrates.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- Cover or incorporate spilled treated seeds.

- Planters must use a closed cab planter when planting treated wheat seeds.

All treated corn seed for sale or use in Canada must also be labeled with the following information:

- Clothianidin is toxic to bees. Dust generated during planting of treated seed may be harmful to bees and other pollinators.
- To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators.
- When using a seed flow lubricant with this treated seed, only a dust reducing fluency agent is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for corn seed treated with these insecticides. Carefully follow use directions for this seed flow lubricant.
- Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds.
- When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.
- Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

Additionally, wheat and all treated CG 15 cereal seed (excluding corn) for sale or use in Canada must be labeled with the following information:

- Clothianidin is toxic to bees. Dust generated during planting of treated seed may be harmful to bees and other pollinators.
- To help minimize the dust generated during planting, refer to the “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators.
- Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds.
- When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.
- Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE: Store in a cool place, away from food and feed. Do not store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

RECYCLABLE CONTAINER:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER:

1. After use, return the container to the point of purchase or designated locations.
2. This container must only be refilled with NIPSIT INSIDE 600 INSECTICIDE at designated locations.
3. Do not reuse container for other purposes.
4. Prior to refilling inspect thoroughly for damage.
5. Check for leaks after refilling and before transportation.
6. Do not refill or transport damaged or leaking containers.
7. If the container is not being refilled, return to the point of purchase.

NON-RECYCLABLE/RETURNABLE/REFILLABLE CONTAINER:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

SEED QUALITY AND STORAGE OF TREATED SEED

- Laboratory and field studies have shown that NIPSIT INSIDE 600 INSECTICIDE applied to seed will not adversely affect germination.
- Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed. Treatment of mechanically scarred or damaged seed, or of seed known to be of low germ, low vigour and poor quality is not recommended.
- Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed.

DIRECTIONS FOR USE:

NIPSIT INSIDE 600 INSECTICIDE is a systemic insecticide seed treatment for use on canola, rapeseed, carinata, corn, and wheat for the control of listed insects, and for control of listed pests

and for suppression of wireworms in potatoes.

For use in commercially available equipment designed for seed treatment only. Not for use in hopper-box, slurry-box, or similar seed treatment applications used at planting. This product is to be used in liquid or slurry treaters. Mix thoroughly before use or use entire container at one time.

This product may be tank mixed with a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Valent Canada, Inc. at 519-767-9262 for information before applying any tank mix that is not specifically recommended on this label.

SPECIAL USE RESTRICTION:

This product contains no colourant. An appropriate colourant must be added when this product is applied. Regulations pertaining to the “Seeds Act” must be strictly adhered to when using this product. A baby blue colourant must be added when this product is applied to canola/rapeseed.

ROTATIONAL CROP RESTRICTIONS:

Corn, canola, rapeseed, carinata, wheat, and potatoes may be replanted at any time. A one year plantback interval is required for leafy, root and tuber vegetables (except potatoes). A 30-day plantback interval on cereal grains (except wheat), grasses, non-grass animal feeds and soybeans and dried beans is required.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that NIPSIT INSIDE 600 INSECTICIDE is a Group 4 insecticide. Any insect population may contain individuals naturally resistant to this product and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of NIPSIT INSIDE 600 INSECTICIDE or other Group 4 insecticides with different groups that control the same pests in a field.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact Valent Canada, Inc. at 1-800-682-5368.

APPLICATION INSTRUCTIONS

Crop	Pest	Rate of NIPSIT INSIDE 600 INSECTICIDE		Remarks
Canola, Rapeseed, carinata	Flea beetle	250 mL, 333 mL or 666 mL per 100 kg seed		<p>250 mL/100 kg seed: For use under low to moderate flea beetle pressure.</p> <p>333 mL/100 kg seed: For a higher level of control under moderate flea beetle pressure.</p> <p>666 mL/ 100 kg seed: For use under high to extreme flea beetle pressure where extended control is required.</p>
		mg AI/kernal	per 80,000 unit of seed	
Corn, field	Corn flea beetle Cutworm, black Seedcorn maggot Wireworm	0.25-0.5	33.3-66.6 mL	
	White grub (larvae of European Chafer, May/June beetle, Japanese beetle)	0.25	33.3 mL	
Corn (including sweet, and pop)	Corn rootworm (including Northern, Western)	1.25	166.7 mL	<p>The application rate for control of corn rootworm also provides control of other listed corn pests.</p> <p>If corn rootworm is not a target pest, use appropriate</p>
	Corn flea beetle Cutworm, black Seedcorn maggot Wireworm	0.25 – 0.5	33.3 – 66.6 mL	

	White grub (larvae of European Chafer, May/June beetle, Japanese beetle)	0.25	33.3 mL	lower application rates for control of other listed corn pests.
Crop	Pest	Rate of NIPSIT INSIDE 600 INSECTICIDE		Remarks
Wheat	Wireworm	17 mL per 100 kg seed (for suppression) 33 – 100 mL per 100 kg seed		Use higher rates on wheat seed to be planted into fields known to have a history of severe wireworm infestations for protection to seed and seedlings.
	Aphids	50 mL per 100 kg seed		For early season protection of wheat seedlings against aphid feeding.

Do not make any subsequent application of a Group 4 Insecticide (i.e., in-furrow or foliar application) following treatment with NIPSIT INSIDE 600 INSECTICIDE.

Crop	Pest	Rate of NIPSIT INSIDE 600 INSECTICIDE		Remarks
Potato (seed- piece treatment)	Wireworm (<i>Agriotes obscurus</i> , <i>A. lineatus</i> , <i>Limonius agonus</i> , <i>Melanotus spp.</i> , <i>M. communis</i> (damage suppression only.) May reduce the damage caused by other wireworm species.	20.8 mL per 100 kg potato seed pieces		Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Apply specified dosage as a diluted spray onto seed pieces using a shielded spray system that is well-contained and will prevent the loss of any liquid.
	Aphid (including potato, green peach, foxglove and buckthorn aphids) Colorado potato beetle Potato leafhopper Potato flea beetle (overwintered adults and	10.4-20.8 mL per 100 kg potato seed pieces		Agitate or stir spray solution as needed. For optimal insect control, good coverage of the seed pieces is required. Do not dilute with any more than 6 parts water to 1 part Nipsit Inside 600 Insectide. As part of the seed cutting and treating process,

	suppression of second generation)		<p>application of a fungicide registered for potato seed treatment and/or an inert absorbent ingredient is recommended to improve suberization.</p> <p>Plant seed pieces as soon as possible after cutting and treating.</p> <p>Do not apply any subsequent application of a Group 4 Insecticide following Nipsit Inside 600 Insecticide as a potato seed piece treatment.</p> <p>For extended residual control of pests other than wireworm, apply the higher rate.</p>
--	-----------------------------------	--	--

In preparing a mixture of NIPSIT INSIDE 600 INSECTICIDE for application to seed, measure proper volume of NIPSIT INSIDE 600 INSECTICIDE along with other seed treatment additives (i.e., seed colourant, registered seed treatments whose labels also allow tank mixing). Triple rinsate from the product container should be incorporated into the mixture and calculated into the total volume recommended for seed volume.

When adding rinsate from triple rinsing the container for disposal, it is recommended to use 2% of the container size 3 times to triple rinse.

Add the appropriate volume of rinse water to the container and swish the contents thoroughly. Empty the rinsate into the seed treater holding tank and repeat this process two more times making sure the total volume of rinsate does not exceed 6% of the container size. Make sure rinsate is thoroughly mixed with seed treatment before treating. CAUTION: DO NOT dilute with rinsate over 6% the total slurry volume. Excess seed wetness may result in seed handling difficulties.

NIPSIT INSIDE® is a registered trademark of Valent U.S.A. LLC