

13-OCT-2009  
2009-2965

**Nufarm**

**Ag-Surf**  
Liquid Spray Adjuvant

For Use With ROUNDUP, ALLY, REGLONE, and ROVRAL 50% W.P

**AGRICULTURAL**

**READ THE LABEL BEFORE USING**

**GUARANTEE:** Nonylphenoxy polyethoxyethanol 92%

**REGISTRATION NO: 27921 PEST CONTROL PRODUCTS ACT**

10 Litres

Nufarm Agriculture Inc.  
5507 1<sup>st</sup> Street SE  
Calgary, Alberta T2H 1H9  
1-800-868-5444

24 Hour Emergency Response Number 1-800-424-9300

**GENERAL INFORMATION:**

Ag-Surf is a surfactant for use with Roundup, Ally and Reglone herbicides, and Rovral 50% W.P fungicide, in situations where a non-ionic surfactant is called for (minimum or zero tillage systems surfactant additive, etc.). It is important to follow label directions for Roundup, Ally, and Reglone herbicides and Rovral fungicide, using the recommended rates in combination with Ag-Surf.

**PRECAUTIONS:**

KEEP OUT OF THE REACH OF CHILDREN.

Do not store or place near open flames or sources of heat. Do not apply more surfactant than recommended or excessive run-off may occur. Avoid contact with eyes.

**DISPOSAL:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple - or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**FIRST AID:**

**If in eyes**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If swallowed**, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:**

Treat symptomatically.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Product Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**DIRECTIONS:**

**With Roundup: Minimum/Zero-Tillage/Summerfallow:** To improve the effectiveness for post-emergent control of annual grasses and broadleaf weeds in Minimum/Zero-Till systems for wheat, oats, barley or on summerfallow, apply Ag-Surf at 350 mL/ha plus Roundup at 1.1 L/ha in 50 - 100 L/ha of clean water. See Minimum and Zero Tillage section of Roundup label for weeds controlled and other details of application.

**Quackgrass:** When applying the “Quackgrass Control” rate of 2.5 L/ha of Roundup and water volumes of 150 - 300 L/ha, add Ag-Surf at 0.5% (0.5 litres/100 litres of water) to maintain adequate surfactant concentration in spray solution. See “Quackgrass Control” section of Roundup label for other details of application.

**With Ally:** For maximum weed killing effect – after the addition of Ally to water in the tank add 200 mL of Ag-Surf for each 100 litres of spray solution (0.2% total spray volume). An antifoaming agent may be needed to reduce foaming.

**With Reglone:** To improve the effectiveness for crop desiccation, post-emergent weed control in non-crop land, chemical mowing and potato vine killing, add Ag-Surf at 1 L per 1000 Litres of spray where the addition of a wetting and spreading agent is recommended on the Reglone label.

**With Rovral 50% W.P.:** For use in rapeseed only. For the control of Sclerotinia Stem Rot add Ag-Surf at the rate of 0.5% of the final spray solution (500 mL in 100 litres).

J04R0B/

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.