

2022-0231

2022-01-31

Advantage™ II Large Dog
Insert (English)

<Elanco Logo>

Advantage™ II large dog

Flea Adulticide/Larvicide/Ovicide and Lice Adulticide

For Dogs 7 weeks and older weighing between 11 and 25 kg
Once-A-Month Topical Flea and Lice Treatment

Liquid

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS
Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING



POISON

EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 27582 P.C.P. ACT

One to Six 2.5 mL Tubes

Elanco Canada Limited
1919 Minnesota Court, Suite 401
Mississauga, Ontario L5N 0C9
1-800-265-5475

42

43 Advantage II large dog provides dual protection against fleas and lice. Advantage II
44 large dog kills 98-100% of the existing fleas on dogs within 1 day after treatment.
45 Protection against further flea infestation can last up to 4 weeks. Retreatment at
46 monthly intervals is required if reinfestation is continuous or if flea control is desired.
47 Fleas, eggs, and larvae are killed following contact with skin debris from an Advantage
48 II large dog treated pet. Advantage II large dog provides multi-stage flea control
49 effectively breaking all flea life-cycle stages for quick and lasting control of flea
50 populations. Advantage II large dog treats lice infestations on dogs.

51

52 Do not shampoo or immerse the animal in water immediately after treatment. The
53 animal can be shampooed, immersed in water or exposed to rain 3 days after
54 treatment. Advantage II large dog retains its ovicidal and larvicidal efficacy up to 4
55 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal
56 effect may be shortened by repeated shampooing, water immersion or exposure to rain.
57 Reapplication may be required after such activities.

58

59 The successive feeding activity of fleas on pets may elicit a hypersensitivity skin
60 disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage II
61 large dog rapidly kills fleas and may reduce the incidence of this condition. For effective
62 flea and lice control, treatment of the pet should be combined with sanitation of any area
63 used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use)
64 and wash the pet's bedding, living quarters and surrounding areas. If pest problems
65 persist, an insecticidal premise treatment may be required. A veterinarian should be
66 consulted when pest infestation continues to be a problem on the pet.

67

68 PRECAUTIONS:

69

70 HUMANS: KEEP OUT OF THE REACH OF CHILDREN.

71 Harmful if swallowed, or absorbed through the skin. Causes eye irritation. Do not get in
72 eyes. Avoid contact with skin or clothing. Wear rubber gloves during application of
73 this product. After handling or applying, wash hands (or any other skin that came in
74 contact with the product), with soap and water. Avoid contact with pets until dry.

75

76 PETS: Do not use on animals under 7 weeks of age. Care should be taken to prevent
77 exposure of other pets who may be in close physical contact with the treated animal.
78 Individual sensitivities, while rare, may occur after using ANY pesticide product for pets.
79 Smaller animals are more likely to experience an adverse reaction. Observe carefully. If
80 signs persist, or become more severe, consult a veterinarian immediately.

81

Side Effects: Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475.

82

83 FIRST AID:

84

85 HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20
86 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue
87 rinsing eye.

88 Call a poison control centre or doctor for treatment advice.

89 IF SWALLOWED, call a poison control centre or doctor immediately for treatment
90 advice. Have person sip a glass of water if able to swallow. Do not induce vomiting
91 unless told to do so by a poison control centre or doctor. Do not give anything by mouth
92 to an unconscious person.

93

94 IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately
95 with plenty of water for 15–20 minutes. Call a poison control centre or doctor for
96 treatment advice.

97

98 PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large
99 amounts of water. If signs continue, consult a veterinarian immediately.

100 Take container, label or product name and Pest Control Product Registration Number
101 with you when seeking medical or veterinary attention.

102

103 TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

104

105 DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and
106 dispose in household garbage.

107

108 DIRECTIONS FOR USE:

109 This product contains imidacloprid and pyriproxyfen. Do not apply another pest control
110 product such as a shampoo, collar, or powder that contains imidacloprid or pyriproxyfen
111 to the treated animal after applying Advantage II Large Dog.

112 Do NOT administer by mouth. For dogs 7 weeks and older weighing between 11 and 25
113 kg. This product is packaged in single dosage tubes. This package contains one to six
114 monthly 2.5 mL dosages. Treat your dog monthly in the manner described below. Do
115 not apply a dosage higher than one 2.5 mL tube at one time. Consult a veterinarian
116 before using on sick, aged, pregnant or nursing animals or animals receiving drug or
117 other pesticide treatment. A veterinarian should be consulted when pest infestation
118 continues to be a problem on the pet. Retreatment at monthly intervals is required if
119 reinfestation is continuous or if flea control is desired.

120

121 Use only on dogs.

122 Do not use on other animals.

123 Do not use on puppies under 7 weeks of age.

124 Remove one applicator from the package.

2022-0231

2022-01-31

125 Hold applicator tube in upright position. Pull cap off tube. Turn cap around and place
126 other end of cap back on tube.

127 Twist cap to break seal, then remove cap from tube.

128 The dog should be standing for easy application. The entire contents of the Advantage®
129 II large dog tube should be applied evenly to four or five spots on the top of the back
130 from the shoulder to the base of the tail. At each spot part the hair until the skin is
131 visible. Place the tip of the tube on the skin and gently squeeze to dispense a portion of
132 the solution on the skin. Do not apply an excessive amount of solution at one spot that
133 could cause some of the solution to run off the side of the dog. Do not get this product
134 in your pet's eyes or administer by mouth. Until Advantage® II large dog dries, keep pet
135 off furniture and away from hardwood floors.

136

137 Discard the empty tube as described in "DISPOSAL".

138

139 RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE

140

141 Advantage, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

142 ©2022 Elanco or its affiliates.

143

2022-0231
2022-01-31

144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163

164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184

Advantage™ II Large Dog
Carton (English)

<Elanco Logo>

Advantage™ II large dog

Flea Adulticide/Larvicide/Ovicide and Lice Adulticide

For Dogs 7 weeks and older weighing between 11 and 25 kg
Once-A-Month Topical Flea and Lice Treatment

Liquid

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS
Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING



POISON

EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 27582 P.C.P. ACT

CONTENTS: One to Six 2.5 mL Tubes

Elanco Canada Limited
1919 Minnesota Court, Suite 401
Mississauga, Ontario L5N 0C9
1-800-265-5475

185
186 Advantage II large dog provides dual protection against fleas and lice. Advantage II
187 large dog kills 98-100% of the existing fleas on dogs within 1 day after treatment.
188 Protection against further flea infestation can last up to 4 weeks. Retreatment at
189 monthly intervals is required if reinfestation is continuous or if flea control is desired.
190 Fleas, eggs, and larvae are killed following contact with skin debris from an Advantage II
191 large dog treated pet. Advantage II large dog provides multi-stage flea control effectively
192 breaking all flea life-cycle stages for quick and lasting control of flea populations.
193 Advantage II large dog treats lice infestations on dogs.

194
195 Do not shampoo or immerse the animal in water immediately after treatment. The
196 animal can be shampooed, immersed in water or exposed to rain 3 days after treatment.
197 Advantage II large dog retains its ovicidal and larvicidal efficacy up to 4 weeks following
198 a shampoo treatment, swimming or exposure to rain. The adulticidal effect may be
199 shortened by repeated shampooing, water immersion or exposure to rain. Reapplication
200 may be required after such activities.

201
202 **PRECAUTIONS:**

203
204 **HUMANS: KEEP OUT OF THE REACH OF CHILDREN.** Harmful if swallowed, or
205 absorbed through the skin. Causes eye irritation. **DO NOT** get in eyes. Avoid contact
206 with skin or clothing. Wear rubber gloves during application of this product. After
207 handling or applying, wash hands (or any other skin that came in contact with the
208 product), with soap and water. Avoid contact with pets until dry.

209
210 **PETS:** Do not use on animals under 7 weeks of age. Care should be taken to prevent
211 exposure of other pets who may be in close physical contact with the treated animal.
212 Individual sensitivities, while rare, may occur after using ANY pesticide product for pets.
213 Smaller animals are more likely to experience an adverse reaction. Observe carefully. If
214 signs persist, or become more severe, consult a veterinarian immediately.

215

Side Effects: Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475.

216
217 **FIRST AID:**

218
219 **HUMANS: IF IN EYES,** hold eye open and rinse slowly and gently with water for 15–20
220 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue
221 rinsing eye.

222 Call a poison control centre or doctor for treatment advice.

223 **IF SWALLOWED,** call a poison control centre or doctor immediately for treatment
224 advice. Have person sip a glass of water if able to swallow. Do not induce vomiting

2022-0231

2022-01-31

225 unless told to do so by a poison control centre or doctor. Do not give anything by mouth
226 to an unconscious person. IF ON SKIN OR CLOTHING, take off contaminated clothing.
227 Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control
228 centre or doctor for treatment advice.

229

230 PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large
231 amounts of water. If signs continue, consult a veterinarian immediately.

232 Take container, label or product name and Pest Control Product Registration Number
233 with you when seeking medical or veterinary attention.

234

235 TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

236

237 DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household
238 garbage.

239

2022-0231
2022-01-31

240
241
242
243
244
245
246
247

248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266

<Elanco Logo>

Advantage™ II Large Dog
Tube (English)

Advantage™ II large dog

READ THE LABEL AND INSERT BEFORE USING

WARNING



POISON

EYE IRRITANT

DOMESTIC

EXTERNAL USE ONLY

ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 27582 P.C.P. ACT

Elanco Canada Limited
Mississauga, Ontario L5N 0C9

2.5 mL