

PRINCIPAL DISPLAY PANEL

GROUP	2	HERBICIDE
GROUP	4	HERBICIDE

TRITON[®] C Toss-N-Go Herbicide

Dry Flowable 75%

FOR SALE FOR USE ON WHEAT (SPRING OR DURUM), AND SPRING BARLEY (GROWN FOR FEED PURPOSES ONLY) IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

AGRICULTURAL

REGISTRATION NO. 28348 PEST CONTROL PRODUCTS ACT

**WARNING – EYE AND SKIN IRRITANT
CAUTION – POISON**

GUARANTEE:	Quinclorac	57.8 %
	Thifensulfuron methyl	11.5 %
	Tribenuron methyl	5.8 %
		<hr/>
		75.1 %

Warning: Contains the allergens milk and sulfites.

NET CONTENTS: 1.4 Kg.
(4 x 350 gram water soluble bags)

READ THE LABEL AND THE ATTACHED BOOKLET BEFORE USING

E.I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
BOX 2300 STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

PRECAUTIONS:

- WARNING - EYE AND SKIN IRRITANT.
- POTENTIAL SKIN SENSITIZER
- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE VENTILATION.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WEAR GOGGLES OR FACE SHIELD, COVERALLS, SOCKS, SUITABLE CHEMICAL RESISTANT GLOVES AND FOOTWEAR DURING MIXING AND LOADING. WEAR LONG SLEEVED SHIRT, LONG PANTS, SOCKS AND CHEMICAL RESISTANT GLOVES AND FOOTWEAR DURING APPLICATION, CLEAN-UP AND REPAIR. CHEMICAL RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. CAUSES EYE AND SKIN IRRITATION.
- WHEN USING, DO NOT EAT, DRINK OR SMOKE. WASH SKIN THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.
- REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE. STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE REUSE.
- DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.
- DO NOT APPLY BY AIR
- THE ENCLOSED POUCHES OF TRITON[®] C TOSS-N-GO HERBICIDE WILL DISSOLVE IN WATER. **DO NOT** ALLOW THE POUCHES TO BECOME WET BEFORE USE. **DO NOT** HANDLE INDIVIDUAL POUCHES WITH WET HANDS OR WET GLOVES. ALWAYS RESEAL THE ORIGINAL CONTAINER TO PROTECT THE REMAINING UNUSED POUCHES.
- DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS FOR 12 HOURS FOLLOWING APPLICATION.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca/.

FIRST AID:

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container or label or the product name and Pest Control Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

REGISTERED TRADEMARKS:

®/™ Registered trademarks/trademarks of E. I. du Pont Company. E. I. du Pont Canada Company is a licensee. Member of CropLife

GROUP	2	HERBICIDE
GROUP	4	HERBICIDE

TRITON[®] C Toss-N-Go[®] Herbicide

Dry Flowable 75%

FOR SALE FOR USE ON WHEAT (SPRING OR DURUM), AND SPRING BARLEY (GROWN FOR FEED PURPOSES ONLY) IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

AGRICULTURAL

REGISTRATION NO. 28348 PEST CONTROL PRODUCTS ACT

**WARNING – EYE AND SKIN IRRITANT
CAUTION – POISON**

GUARANTEE:	Quinclorac	57.8 %
	Thifensulfuron methyl	11.5 %
	Tribenuron methyl	5.8 %
		<hr/>
		75.1%

Warning: Contains the allergens milk and sulfites.

NET CONTENTS: 1.4 Kg.
(4 x 350 gram water soluble bags)

READ THE LABEL AND THIS ATTACHED BOOKLET BEFORE USING

E.I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
BOX 2300 STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

PRECAUTIONS:

- WARNING - EYE AND SKIN IRRITANT.
- POTENTIAL SKIN SENSITIZER
- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE VENTILATION.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WEAR GOGGLES OR FACE SHIELD, COVERALLS, SOCKS, SUITABLE CHEMICAL RESISTANT GLOVES AND FOOTWEAR DURING MIXING AND LOADING. WEAR LONG SLEEVED SHIRT, LONG PANTS, SOCKS AND CHEMICAL RESISTANT GLOVES AND FOOTWEAR DURING APPLICATION, CLEAN-UP AND REPAIR. CHEMICAL RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS..
- DO NOT CONTAMINATE ANY BODY OF WATER.
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. CAUSES EYE AND SKIN IRRITATION.
- WHEN USING, DO NOT EAT, DRINK OR SMOKE. WASH SKIN THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.
- REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE. STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE REUSE.
- DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.
- DO NOT APPLY BY AIR
- THE ENCLOSED POUCHES OF TRITON[®] C TOSS-N-GO HERBICIDE WILL DISSOLVE IN WATER. **DO NOT** ALLOW THE POUCHES TO BECOME WET BEFORE USE. **DO NOT** HANDLE INDIVIDUAL POUCHES WITH WET HANDS OR WET GLOVES. ALWAYS RESEAL THE ORIGINAL CONTAINER TO PROTECT THE REMAINING UNUSED POUCHES
- DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS FOR 12 HOURS FOLLOWING APPLICATION.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca/.

FIRST AID:

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container or label or the product name and Pest Control Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

ENVIRONMENTAL HAZARDS:

- Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided. Observe buffer zones specified under DIRECTIONS FOR USE.
- Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls, or calcium deposits.
- Do not remove soil from treated fields for use on lawns, in gardens or for back fill.
- Do not contaminate irrigation water. Do not apply to land where tail water will be used to irrigate cropland.
- Do not apply more than 87 g per hectare of Triton[®] C Toss-N-Go[®] Herbicide per season.
- Do not apply when weather conditions may cause spray drift from treated areas to adjacent crops. Certain crops, such as alfalfa, canola, clover species, fababeans, flax, lentils, ornamentals, potatoes and vegetables will be injured by spray drift.
- Quinclorac is persistent and will carryover. It is recommended that any products containing quinclorac not be used in areas treated with this product in the previous season.
- TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

TABLE OF CONTENTS:

PRINCIPAL DISPLAY PANEL..... 1
PRECAUTIONS:..... 2
 TOXICOLOGICAL INFORMATION:..... 3
STORAGE:..... 3
DISPOSAL:..... 3
NOTICE TO USER:..... 3
REGISTERED TRADEMARKS:..... 3
PRECAUTIONS:..... 5
 TOXICOLOGICAL INFORMATION:..... 6
TREAT SYMPTOMATICALLY. 6
 ENVIRONMENTAL HAZARDS:..... 6
PRODUCT DESCRIPTION: 7
 REPLANTING TO OTHER CROPS: 8
 GRAZING/HARVEST RESTRICTIONS:..... 9
 VARIETY RESTRICTIONS:..... 9
APPLICATION INFORMATION: 9
 APPLICATION PRECAUTIONS:..... 9
 APPLICATION INSTRUCTIONS: 10
 BUFFER ZONES: 10
DIRECTIONS FOR USE: 10
 SPRING WHEAT, DURUM, SPRING BARLEY (GROWN FOR FEED PURPOSES ONLY), NOT
 UNDERSEEDED TO FORAGES. 10
 TRITON® C TOSS-N-GO® 11
 TRITON® C TOSS-N-GO® + PUMA* 120 SUPER..... 12
 MIXING INFORMATION: 12
 MIXING PRECAUTIONS: 12
 MIXING DIRECTIONS:..... 13
 EQUIPMENT CLEANING INFORMATION: 13
 CLEANING PRECAUTIONS: 14
 CLEANING INSTRUCTIONS: 14
RESISTANCE-MANAGEMENT RECOMMENDATIONS: 14
STORAGE:..... 15
DISPOSAL:..... 15
NOTICE TO USER:..... 15
WATER SOLUBLE BAG LABEL 16

PRODUCT DESCRIPTION:

TRITON® C TOSS-N-GO® Herbicide is recommended for selective post-emergent control or suppression of broadleaf weeds in spring wheat, durum, and spring barley (grown for feed purposes only), not underseeded to legumes or grasses.

TRITON® C TOSS-N-GO® Herbicide contains 3 active ingredients formulated as dry flowable, water dispersible granules, blended in a water soluble Toss-N-Go® bags. It is non-corrosive, non-flammable, non-volatile and does not freeze. Should product be exposed to freezing temperatures, warm to room temperature prior to use.

Use gloves and goggles when handling the water-soluble bags. Reseal unused Toss-N-Go® Bags into original bag when not in use. Avoid exposure of Toss-N-Go® Bags to excess moisture or physical damage during storage and prior to mixing in spray tank.

TRITON® C TOSS-N-GO® Herbicide is to be mixed with water and Merge* adjuvant and applied as a uniform broadcast spray. Apply early post emergence, to the main flush of actively growing broadleaf weeds. APPLY WITH GROUND EQUIPMENT ONLY.

Warm, moist growing conditions promote active weed growth and enhance the activity of TRITON® C TOSS-N-GO® by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

TRITON® C TOSS-N-GO® stops growth of susceptible weeds rapidly. However, typical symptoms of dying weeds, including twisting, chlorosis, stunting and discolouration, may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

TRITON® C TOSS-N-GO® may provide residual activity in soil, and for this reason treated fields cannot be rotated to crops and recrop intervals other than specified in the section Replanting to Other Crops.

IMPORTANT INFORMATION: READ BEFORE USING

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, recreational grounds or similar areas.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water that may be used on other crops.
- Carefully observe sprayer clean-up instructions, as spray tank residues may damage crops other than wheat, spring barley or oats.

REPLANTING TO OTHER CROPS:

CROP	MINIMUM RECROPPING INTERVAL(months)	NOTES
Spring wheat, durum, spring barley	0	In the event of a crop failure, only these crops can be reseeded in treated fields.

Canola, field peas, sunflower	10	These crops can be planted in the year following treatment.
Flax, lentils*	22	These crops can be planted in treated fields in the second year after treatment.

On lighter soils with low organic matter or under dry conditions, some crop injury may occur particularly in flax and lentils but will not reduce yield. Under these conditions, the minimum recropping interval for flax and lentils should be extended by 12 months.

TRITON® C TOSS-N-GO® should not be used on land where potatoes or vegetables are part of the rotation.

Other crops should only be planted after a field bioassay has shown that it is safe to do so. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year.

GRAZING/HARVEST RESTRICTIONS:

Spring wheat, durum, or spring barley may not be grazed or fed to livestock within 77 days of application.

Do not apply TRITON® C TOSS-N-GO® within 77 days of harvest of spring wheat or durum or 80 days of harvest of spring barley.

VARIETY RESTRICTIONS:

Limit use of TRITON® C TOSS-N-GO® to currently licensed varieties of spring wheat, durum and spring barley. Consult E.I. du Pont Canada Company prior to use on unlicensed varieties for crop tolerance information. **DO NOT USE ON LEGER SPRING BARLEY OR BELVEDERE WHEAT VARIETIES.**

APPLICATION INFORMATION:

APPLICATION PRECAUTIONS:

- DO NOT APPLY BY AIR.
- Do not apply to irrigated land where tail water will be used to irrigate cropland.
- Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls or calcium deposits.
- Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided. Observe buffer zones specified under DIRECTIONS FOR USE.
- Apply with an adjuvant registered for use with TRITON® C TOSS-N-GO®. Use only water as a carrier. Other carriers may accelerate breakdown of TRITON® C TOSS-N-GO® and reduce its effectiveness.
- Do not apply to crops under seeded to legumes or grasses as injury to forage may result.
- Application to crops that are stressed by severe weather conditions, frost, low fertility, water logged soil (soil at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury.
- Under certain conditions, such as heavy rainfall, prolonged cool weather, frost or wide fluctuations in day/night temperatures, temporary lightening in crop colour, and occasionally, a slight reduction in crop height may occur.

- Do not apply more than 87 grams/hectare of TRITON® C TOSS-N-GO® per year.
- Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.

APPLICATION INSTRUCTIONS:

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh or larger filter screens (metal or nylon). A spray pressure of 210-275 kPa is recommended.

Continuous agitation is required to keep TRITON® C TOSS-N-GO® in suspension. Apply uniformly. Avoid overlapping and shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

Field Sprayer Application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, rangelands, riparian areas and shrub lands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Buffer Zones (metres) Required for the Protection of:		
	Aquatic Habitats of Depths:		Terrestrial Habitats
	Less than 1 m	Greater than 1 m	
Field sprayer*	1	0	15

* For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

DIRECTIONS FOR USE:

SPRING WHEAT, DURUM, SPRING BARLEY (Grown for feed purposes only), NOT UNDERSEEDED TO FORAGES.

TRITON® C TOSS-N-GO®

APPLY	TRITON® C TOSS-N-GO® Merge* Adjuvant	87 grams/Ha 1% v/v (10 L/1000L)																																														
TIMING	Make application when crop is in the 2-5 leaf stage. For best results apply to young, actively growing weeds that are less than 10 cm in height or diameter(cleavers 1-4 whorls). Thorough coverage of foliage is important for consistent weed control.																																															
<table style="width: 100%; border: none;"> <thead> <tr> <th style="text-align: left; width: 50%;">Weeds Controlled</th> <th style="text-align: left; width: 50%;">Weeds Suppressed*</th> </tr> </thead> <tbody> <tr> <td>Annual smartweed (green smartweed, lady's thumb)</td> <td>Canada thistle¹</td> </tr> <tr> <td>Annual sowthistle (1-4 leaf)</td> <td>Round-leaved mallow⁴</td> </tr> <tr> <td>Ball mustard</td> <td></td> </tr> <tr> <td>Chickweed²</td> <td>Scentless chamomile</td> </tr> <tr> <td>Cleavers(1-4 whorls)</td> <td>Sow thistle¹</td> </tr> <tr> <td>Corn spurry</td> <td>Stork's-bill (2-6 leaf)</td> </tr> <tr> <td>Cow cockle</td> <td>Toadflax⁵</td> </tr> <tr> <td>Flixweed</td> <td></td> </tr> <tr> <td>Groundsel, common</td> <td></td> </tr> <tr> <td>Hemp-nettle</td> <td></td> </tr> <tr> <td>Kochia</td> <td></td> </tr> <tr> <td>Lamb's-quarters</td> <td></td> </tr> <tr> <td>Narrow-leaved hawk's-beard</td> <td></td> </tr> <tr> <td>Redroot pigweed</td> <td></td> </tr> <tr> <td>Russian thistle</td> <td></td> </tr> <tr> <td>Shepherd's purse</td> <td></td> </tr> <tr> <td>Stinkweed</td> <td></td> </tr> <tr> <td>Tartary buckwheat</td> <td></td> </tr> <tr> <td>Volunteer rapeseed⁶</td> <td></td> </tr> <tr> <td>Volunteer sunflower</td> <td></td> </tr> <tr> <td>Wild buckwheat³</td> <td></td> </tr> <tr> <td>Wild mustard</td> <td></td> </tr> </tbody> </table>			Weeds Controlled	Weeds Suppressed*	Annual smartweed (green smartweed, lady's thumb)	Canada thistle ¹	Annual sowthistle (1-4 leaf)	Round-leaved mallow ⁴	Ball mustard		Chickweed ²	Scentless chamomile	Cleavers(1-4 whorls)	Sow thistle ¹	Corn spurry	Stork's-bill (2-6 leaf)	Cow cockle	Toadflax ⁵	Flixweed		Groundsel, common		Hemp-nettle		Kochia		Lamb's-quarters		Narrow-leaved hawk's-beard		Redroot pigweed		Russian thistle		Shepherd's purse		Stinkweed		Tartary buckwheat		Volunteer rapeseed ⁶		Volunteer sunflower		Wild buckwheat ³		Wild mustard	
Weeds Controlled	Weeds Suppressed*																																															
Annual smartweed (green smartweed, lady's thumb)	Canada thistle ¹																																															
Annual sowthistle (1-4 leaf)	Round-leaved mallow ⁴																																															
Ball mustard																																																
Chickweed ²	Scentless chamomile																																															
Cleavers(1-4 whorls)	Sow thistle ¹																																															
Corn spurry	Stork's-bill (2-6 leaf)																																															
Cow cockle	Toadflax ⁵																																															
Flixweed																																																
Groundsel, common																																																
Hemp-nettle																																																
Kochia																																																
Lamb's-quarters																																																
Narrow-leaved hawk's-beard																																																
Redroot pigweed																																																
Russian thistle																																																
Shepherd's purse																																																
Stinkweed																																																
Tartary buckwheat																																																
Volunteer rapeseed ⁶																																																
Volunteer sunflower																																																
Wild buckwheat ³																																																
Wild mustard																																																

*Weed suppression is a visual reduction in weed competition (reduced population and/or vigour) as compared to an untreated check. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

Specific Weed Recommendations:

1. Canada thistle, Sow thistle: Apply TRITON® C TOSS-N-GO® plus recommended surfactant postemergence when the majority of the thistles have emerged and are actively growing. For best top growth control, apply when thistles are no larger than 15 cm tall, before bud stage, and before crop canopy prevents thorough coverage of thistles. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.
2. Chickweed: Apply TRITON® C TOSS-N-GO® plus recommended surfactant when the chickweed is small (1-6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.
3. Wild Buckwheat: Apply TRITON® C TOSS-N-GO® plus surfactant to actively growing wild buckwheat in the cotyledon - 3 leaf stage. Under stress conditions such as dry weather control may be reduced. Large plants may regrow after treatment.

4. Round-leaved Mallow: Apply TRITON® C TOSS-N-GO® plus a surfactant to actively growing round leaved mallow in the 2-6 leaf stage (10-12 cm in height). TRITON® C TOSS-N-GO® will keep mallow stunted but may not reduce the overall populations.
5. Toadflax: Apply TRITON® C TOSS-N-GO® plus recommended surfactant when toadflax is no taller than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.
6. Volunteer rapeseed: TRITON® C TOSS-N-GO® alone will not control imazethapyr tolerant canola varieties, (e.g. canola varieties with the PURSUIT SMART™) trait.

NOTE:

Avoid making applications to weeds when rainfall is threatening. Rainfall within 6 hours will result in reduced weed control of TRITON® C TOSS-N-GO®.

TRITON® C TOSS-N-GO® + PUMA* 120 SUPER

APPLY	TRITON® C TOSS-N-GO® Puma 120 Super	87 grams/Ha 770 mL/Ha
TIMING	Apply to spring and durum wheat at the 2 to 5-leaf stage or to spring barley (grown for feed purposes) at the 2 to 4-leaf stage. Apply the tank mix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller), or to cleavers in the 1-3 whorl. Consult the Puma* 120 Super label for additional information on appropriate weed staging, application instructions, environmental and use precautions	
Weeds Controlled		
Weeds controlled or suppressed by TRITON® C TOSS-N-GO® alone, plus Green Foxtail Wild Oats Barnyard Grass		

MIXING INFORMATION:

MIXING PRECAUTIONS:

- TRITON® C TOSS-N-GO® is formulated as a soluble granule sealed within a water-soluble package (Toss-N-Go® bag). Drop intact Toss-N-Go® bag directly into spray tank. The Toss-N-Go® bag and pesticide will dissolve readily in water.
- Do not allow the Toss-N-Go® bag to become wet prior to use. Do not handle individual Toss-N-Go® bag with wet hands or wet gloves as this may cause breakage. Do not remove Toss-N-Go® bag from over-wrap container except for immediate use.
- Do not open Toss-N-Go® bag or attempt to remove the contents from them. The contents are not designed to be re-measured or subdivided.
- Do not use opened or punctured Toss-N-Go® bag.
- If broken Toss-N-Go® bag is found when the over-wrap container is opened, avoid contact with, and inhalation of the product. Handle with chemical-resistant gloves and dispose of broken Toss-N-Go® bag according to the DISPOSAL section.

Do not allow the spray mixture to remain in the tank for more than 24 hours before spraying or the effectiveness may be reduced. If the spray mixture has been allowed to stand in the tank, use vigorous agitation to thoroughly disperse before resuming spraying.

NOTE: Do not mix the soluble bags with liquid fertilizers, substances that contain boron or substances that release free chlorine. Mixing the soluble bags with any of these compounds will result in an insoluble substance in the tank.

MIXING DIRECTIONS:

Select a spray volume that will ensure thorough coverage and a uniform spray pattern. Use a minimum of 55 L of spray volume per hectare. If tank mixing with other herbicides, always consult the label of the tank mix partner(s) for minimum spray volume requirements and apply the tank mixture using a water volume recommended for all products.

1. Fill the spray tank 1/4 to 1/3 full with fresh water.
2. Add the recommended number of TRITON® C TOSS-N-GO® bags to the spray tank while the agitator is running. Continuous agitation is recommended through the entire mixing procedure to ensure products are completely dissolved and/or in suspension.
3. After TRITON® C TOSS-N-GO® bags have fully dissolved and TRITON® C TOSS-N-GO® is in suspension, other tank mix components may be added while continuing to fill the tank with water. The proper sequence for tank mix components is:
 - a. dry formulations
 - b. emulsifiable concentrates
 - c. adjuvants (a recommended adjuvant must be added to improve wetting and contact activity)
 - d. If a drift control agent is to be tank mixed, it is to be added last.

For repeat loads, reduce the tank heel to 10% or less of the previous load. Fill tank with fresh water as described in step 1, and continue as directed. Emulsifiable concentrates may make dispersion of TRITON® C TOSS-N-GO® more difficult. For that reason, tank heels of the previous tank mix should be kept to 10% or less of the spray tank volume.

EQUIPMENT CLEANING INFORMATION:

CLEANING PRECAUTIONS:

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour, which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.

CLEANING INSTRUCTIONS:

To avoid subsequent injury to crops other than spring wheat, durum, or spring barley, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of TRITON® C TOSS-N-GO® from mixing and spray equipment as follows:

1. Drain tank and flush tank, boom and hoses with clean water for a minimum of ten minutes.
2. Visually inspect tank to ensure removal of all visible residues of TRITON® C TOSS-N-GO®. If necessary, repeat Step 1.
3. Fill the tank with clean water, and then add 1 litre household AMMONIA (containing minimum of 3% ammonia) per 100 litres of water, or a commercial tank cleaner such as Finnish*
4. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation.
5. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
6. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
7. Repeat Steps 3-5.
8. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, TRITON® C TOSS-N-GO® contains Group 2 and Group 4 herbicides. Any weed population may contain or develop plants naturally resistant to TRITON® C TOSS-N-GO®. Bags and other Group 2 and Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of TRITON® C TOSS-N-GO® or other Group 2 or Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted.
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.

- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local du Pont representative or the E. I. du Pont Canada Company hotline at 1-800-667-3925 for further information.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

®/™ Registered trademarks/trademarks of E. I. du Pont Company. E. I. du Pont Canada Company is a licensee. Member of CropLife

* All other products mentioned are trademarks of their respective companies.

WATER SOLUBLE BAG LABEL

TRITON® C TOSS-N-GO® HERBICIDE TOSS-N-GO® BAG

KEEP OUT OF REACH OF CHILDREN

CAUTION: SEE DIRECTIONS FOR USE FOLDER
FOR COMPLETE PRECAUTIONARY STATEMENTS
CAUTION: EYE AND SKIN IRRITANT

REGISTRATION NO. 28348 PEST CONTROL PRODUCTS ACT

GUARANTEE:	Quinclorac	57.8 %
	Thifensulfuron methyl	11.5 %
	Tribenuron methyl	5.8 %
		<hr/>
		75.1%

Warning: Contains the allergens milk and sulfites.

DRY FLOWABLE

NET CONTENTS: 350 GRAMS

THIS BAG TREATS 4 HECTARES (10 ACRES)

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.