

2023-1413
2024-01-10

This label has been updated according to the re-evaluation decision of Corn Gluten Meal RVD2023-02. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 01-16-2025 in accordance with the phase out period set out in the re-evaluation decision of Corn Gluten Meal RVD2023-02. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

[FRONT PANEL - ENGLISH]

Natural Active Ingredient*

With Without
[images to show treatment with and without]

**SCOTTS® TURF BUILDER® WEED PREVENT
CORN GLUTEN MEAL**

HERBICIDE
DOMESTIC

SOLID

Pre-Emergent Dandelion and Large Crabgrass Weed Seed Germination
Inhibitor

APPLY EARLY SPRING AND LATE SUMMER BEFORE WEEDS APPEAR

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Corn gluten meal 99.975%
REG. NO.: 29060 P.C.P. ACT

CAUTION: EYE IRRITANT
POTENTIAL SENSITIZER

5-25 kg

**If used inappropriately, any pest control product, including those with natural ingredients, may have health, safety or environmental risks: use this product only in accordance with label instructions.*

Imported by:
Scotts Canada Ltd.
2000 Argentia Road
Plaza #2, Suite 300
Mississauga, Ontario
L5N 1V8

1-866-436-7645

[BOTTOM GUSSETS - ENGLISH]

**SCOTTS® TURF BUILDER® WEED PREVENT
CORN GLUTEN MEAL**

[SIDE GUSSETS - ENGLISH]

**SCOTTS® TURF BUILDER® WEED PREVENT
CORN GLUTEN MEAL**

[BACK PANEL - ENGLISH]

**SCOTTS® TURF BUILDER® WEED PREVENT
CORN GLUTEN MEAL**

DIRECTIONS FOR USE:

May inhibit the germination of dandelion and large crabgrass seed in lawns where established perennial ryegrass and Kentucky bluegrass is predominant.

- For best results, apply with a Scotts® spreader at the recommended setting listed.
- Apply in spring before dandelion and large crabgrass seeds have germinated.
- Apply when soil is moist and when rain is forecasted within 2 days of treatment. If rainfall does not occur within 2 days of treatment, irrigation is required. Excessive moisture at time of treatment may reduce the effectiveness of the product.
- For best results: Apply to established lawns twice a year; once in the early spring 2 weeks before weed seed germination, and once in the late summer or early fall after heat stress has passed.
- Do not apply under windy conditions.
- Do not apply the product on newly seeded grass as it may inhibit seeds from germinating, wait until after first mowing when the grass root systems are established.
- Do not apply this product if over-seeding or re-sodding in the spring or fall.
- Not recommended for use with a fertilizer containing nitrogen.
- Established dandelion and large crabgrass, at the time of application, will not be controlled. The inhibitory effect of the product to weed seeds generally dissipates in five weeks following application.

- May inhibit weed seed germination when used in conjunction with a sound lawn maintenance program.

SPREADER SETTINGS:

Spreader	Setting
Scotts® Broadcast/Rotary Spreaders Apply header strips. Then apply product in longest direction of lawn. Shut off spreader or stop each time you reach a header strip. Each pass should be within 1.5 metres (5 feet) of each other to ensure an overlap.	15 (Go over lawn twice)
Scotts® Drop Spreader Mow lawn first. Apply 2 header strips. Apply in longest direction of lawn, overlapping wheel marks.	17 (Go over lawn three times)

[Illustration of spreading patterns]

- If not using a Scotts® spreader, apply at the rate of 9.8 kg per 100 sq. metres (20 lb per 1,000 sq. feet).

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Keep pets off the treated area until a thorough watering or rainfall moves the applied product into the thatch layer. May cause sensitization. Should not be applied if the applicator or a member of the household has a sensitivity or allergy to corn. Avoid contact with skin, eyes or clothing. Avoid inhaling/breathing dusts. For good hygiene practice, wear a long-sleeved shirt, long pants, closed footwear, waterproof gloves, and a dust mask when handling and applying the product.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Wash skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **IF INHALED:** Move to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

STORAGE: Keep in cool, dry conditions. Store this product away from food or feed, seed, fertilizer and other pesticides.

DISPOSAL: DO NOT reuse empty container. Dispose the empty container with household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites.

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund cheque promptly.

HELP IS ONLY A CALL OR CLICK AWAY!

Consumer Helpline:

1-800-543-TURF (8873)

Or visit our website www.scotts.ca

® used under license by Scotts Canada Ltd.

© 2023 Scotts Canada Ltd. World rights reserved.

OPTIONAL STATEMENTS

- [FOR 9.1 kg] Covers 93 sq. metres (1,000 sq. feet)
- [FOR 18.2 kg] Covers 186 sq. metres (2,000 sq. feet)