

2012-1107
2013-06-10

NUTRITE PRE-EMERGENT WEED SEED GERMINATION INHIBITOR

WITH CORN GLUTEN

For the inhibition of large crabgrass, white clover and dandelion seed germination in residential lawns where established perennial ryegrass or established Kentucky bluegrass are the predominant grass species.

To be used only in the manufacture of a fertilizer-pesticide that is registered under the Fertilizers Act.

DOMESTIC

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: corn gluten meal 100%

REG. NO. 27727 PEST CONTROL PRODUCTS ACT

CAUTION EYE IRRITANT

POTENTIAL SENSITIZER

NET CONTENTS: 9-25 kg

FERTI TECHNOLOGIES INC.
560 Chemin Rhéaume
St-Michel QC
J0L 2J0

Telephone: (450) 462-2555

Nutrite Pre-emergent Weed Seed Germination Inhibitor With Corn Gluten may inhibit large crabgrass, white clover and dandelion seed germination when used in conjunction with a sound lawn maintenance program. Established large crabgrass, white clover and dandelion at the time of application will not be inhibited. This product is formulated for use in residential lawns where established perennial ryegrass or established Kentucky bluegrass are the predominant grass species. Nutrite Pre-emergent Weed Seed Germination Inhibitor With Corn Gluten is to be used only in the manufacture of a fertilizer-pesticide that is registered under the Fertilizers Act.

DIRECTIONS FOR USE:

Do not apply under windy conditions. Product application dates may vary for both the spring and late summer/early fall application from year to year according to weather conditions. The inhibitory effect of the product to weed seeds generally dissipates in five weeks following application.

Nutrite Pre-emergent Weed Seed Germination Inhibitor With Corn Gluten is to be applied to a mature lawn having a well developed root mass. Do not apply the product on newly seeded grass as it may inhibit seeds from germinating, wait until after first mowing when root systems are established. If over-seeding or re-sodding in the spring, do not apply the product in the spring. If over-seeding or re-sodding in the fall, do not apply the product in the fall.

For best results: Apply to established turf twice a year; once in the early spring and once in the late summer/early fall.

SPRING APPLICATION:

Large Crabgrass, White Clover and Dandelion Inhibition:

Apply to established perennial ryegrass or established Kentucky bluegrass in early spring two weeks before weed seed germination.

LATE SUMMER/EARLY FALL APPLICATION:

Large Crabgrass, White Clover and Dandelion Inhibition:

Apply to established perennial ryegrass or established Kentucky bluegrass in late summer or early fall after heat stress has passed.

APPLICATION RATE:

Apply at a rate of 9.8 kg/100 m² (20 lbs. /1000 ft²).

METHOD OF APPLICATION:

1. Measure the area to be applied by taking the length multiplied by the width of the total area in metres. One normal step is approximately one metre. Measure the size of any buildings or driveway in the same manner and subtract this amount from the total area. The remaining total will represent the grass area to be treated.
2. Apply **Nutrite Pre-emergent Weed Seed Germination Inhibitor With Corn Gluten** using a rotary spreader for large areas. For smaller areas, apply **Nutrite Pre-emergent Weed Seed Germination Inhibitor With Corn Gluten** using a small hand spreader.
3. Apply **Nutrite Pre-emergent Weed Seed Germination Inhibitor With Corn Gluten** in a criss-cross pattern. Use one half the rate in each direction. Apply when soil is moist

and when rain is forecasted within 2 days of treatment. If rainfall does not occur within 2 days of treatment, irrigation is required. Excessive moisture at time of treatment may reduce the effectiveness of the product.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May cause sensitization. **Nutrite Pre-emergent Weed Seed Germination Inhibitor With Corn Gluten** should not be applied if a member of the household has a sensitivity or allergy to corn. Avoid contact with skin, eyes or clothing. Avoid inhaling/breathing dusts. For good hygiene practice, wear a long-sleeved shirt, long pants, closed footwear and gloves when handling the product. In addition, it is recommended that a dust mask be worn when transferring the product to the spreader.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN/CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

STORAGE: Keep in cool, dry conditions, away from seed, fertilizer and other pesticides.

DISPOSAL: Do not re-use the empty container. Wrap and dispose of empty container with household garbage.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.