KILLS RATS & MICE

VICTOR Mouse & Rat Paraffin Block

With Bitrex

GUARANTEE: bromadiolone...... 0.005% Warning, contains the allergens peanuts and wheat REG.NO. 27701 PCP ACT

DOMESTIC KEEP OUT OF REACH OF CHILDREN READ THE LABEL BEFORE USING

WARNING POISON

NET CONTENTS: 450 g

Woodstream Canada Corporation, Unit 1, 25 Bramtree Court, Brampton, ON L6S 6G2 Tel: 1-800-800-1819 Use Victor Mouse & Rat Paraffin Block with Bitrex for the control of Norway rats, Roof rats and House mice indoors and against the outside walls of homes.

Areas of use:

General: May be used inside homes in areas where no feed or food is stored.

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Selection of Treatment Areas: Place bait indoors in locations where rats and mice travel and will most likely find and consume the bait. Placement along outside walls of buildings, in corners, by gnawed openings, in and around burrows, and in concealed locations where rats and mice feed is desirable. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly.

Application Directions:

Each Victor Mouse & Rat Paraffin Block consists of a 450 g slab which is to be broken into 8 pieces (i.e. each piece is 56 grams).

For Norway rats (*Rattus norvegicus*) and Roof rats (*Rattus rattus*): Norway rats usually inhabit the ground floor of structures and burrows in the ground. They will range approximately 15 to 100 m from their nesting area seeking food and water. Roof rats, however, are excellent climbers and usually infest the upper stories of structures. Hence,

special attention must be paid to this species ability to climb by placing baits in the roof rats activity area. For both species, eliminate as much of the naturally occurring food as possible and place 3 to 10 pieces of block (usually at intervals of 5-10 m) within the treatment area. Maintain an uninterrupted supply of fresh bait for at least 10 days, or until signs of rat activity cease.

<u>For House Mice (*Mus musculus*)</u>: House mice are excellent climbers and will infest all floors in a structure. Mice will only range approximately 10 m from their nesting area seeking food. Eliminate as much of the naturally occurring food as possible and place 1 piece of block at intervals of 2-4 m within the treatment area. If a bait placement is not disturbed after several days, move the placement approximately 1 metre. Two (2) pieces of block per placement may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

Burrow Application: Place 1-2 pieces of block per burrow.

<u>Note:</u> Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as necessary.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS, AND

LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. Treated baits must be maintained in tamperresistant bait boxes or in locations not accessible to children, pets, or livestock. This material may reduce the clotting ability of blood and cause bleeding. This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID: **If in eyes:** flush with water for 15 minutes, If irritation persists, seek medical attention. **If on skin**, wash affected area thoroughly with soap and water. If swallowed: give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or blunt object; repeat until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATON: Vitamin K_1 , in the form of intramuscular or subcutaneous injections, or by oral ingestion, are suggested remedial treatements for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap and dispose of empty container with household garbage.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.