

2017-7405 2018-01-12

UNIT

**WILSON<sup>®</sup>**  
**ANTOUT<sup>®</sup>**  
**ANT BAIT GEL**

Insecticide  
Paste

DOMESTIC

REG. NO. 30596 P.C.P. ACT

GUARANTEE: Boron 0.61% (present as disodium tetraborate decahydrate)

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

42.5g syringe

Sure-Gro IP Inc./ Lic. Premier Tech Home & Garden Inc.  
P.O. Box 21001, Brantford, ON N3R 7W9  
1-800-268-2806

CARTON

**WILSON<sup>®</sup> ANTOUT<sup>®</sup>  
ANT BAIT GEL**

Insecticide  
Paste

DOMESTIC

REG. NO. 30596 P.C.P ACT

GUARANTEE: Boron 0.61% (present as disodium tetraborate decahydrate)

**READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

42.5g syringe (1 – 6 syringes per box)  
1-24 Bait Stations

Sure-Gro IP Inc. / Lic. Premier Tech Home & Garden Inc.  
P.O. Box 21001, Brantford, ON N3R 7W9  
1-800-268-2806

Wilson<sup>®</sup> AntOut<sup>®</sup> Ant Bait Gel incorporates boron (in the form of Borax) in special food bait which is highly attractive to ants. Ants pick up the bait and take it to their nests to feed the colony. The boron works slowly and sufficient bait should be made available for continuous feeding. The ants will return to their nests after consuming the bait and will feed their young and the queen, destroying the colony. Repeat placement of the bait until satisfactory control has been achieved.

Do not use sprays, foggers or any chemicals or cleaners near the bait containers or placements. This may contaminate the bait and cause the insects to cease feeding. For control of ants, use indoors or outdoors near homes, apartments and condominiums.

**DIRECTIONS:** Wilson® AntOut® Ant Bait Gel is applied with the syringe. Remove a syringe from the package and inject gel into cracks and crevices around windows and doors, inside and out. Use along baseboards and void areas in walls, where the bait is protected from exposure and inaccessible to children and pets. Make small pea-sized placements into cracks and crevices where ants are present, up to 5 pea-sized placements per metre. Each syringe will cover about 8 metres.

Wilson® AntOut® Ant Bait Gel syringe may also be used to inject the gel into devices that provide entry to ants, but protect the bait from exposure to humans, domestic animals and the elements. The bait can be used to refill ant bait stations sold with the bait.

**Indoors:** Place bait stations 1.5-3.0 metres apart where ants are numerous. Replace bait stations as needed. Place stations under food cabinets, sinks, stoves, refrigerators, food storage areas, attics, garages, basements, storage areas and closets. Place on exterior windowsills and in doorways to help prevent entry of ants. Some bait stations may be designed to be affixed to a wall or placed on a flat surface. When using bait stations, wash hands thoroughly before handling to remove any odors that may be repellent to ants, i.e. tobacco, garlic, chemicals/cleaners.

**Outdoors:** Place bait stations 1.5-3.0 metres apart near where ants enter dwellings. For best results place stations in shaded areas or create shade for the bait station. [Avoid placing bait station where it will receive direct sunlight whenever possible.] (Optional text)

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. Avoid contamination of water, foodstuffs and utensils. Wash after use. Use only in areas inaccessible to children and pets.

**FIRST AID: If swallowed,** call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If in eyes,** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If inhaled,** move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**DISPOSAL:** Do not reuse empty container. Wrap empty container and discard in household garbage.

**For additional product information or in case of emergency, spills or fire, call toll-free 1-800-268-2806. [www.pthomeandgarden.com](http://www.pthomeandgarden.com)**

**OPTIONAL TEXT:**

RTU

Refill

No Mess – No Ants

Use Indoors or Out

Honeydew Formula

Ants return to their nests after consuming the bait and feed their young and the queen, destroying the colony