


(Container)

 **CORTEVA™**
agriscience
GF-2654 TSOY
Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

For control of annual and perennial weeds and use in Enlist™ soybeans

AGRICULTURAL

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

GUARANTEE: 2,4-D, present as choline salt 456 g a.e./L
Solution

REGISTRATION NO. 30959 PEST CONTROL PRODUCTS ACT

DANGER  **POISON**

EYE IRRITANT

NET CONTENTS: 1 L - Bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

CAUSES IRRITATION OF SKIN AND EYES

DO NOT GET IN EYES

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY

AVOID CONTACT WITH SKIN AND CLOTHING

DO NOT APPLY BY AIR

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

PROTECTIVE CLOTHING AND EQUIPMENT

All Application Equipment

- When mixing/loading, wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eye wear (face shield or safety glasses). Rinse gloves before removal.
- When applying and during clean-up and repair, wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks and shoes. Rinse gloves before removal.
- Gloves are not required during application when the applicator is in an enclosed tractor.
- When handling more than 582 L per day workers must also use a closed system.

Application using Groundboom Equipment

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities.

Application using Handheld Equipment

- Applicators must wear coveralls over a long-sleeved shirt, long pants, respirator and chemical-resistant gloves.

Refer to Directions for Use section for any additional requirements.

MIXING

Transfer System

For containers larger than 20 L, use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

OPERATOR USE PRECAUTIONS

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. This product may cause severe irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness or nausea. Treat symptomatically.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL HAZARDS

- Toxic to small mammals, birds, aquatic organisms and non-target terrestrial plants.
- Observe buffer zones specified under DIRECTIONS FOR USE.
- This product will harm other broadleaved plants in the vicinity of the treatment area. If applying this product using a handheld sprayer, do not directly spray or allow the spray to drift onto ornamentals or gardens.
- Do not spray exposed roots of trees and ornamentals.

LEACHING

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

RUNOFF

- To reduce runoff from treated areas into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a strip of untreated vegetation between the treated area and the edge of the water body.
- To prevent runoff from domestic uses avoid spraying on driveways, sidewalks or other hard surfaces. Do not irrigate within 24 hours after application.

STORAGE

This product must be stored away from fertilizers, seeds, insecticides, fungicides or other herbicides intended for use on 2,4-D sensitive crops.

USE

When the container is empty, triple rinse and add the rinse water to the spray tank. Care must be taken during application to avoid contaminating irrigation ditches or domestic water supplies. Avoid using sprayers which contained 2,4-D to apply other pesticides to sensitive crops.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

If this product is exposed to temperatures below 0°C, it should be warmed to at least 5°C and mixed thoroughly before using.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Booklet)



GF-2654 TSOY Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

For control of annual and perennial weeds and use in Enlist™ soybeans

AGRICULTURAL

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

GUARANTEE: 2,4-D, present as choline salt 456 g a.e./L
Solution

REGISTRATION NO. 30959 PEST CONTROL PRODUCTS ACT

DANGER  **POISON**

EYE IRRITANT

NET CONTENTS: 1 L - Bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9

1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

CAUSES IRRITATION OF SKIN AND EYES

DO NOT GET IN EYES

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY

AVOID CONTACT WITH SKIN AND CLOTHING

DO NOT APPLY BY AIR

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

PROTECTIVE CLOTHING AND EQUIPMENT

All Application Equipment

- When mixing/loading, wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eye wear (face shield or safety glasses). Rinse gloves before removal.
- When applying and during clean-up and repair, wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks and shoes. Rinse gloves before removal.
- Gloves are not required during application when the applicator is in an enclosed tractor.
- When handling more than 582 L per day workers must also use a closed system.

Application using Groundboom Equipment

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities.

Application using Handheld Equipment

- Applicators must wear coveralls over a long-sleeved shirt, long pants, respirator and chemical-resistant gloves.

Refer to Directions for Use section for any additional requirements.

MIXING

Transfer System

For containers larger than 20 L, use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

OPERATOR USE PRECAUTIONS

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. This product may cause severe irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness or nausea. Treat symptomatically.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL HAZARDS

- Toxic to small mammals, birds, aquatic organisms and non-target terrestrial plants.
- Observe buffer zones specified under DIRECTIONS FOR USE.
- This product will harm other broadleaved plants in the vicinity of the treatment area. If applying this product using a handheld sprayer, do not directly spray or allow the spray to drift onto ornamentals or gardens.
- Do not spray exposed roots of trees and ornamentals.

LEACHING

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

RUNOFF

- To reduce runoff from treated areas into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a strip of untreated vegetation between the treated area and the edge of the water body.
- To prevent runoff from domestic uses avoid spraying on driveways, sidewalks or other hard surfaces. Do not irrigate within 24 hours after application.

STORAGE

This product must be stored away from fertilizers, seeds, insecticides, fungicides or other herbicides intended for use on 2,4-D sensitive crops.

USE

When the container is empty, triple rinse and add the rinse water to the spray tank. Care must be taken during application to avoid contaminating irrigation ditches or domestic water supplies. Avoid using sprayers which contained 2,4-D to apply other pesticides to sensitive crops.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

If this product is exposed to temperatures below 0°C, it should be warmed to at least 5°C and mixed thoroughly before using.

DIRECTIONS FOR USE

USE PRECAUTIONS

- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Restricted Entry Interval (REI)

- For scouting, do not enter or allow worker re-entry until 24 hours after treatment.
- For all other activities, do not enter or allow worker re-entry until 12 hours after treatment.

Ground Application - Cropland

Apply 50 to 200 L of spray solution per hectare depending on the type of application equipment used. Use sufficient water for even distribution. Spray at low pressures (200 to 275 kPa) when the weeds are actively growing.

Droplet Size: Apply as a coarse to extremely coarse spray (ASABE S-572 Standard). Use drift reducing nozzle tips in accordance with manufacturer directions that produce a droplet classification of coarse to extremely coarse to significantly reduce the potential for drift.

Refer to the Product Use Guide for further information.

To Prepare a Spray

Add half the required amount of water to the spray tank, then the GF-2654 TSOY Herbicide with agitation, and finally the balance of the water with continued agitation.

Spot Treatment

For knapsack application of GF-2654 TSOY Herbicide for spot treatment of weeds such as thistles, mix 0.2 L of product in 10 L of water. Wet all foliage thoroughly. Maximum 2 applications per year.

Maximum amount of product applied/day/person with handheld equipment: 6.1 L.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

FOR USE IN ENLIST™ SOYBEAN ONLY

GF-2654 TSOY Herbicide is a systemic herbicide with very limited soil residual activity and is intended for control of emerged annual and perennial weeds. GF-2654 TSOY Herbicide is selective to Enlist soybeans. Enlist soybeans contain an AAD-12 expressing event. AAD-12 provides resistance to 2,4-D Choline Salt herbicides. Information on Enlist soybean varieties containing AAD-12 expressing events may be obtained from your seed supplier.

Rates of application

Use the rates given below for each weed list. Use the lower rate in each group for seedling weeds (2-4 leaves) and growing rapidly; and the higher rate for weeds that are in bud, during dry, cool periods or where heavy infestations exist.

Susceptible Weeds (0.73 to 1.1 L/ha):

bluebur	mustards (except dog and green tansy)
burdock (before 4 leaf stage)	pigweed, Russian
canola, volunteer*	pigweed, redroot
cocklebur	plantain, common
false flax	ragweed, common
fleabane, daisy	ragweed, giant
flixweed	Russian thistle
goatsbeard	shepherd's purse
hoary cress	sunflower, annual
horsetail, field	sweet clover
kochia	vetch
lamb's-quarters	wild radish

*including glyphosate tolerant and Clearfield canola varieties.

Less susceptible weeds (1.3 - 1.8 L/ha):

biennial wormwood*	knotweed (before 4 leaf stage)
bindweed*, field	lady's thumb
bindweed, hedge	leafy spurge*
blue lettuce*	mustard, dog
buckwheat, tartary	oak leaf goosefoot
buckwheat, wild*	pineappleweed
burdock*	purslane, common
Canada thistle*	smartweed green
dandelion	smartweed, Pennsylvania
chickweed, common	sow-thistle, annual
Chickweed, mouse-eared*	sow-thistle, perennial*
field peppergrass	tansy, common
hairy galinsoga	velvetleaf

*Top growth control only.

Mixing Directions

Sprayer Clean-Out Instructions

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
3. Second rinse:
 - Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator, or Clean-Out Spray Tank Cleaner, or 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water, or similar tank cleaning agent as per manufacturer's recommendations while filling the tank with clean water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.
 - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
 - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

Timing of Application

Apply when weeds are small and Enlist soybean growth stage is no later than R2 (full flowering stage). Make one to two applications with a minimum of 12 days between applications. Do not apply GF-2654 TSOY Herbicide to Enlist soybeans later than the R2 stage.

Pre-Harvest Interval

- Do not graze treated Enlist soybeans.
- Do not harvest for forage or hay.

Restrictions

- Do not apply more than two post emergent applications per use season.
- Do not apply more than 3.6 L/ha of GF-2654 TSOY Herbicide (1.64 kg acid equivalent) per use season.
- Read and follow the DAS Stewardship Program (<http://www.traitsstewardship.com>) that accompanies the use of soybean seed containing the DAS-68416-4 gene.

Sensitive Plants

Vegetables, flowers, grapes, fruit trees and other desirable plants are sensitive to 2,4-D, even in minute quantities. Care should be taken to avoid spraying these types of plants or allowing spray mist to drift onto these plants during both their growing and dormant periods. Coarse sprays are less likely to drift. At higher temperatures, vaporization may cause injury to susceptible plants growing nearby.

This product may cause damage to lawns or pastures if applied before the grass is well established. In addition, most legumes may be damaged or killed.

BUFFER ZONES TO PROTECT SENSITIVE HABITATS

Buffer zones are not required for applications with handheld equipment and for spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Seasonal water bodies require buffer zones if there is water in them during application. Water bodies which do not fill on an annual basis need not be buffered.

BUFFER ZONE TABLE

Method of Application	Crop	Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer*	Enlist soybeans	1	0	1	0	1

*For field sprayer application, buffer zones can be reduced with the use of drift-reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labeled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labeled buffer zone can be reduced by 30%.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, GF-2654 TSOY Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to GF-2654 TSOY Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of GF-2654 TSOY Herbicide or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852 or at www.corteva.ca.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

™Trademark of Corteva Agriscience and its affiliated companies.