

2023-6119  
2024-04-03

Container Label

GROUP	14	15	HERBICIDE
-------	----	----	-----------

**INTEGRITY®**

For weed control in corn (field and sweet) and soybeans

**HERBICIDE**

**EMULSIFIABLE CONCENTRATE**

**COMMERCIAL (AGRICULTURAL)**

**ACTIVE INGREDIENTS:** Saflufenacil..... 68 g/L  
Dimethenamid-P ..... 600 g/L

**REGISTRATION NO. 29371**

**PEST CONTROL PRODUCTS ACT**

**WARNING - EYE AND SKIN IRRITANT  
POTENTIAL SKIN SENSITIZER**

**READ THE LABEL AND BOOKLET BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY  
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT  
1-800-454-2673**

**NET CONTENTS:** 1 L - 1000 L, Bulk

BASF Canada Inc.  
5025 Creekbank Road  
Building A, 2nd Floor  
Mississauga, Ontario L4W 0B6  
1-877-371-2273

INTEGRITY is a registered trade-mark of BASF SE, used under license by  
BASF Canada Inc.

## PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Potential skin sensitizer.
3. **DO NOT** take internally. Harmful if swallowed.
4. **DO NOT** get in eyes or on skin. Causes eye and skin irritation.
5. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.
6. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. In addition, wear goggles or face shield during mixing/loading.

Applicators performing mixing, loading, and application must wear chemical-resistant coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, and additionally, goggles or a face shield during mixing and loading.

7. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.
8. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

## FIRST AID

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## TOXICOLOGICAL INFORMATION

Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

## **ENVIRONMENTAL PRECAUTIONS**

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under Restrictions and Limitations. This product contains an aromatic petroleum distillate which is toxic to aquatic organisms. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

## **STORAGE**

1. Store the product in original, tightly-closed container and do not allow water to be introduced into this container.
2. **DO NOT** ship or store the product near food, feed, seed or fertilizers.
3. Store the product in a cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.

## **DISPOSAL**

### **Recyclable Containers**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

### **Returnable-Refillable Containers**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

## **Unused or unwanted product**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

#### **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Booklet

GROUP	14	15	HERBICIDE
-------	----	----	-----------

**INTEGRITY®**

For weed control in corn (field and sweet) and soybeans

**HERBICIDE**

**EMULSIFIABLE CONCENTRATE**

**COMMERCIAL (AGRICULTURAL)**

**ACTIVE INGREDIENTS:** Saflufenacil..... 68 g/L  
Dimethenamid-P ..... 600 g/L

**REGISTRATION NO. 29371**

**PEST CONTROL PRODUCTS ACT**

**WARNING - EYE AND SKIN IRRITANT  
POTENTIAL SKIN SENSITIZER**

**READ THE LABEL AND BOOKLET BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY  
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT  
1-800-454-2673**

NET CONTENTS: 1 L - 1000 L, Bulk

BASF Canada Inc.  
5025 Creebank Road  
Building A, 2nd Floor  
Mississauga, Ontario L4W 0B6  
1-877-371-2273

INTEGRITY is a registered trade-mark of BASF SE, used under license by  
BASF Canada Inc.

## Table of Contents

GENERAL INFORMATION .....	3
SAFETY AND HANDLING .....	3
PRECAUTIONS .....	3
FIRST AID .....	4
TOXICOLOGICAL INFORMATION .....	4
ENVIRONMENTAL PRECAUTIONS .....	4
STORAGE .....	5
DISPOSAL .....	5
NOTICE TO USER .....	5
DIRECTIONS FOR USE .....	6
WEEDS, RATES AND TIMING .....	6
CORN TANK MIXES .....	9
LIQUID FERTILIZER – CORN ONLY .....	10
MIXING INSTRUCTIONS .....	13
SPRAYING INSTRUCTIONS .....	13
RECROPPING GUIDELINES .....	14
RESTRICTIONS AND LIMITATIONS .....	14
RESISTANCE-MANAGEMENT RECOMMENDATIONS .....	16

## GENERAL INFORMATION

**INTEGRITY** will provide control of most annual grass and broadleaf weeds in corn (field and sweet), soybeans, miscanthus and big blue stem. Reduced weed control may occur if adequate moisture is not received.

**INTEGRITY** can be applied as a pre-emergence, pre-plant surface or pre-plant incorporated treatment in corn (field and sweet) either alone or in tank-mix combination. **INTEGRITY** will provide most effective weed control in corn when applied and subsequently incorporated into soil by rainfall or mechanical tillage prior to weed seedling emergence from soil.

**INTEGRITY** may also be applied as a pre-plant surface or pre-emergence application in reduced or no-till soybeans in tank-mix combination with glyphosate. MERGE Adjuvant is required for optimum burndown activity.

**INTEGRITY** combines the two active ingredients: saflufenacil, a potent inhibitor of protoporphyrinogen-oxidase (PPO) belonging to herbicide mode of action Group 14, and dimethenamid-P, a chloroacetamide belonging to the herbicide mode of action Group 15. Saflufenacil is rapidly absorbed by roots and foliage with limited systemic activity. Under active growing conditions, susceptible emerged weeds usually develop chlorotic and necrotic injury symptoms within hours and die within a few days. Susceptible germinating weed seeds will usually die as they reach the soil surface or shortly after emergence. Dimethenamid-P is a root- and shoot-inhibitor that controls susceptible weed seedlings before or soon after they emerge from the soil.

## SAFETY AND HANDLING

### PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Potential skin sensitizer.
3. **DO NOT** take internally. Harmful if swallowed.
4. **DO NOT** get in eyes or on skin. Causes eye and skin irritation.
5. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.
6. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. In addition, wear goggles or face shield during mixing/loading.

Applicators performing mixing, loading, and application must wear chemical-resistant coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, and additionally, goggles or a face shield during mixing and loading.

7. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.
8. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

## **FIRST AID**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## **TOXICOLOGICAL INFORMATION**

Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

## **ENVIRONMENTAL PRECAUTIONS**

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under Restrictions and Limitations. This product contains an aromatic petroleum distillate which is toxic to aquatic organisms. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.



## STORAGE

1. Store the product in original, tightly-closed container and do not allow water to be introduced into this container.
2. **DO NOT** ship or store the product near food, feed, seed or fertilizers.
3. Store the product in a cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.

## DISPOSAL

### Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

### Returnable-Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**Unused, unwanted product** For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## DIRECTIONS FOR USE

### WEEDS, RATES AND TIMING

#### Corn (Field and Sweet)

**INTEGRITY** may be applied to corn as a pre-plant surface<sup>1</sup>, pre-plant incorporated or pre-emergent treatment at a rate of 1.1 L/ha for season long control of weeds as listed below:

- Barnyard grass
- Canada fleabane <sup>2</sup>
- Common ragweed<sup>2</sup>
- Crabgrass (smooth, large)
- Eastern black nightshade <sup>3</sup>
- Fall panicum
- Foxtail (green, yellow, giant)
- Giant ragweed <sup>2</sup>
- Hairy galinsoga
- Lamb's-quarters
- Palmer amaranth <sup>2</sup>
- Redroot pigweed
- Nutsedge, yellow <sup>3</sup>
- Old witchgrass
- Smallflower galinsoga
- Velvetleaf
- Waterhemp <sup>2</sup>
- Wild buckwheat
- Wild mustard

<sup>1</sup> Apply in tank mix with glyphosate

<sup>2</sup> Includes glyphosate-resistant biotypes

<sup>3</sup> PPI application only

**INTEGRITY** may be applied to corn as a pre-emergent treatment at a reduced rate of 0.73 L/ha for early season control of the following weeds, when an in-crop application of another registered herbicide is planned.

- Canada fleabane <sup>1</sup>
- Common ragweed<sup>1</sup>
- Crabgrass (smooth, large)
- Fall panicum
- Foxtail (green, yellow, giant)
- Giant ragweed <sup>1</sup>
- Hairy galinsoga
- Lamb's-quarters
- Palmer amaranth <sup>1</sup>
- Redroot pigweed
- Smallflower galinsoga

- Velvetleaf
- Waterhemp <sup>1</sup>
- Wild buckwheat
- Wild mustard

<sup>1</sup> Includes glyphosate-resistant biotypes

## Soybeans

**INTEGRITY** at a rate of 0.37 L/ha is applied in tank-mix combination with glyphosate and MERGE Adjuvant as a surface application prior to planting or pre-emergent application to reduced or no-till soybeans for burndown control of the following emerged broadleaf weeds and early-season suppression of the following grass weeds. This treatment is to be used when there is a planned in-crop application of another registered herbicide. Consult the glyphosate label for rates and weeds controlled by glyphosate in addition to those listed below, as well as for further recommendations, precautions and restrictions. **INTEGRITY** is compatible with all liquid glyphosate formulations registered for use as a pre-plant treatment prior to soybeans in which glyphosate is present as an isopropylamine salt, di-ammonium salt or potassium salt.

<b>Broadleaf Weeds</b>	<b>Maximum Weed Stage</b>
Broadleaf plantain	beginning of flowering
Canada fleabane*	20 leaf
Common ragweed*	8 leaf
Giant ragweed*	8 leaf
Lady's-thumb	6 leaf
Lamb's-quarters	8 leaf
Palmer amaranth*	6 leaf
Perennial sow thistle (top growth burndown control)	8 leaf
Prickly lettuce (top growth only)	9 leaf
Redroot pigweed	8 leaf
Waterhemp*	6 leaf
Wild mustard	8 leaf
Shepherd's-purse	full flower
Stinkweed	full flower
Wild buckwheat	6 leaf
Dandelion (suppression only)	6 leaf
<b>Grasses (Early season suppression)</b>	
Barnyard grass	
Crabgrass (large)	
Foxtail (green, yellow)	

\*Includes glyphosate-resistant biotypes

## USE RATES

### Corn

Timing	Pre-emergent
Rate	0.73 L/ha – 1.1 L/ha
Remarks	<p>The 0.73 L/ha use rate will only provide early season control and should be used in a program where an in-crop application of another registered herbicide is planned.</p> <p>Some sweet corn hybrids may be more sensitive to the combination of saflufenacil and dimethenamid-p and injury might occur.</p> <p>DO NOT apply <b>INTEGRITY</b> at the 1.1 L/ha use rate to sweet corn grown on coarse soils as injury may occur.</p> <p>DO NOT apply <b>INTEGRITY</b> at the 1.1 L/ha use rate to field corn grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, as injury may occur.</p>

Timing	Pre-plant Incorporated
Rate	1.1 L/ha
Remarks	<p>Do not incorporate <b>INTEGRITY</b> greater than 3 cm deep.</p> <p>Some sweet corn hybrids may be more sensitive to the combination of saflufenacil and dimethenamid-P and injury might occur.</p> <p>DO NOT apply to sweet corn grown on coarse soils as injury may occur.</p> <p>DO NOT apply <b>INTEGRITY</b> at the 1.1 L/ha use rate to field corn grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, as injury may occur.</p>

Timing	Pre-plant Surface
Rate	1.1 L/ha
Remarks	<p>Apply in tank mix with glyphosate.</p> <p>DO NOT apply <b>INTEGRITY</b> at the 1.1 L/ha use rate to field corn grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, as injury may occur.</p>

Apply **INTEGRITY** in a water volume of 100 to 200 L/ha. For corn, a nitrogen solution or complete liquid fertilizer carrier may be used as a replacement for water for use in pre-plant incorporated or pre-emergent applications. Refer to Liquid Fertilizer section for additional details. DO NOT apply **INTEGRITY** in liquid fertilizer to soybeans.

## Soybeans

Timing	Pre-plant Surface Application or Pre-emergent Application
Rate	0.37 L/ha <b>INTEGRITY</b> + glyphosate
Water Volume	100 to 200 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 1 L/ha.
Weeds Controlled	Refer to the glyphosate label for rates and weeds controlled in addition to weeds listed in the Soybeans Weeds, Rates and Timing section.
Remarks	<p>Do not use rates higher than 0.37 L/ha or crop injury may result.</p> <p>Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.</p> <p>For Pre-plant timing, apply as a pre-plant application from 21 days prior to planting up to the day of planting.</p> <p>For Pre-emergent timing, apply prior to when the soybeans cause the ground to crack and no more than 3 days after planting.</p> <p>For pre-emergent timing, DO NOT apply to coarse textured soils with less than 2% organic matter.</p> <p>DO NOT incorporate as injury may result.</p>

## CORN TANK MIXES

### **INTEGRITY + Atrazine**

**INTEGRITY** may be tank mixed with atrazine at 750 g ai/ha and applied as a pre-emergent or pre-plant incorporated treatment to corn.

Timing	Pre-emergent or Pre-plant Incorporated
Rate	1.1 L/ha <b>INTEGRITY</b> + 1.56 L/ha Aatrex® Liquid 480*
Water Volume	100 to 200 L/ha
Weeds Controlled	Refer to the atrazine label for weeds controlled in addition to those controlled by <b>INTEGRITY</b> applied alone.
Remarks	<p>Do not incorporate <b>INTEGRITY</b> greater than 3 cm deep. DO NOT apply this tank mix to sweet corn grown on coarse soils as injury may occur.</p> <p>DO NOT apply <b>INTEGRITY</b> at the 1.1 L/ha use rate to field corn grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, as injury may occur.</p> <p>Read and follow the label of the tank mix partner product for further use instructions, precautions and restrictions.</p>

\* For atrazine with a different concentration, adjust the rate accordingly.

## **INTEGRITY + Glyphosate**

For burndown and season long residual control of weeds labeled for **INTEGRITY** alone treatment in no-till and reduced-till corn, **INTEGRITY** may be tank mixed at a rate of 1.1 L/ha with glyphosate and applied as an early pre-plant surface application. Consult the glyphosate label for weeds controlled by glyphosate, as well as for further recommendations, precautions and restrictions. **INTEGRITY** is compatible with all liquid glyphosate formulations registered for use as a pre-plant treatment prior to corn in which glyphosate is present as an isopropylamine salt, di-ammonium salt or potassium salt.

Timing	Pre-plant Surface Application
Rate	1.1 L/ha <b>INTEGRITY</b> + 2.5 L/ha of glyphosate (360 g/L equivalent)
Water Volume	100 to 200 L/ha
Weeds Controlled	Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>INTEGRITY</b> applied alone.
Remarks	DO NOT apply <b>INTEGRITY</b> at the 1.1 L/ha use rate to field corn grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, as injury may occur.  Read and follow the label of the tank mix partner product for further use instructions, precautions and restrictions.

## **LIQUID FERTILIZER – CORN ONLY**

**INTEGRITY** may be applied in a nitrogen solution or complete liquid fertilizer carrier as a replacement for water for use in pre-plant incorporated or pre-emergent applications in corn. Follow the tank mix partner product label for instructions regarding application in liquid fertilizers. Compatibility of **INTEGRITY** with the liquid fertilizer should be determined by mixing a small quantity of herbicide with a proportional quantity of liquid fertilizer in a jar test prior to loading the spray tank. DO NOT apply **INTEGRITY** in liquid fertilizer to soybeans.

## BIG BLUE STEM

The **DIRECTIONS FOR USE** for the uses described in this section of the label were developed by persons other than BASF Canada under the User Requested Minor Use Label Expansion program. For these uses, BASF Canada has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

**INTEGRITY** may be applied to big blue stem for early season control of the following weeds:

- Common ragweed
- Crabgrass (smooth, large)
- Fall panicum
- Foxtail (green, yellow, giant)
- Lamb's-quarters
- Redroot pigweed
- Velvetleaf
- Wild buckwheat
- Wild mustard

### USE RATES

Timing	First year of production: Pre-emergent to the crop (application can be made on same day as planting, as long as shoots have not emerged).
Rate	0.73 L/ha
Water Volume	100 – 200 L/ha
Remarks	<p>The 0.73 L/ha use rate will only provide early season control.</p> <p>DO NOT apply <b>INTEGRITY</b> to big blue stem grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, or on soils with less than 2% OM as injury may occur.</p> <p>DO NOT apply to big blue stem within 365 days of harvest.</p> <p>DO NOT graze or feed treated big blue stem to livestock.</p>

Initial transient crop injury to new plantings may be observed after application but should be outgrown and not affect yield.

## MISCANTHUS

The **DIRECTIONS FOR USE** for the uses described in this section of the label were developed by persons other than BASF Canada under the User Requested Minor Use Label Expansion program. For these uses, BASF Canada has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

**INTEGRITY** may be applied to miscanthus for early season control of the following weeds:

- Common ragweed
- Crabgrass (smooth, large)
- Fall panicum
- Foxtail (green, yellow, giant)
- Lamb's-quarters
- Redroot pigweed
- Velvetleaf
- Wild buckwheat
- Wild mustard

### USE RATES

Timing	<p>First year of production: Pre-emergent to the crop (application can be made on same day as planting, as long as shoots have not emerged from rhizomes)</p> <p>Second year of production and subsequent years: Broadcast post-emergent application to established miscanthus crowns in the spring</p>
Rate	0.73 L/ha
Water Volume	100 – 200 L/ha
Remarks	<p>The 0.73 L/ha use rate will only provide early season control.</p> <p>DO NOT apply <b>INTEGRITY</b> to miscanthus grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, or on soils with less than 2% OM as injury may occur.</p> <p>DO NOT apply to miscanthus within 365 days of harvest.</p> <p>DO NOT graze or feed treated miscanthus to livestock.</p>



## MIXING INSTRUCTIONS

1. When applying **INTEGRITY**, always start with a clean sprayer. Thoroughly clean the sprayer by flushing the system with water containing detergent. Refer to previously applied product labels for specific cleaning instructions.
2. Fill clean spray tank half full with clean water or liquid fertilizer (corn only) and start agitation system. Prior to mixing **INTEGRITY** or **INTEGRITY** tank mixtures in liquid fertilizer, refer to appropriate label sections for recommended uses in liquid fertilizer, application instructions and compatibility determinations. For applications in liquid fertilizer, ensure that the mixture is applied within 24 hours of mixing. **DO NOT** apply **INTEGRITY** in liquid fertilizer to soybeans.
3. If using a wettable powder or water dispersible granule tank mix partner, add correct amount to tank and continue agitation until thoroughly dispersed. Do not add a wettable powder or water dispersible granule to liquid fertilizer.
4. If required, add the correct amount of liquid or solution tank mix partner. Continue to agitate.
5. Add the correct amount of **INTEGRITY** while continuing agitation.
6. For applications to soybeans, add the correct amount of MERGE Adjuvant to the spray tank.
7. Continue agitation while filling the spray tank with the remaining amount of water or liquid fertilizer.
8. Maintain continuous and constant agitation throughout application until spraying is complete.
9. After any break in spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit overnight.
10. If an oil film starts to build up in the tank, drain it and clean the tank with strong detergent solution.
11. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

Dispose of all rinsings in accordance with provincial regulations.

## SPRAYING INSTRUCTIONS

Select proper nozzle to avoid spraying fine mist. For best results, use sprayers equipped with flat fan or similar nozzles to ensure coverage. Apply in a spray volume of 100 to 200 L/ha and at a pressure of 275 kPa. For applications to dense weed infestations and thick canopies, use the higher spray volume. Better coverage of the product results in enhanced control of weeds.

Keep bypass line on or near bottom of tank to minimize foaming. Use 16-mesh suction screens, 50-mesh screens elsewhere on sprayer.

## RECRIPPING GUIDELINES

<b>INTEGRITY Rate</b>	<b>Follow Crop</b>
<b>INTEGRITY</b> at 1.1 L/ha	Anytime: corn (field and sweet) 100 days: cereals other than corn 22 months: peppers, onions and sugar beets 11 months: all other crops
<b>INTEGRITY</b> at 0.73 L/ha	Anytime: corn (field and sweet) 100 days: cereals other than corn 22 months: peppers and onions 11 months: all other crops
<b>INTEGRITY</b> at 0.37 L/ha	Anytime: corn (field and sweet) 100 days: cereals other than corn 11 months: all other crops

## RESTRICTIONS AND LIMITATIONS

1. Clean sprayer thoroughly after use to avoid damage to the next crop sprayed.
2. DO NOT apply to field corn within 100 days of harvest.
3. DO NOT apply to sweet corn or soybeans within 60 days of harvest.
4. DO NOT apply **INTEGRITY** at the 1.1 L/ha use rate to sweet corn grown on coarse soils as injury may occur. DO NOT apply **INTEGRITY** at the 1.1 L/ha use rate to field corn grown on coarse soils with a Cation Exchange Capacity (CEC) less than or equal to 15 MEQ/100 g, as injury may occur.
5. DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT.
6. DO NOT graze or feed the treated immature corn to livestock prior to 40 days after application of **INTEGRITY**.
7. **INTEGRITY** is only to be applied to soybeans in no-till or minimum tillage situations. DO NOT apply to soybeans where conventional tillage has been implemented.
8. For applications in soybeans, DO NOT incorporate **INTEGRITY** as injury may result.
9. For use on soybeans in Eastern Canada only.
10. Do not graze the treated soybean crops or feed soybean forage, hay or straw to livestock.
11. DO NOT enter or allow worker entry into treated areas for 12 hours after application.
12. DO NOT apply directly to water. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

13. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
14. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
15. DO NOT apply during periods of dead calm. Avoid application when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.
16. This product may be tank mixed with a fertilizer, a supplement or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact BASF Canada Inc. at 1-877-371-2273 for information before mixing any pesticide that is not specifically recommended on this label.

Not all varieties and cultivars have been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced efficacy, or crop injury may result from mixing **INTEGRITY** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Consult a BASF representative for more information concerning additives.

17. Tank Mix Compatibility Determination: To determine compatibility, mix all components of the finished spray in proportionate quantities in a small jar. Add components to the test jar in the following order: wettable powders and water dispersible granules, liquid flowables and suspensions, emulsifiable concentrate formulations, and solutions. If the mixture does not ball-up or form flakes, sludge, jelly, oily films or layers, or other precipitates within 2 hours after mixing and can easily be resuspended with agitation, then the tested spray-mix is compatible. Always follow the most restrictive label regarding any precautions when tank mixing.

## **BUFFER ZONES**

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater

habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Buffer zones (metres) required for the protection of:	
		Aquatic habitats	Terrestrial habitats
Field sprayer*	Corn	1	10
	Soybeans	1	4
	Miscanthus	1	10
	Big Blue Stem	1	10

\*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labeled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labeled buffer zone can be reduced by 30%.

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

## RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that **INTEGRITY** is both a Group 14 and a Group 15 herbicide. Any weed population may contain plants naturally resistant to Group 14 and/or Group 15 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **INTEGRITY** or other Group 14 and Group 15 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative

herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273 or at [www.agsolutions.ca](http://www.agsolutions.ca).

®All other products listed are registered trademarks of their respective companies.