

**ON GUARD PROFESSIONAL P-1D-RTU  
READY-TO-USE  
RESIDUAL HOUSEHOLD INSECT SPRAY**

Kills ants, cockroaches, spiders, earwigs, millipedes  
and certain other specified crawling insects

DOMESTIC

REG. NO.31130 P.C.P. ACT

NET CONTENTS -----L

GUARANTEE: Propoxur 1%

DANGER

POISON

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

UR-CAN INC.

3325 North Service Rd. Unit 104

Burlington, ON

L7N 3G2

289-426-1363

Residual Household Insect Spray provides rapid kill of such crawling insects as listed below.

Ants	Flies	Sawtoothed grain beetles
Brown dog ticks	Gnats (exposed stage)	
Clover mites	Hornets	Silverfish
Crickets	Millipedes	Sowbugs
Earwigs	Mosquitoes	Spiders
Fleas	Cockroaches	Wasps

In addition to rapid kill, this Residual Household Spray stays on treated surfaces and kills insects many weeks after the spray deposit has dried.

**DIRECTIONS FOR USE** Insect Spray is ready to use undiluted as it comes from the can, and may be applied with either power-operated or hand sprayers. Apply as a coarse spray to infested surfaces until wet, moving the spray stream rapidly enough to prevent excessive wetting of surfaces and waste of material. Direct spray away from people, plants and pets.

**Indoor Use: ANTS, COCKROACHES, CRICKETS, SILVERFISH, SPIDERS, SAWTOOTHED GRAIN BEETLES (EXPOSED STAGE):** Spray into cracks, crevices, around baseboards and other places where these insects hide. Apply behind and beneath cabinets, refrigerators, sinks, stoves and in and around waste containers. For ants, spray trails and places where ants enter premises. Repeat as needed. Do not spray foodstuffs or houseplants. Remove pets and cover fishbowls before spraying. Do not allow children to contact treated areas until surfaces are thoroughly dry.

**BROWN DOG TICKS:** Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

**Outdoor use: ANTS, CLOVER MITES, CRICKETS, EARWIGS, FLEAS, FLIES, GNATS, HORNETS, MILLIPEDES, MOSQUITOES, SOWBUGS, BROWN DOG TICKS AND WASPS:** Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios, or other places where insects may enter. For ants, spray hills and runways. For control of wasps and hornets, spray nests at night. Repeat as necessary. Do not spray vegetation.

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.** May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing of spray mist. Contact with skin, eyes, or clothing should also be avoided. Wash thoroughly with soap and warm water after handling. Do not spray near open flame or spark. Do not smoke while using. Do not store near excessive heat. Do not contaminate feed or food. Prior to spraying, test a small area of rugs, carpets, draperies, wallpaper or similar materials that may be stained. Avoid excessive wetting of plastics, rubber and asphalt materials such as tiles and floor coverings. Do not use as a space spray. Remove pets and cover fishbowls before spraying. Do not treat animals, humans, plants, or food utensils. Provide adequate ventilation of area being treated.

**FIRST AID: If swallowed,** call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **If in eyes,** hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Propoxur inhibits cholinesterase resulting in stimulation of the central nervous system. This results in a sense of "tightness" in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhoea. The antidote is atropine sulphate administered in large therapeutic doses repeated as necessary to the point of tolerance. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically. This product contains petroleum distillate. Vomiting may cause aspiration pneumonia.

**DISPOSAL:** Do not reuse empty container. Wrap and dispose of empty container in household garbage.