

2021-2764  
2021-10-12

This label has been updated according to the re-evaluation decision of Imidacloprid RVD2021-05. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 19/05/2023 in accordance with the phase out period set out in the re-evaluation/special review decision of Imidacloprid RVD2021-05. This previously approved label will be provided upon request by emailing [hc.pmra.info-arla.sc@canada.ca](mailto:hc.pmra.info-arla.sc@canada.ca). In your email please include the product name and Registration number of the label you are requesting.



GROUP 4A INSECTICIDE

\*container\*

**ACCELERON® IX-409**  
**Insecticide Seed Treatment**

ACCELERON IX-409 INSECTICIDE SEED TREATMENT is a systemic Insecticide Seed Treatment for protection of soybean against listed insect pests.

Insecticide

COMMERCIAL

Suspension



WARNING

POISON

ACTIVE INGREDIENT: Imidacloprid ..... 600 g/L

Contains 1,2-benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 31068 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 0.25 - 1000 L

Bayer CropScience Inc.  
Suite 200, 160 Quarry Park Blvd. S.E.  
Calgary, Alberta T2C 3G3

In case of spills, poisoning, fire or other emergencies,  
call: 1-800-334-7577 (24 hours a day)

For product information, call: 1-800-283-6847

**SPECIAL USE RESTRICTIONS:** This product contains no colourant. Seed treated with ACCELERON IX-409 Insecticide Seed Treatment must be conspicuously coloured. Regulations pertaining to the *Seeds Act* must be strictly adhered to when using this product.

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## PRECAUTIONS

### KEEP OUT OF REACH OF CHILDREN.

Harmful or fatal if swallowed and harmful if inhaled. Avoid breathing vapour or spray mist. When handling ACCELERON IX-409 Insecticide Seed Treatment or seed treated with ACCELERON IX-409 Insecticide Seed Treatment, work in a well-ventilated area. When treating seeds, handling and planting treated seeds, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear. In commercial facilities, worker involved with cleaning seed treatment equipment must wear chemical-resistant coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides. Use good personal hygiene, washing hands and exposed skin before eating, drinking or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

### FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed	Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

### TOXICOLOGICAL INFORMATION:

Treat symptomatically.

### ENVIRONMENTAL PRECAUTIONS

Toxic to birds, small wild mammals and aquatic organisms.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites).

Imidacloprid is toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatment applications. When used according to label directions minimal exposure or risk is expected. Dust generated during planting of treated seed may be harmful to bees and other pollinators. To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at [www.canada.ca/pollinators](http://www.canada.ca/pollinators).

When using a seed flow lubricant with this treated seed, only a dust-reducing seed flow lubricant is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for soybean seed treated with this insecticide. Carefully follow use directions for this seed flow lubricant. Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds. When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.

Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

#### STORAGE

Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep container closed.

Do not store ACCELERON IX-409 Insecticide Seed Treatment in direct sunlight.

Do not store ACCELERON IX-409 Insecticide Seed Treatment above 35°C.

#### DISPOSAL

For non-returnable containers:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Triple rinse instructions for container disposal (for recyclable and disposable containers only):

This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment provided the total volume of water does not exceed 6 % of the container size. Add 1/3 of the rinse water to the container and swish the contents thoroughly. Empty the rinsate into the seed treater holding tank and repeat this process two more times making sure the total volume of rinsate does not exceed 6%. Make sure rinsate is thoroughly mixed with seed treatment before treating. Close the container.

**CAUTION:** Do not dilute beyond 6% or excess seed wetness may result in seed handling difficulties. Using excess water can also cause ACCELERON IX-409 Insecticide Seed Treatment to thin resulting in settling of solids in the product. Be sure to adjust the undiluted application rate up by 6% to compensate for the dilution of the product caused by the addition of rinsate.

**DISPOSAL OF UNUSED, UNWANTED PRODUCT:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.



\*booklet\*

GROUP 4A INSECTICIDE

**ACCELERON® IX-409**  
**Insecticide Seed Treatment**

ACCELERON IX-409 INSECTICIDE SEED TREATMENT is a systemic seed treatment insecticide for protection of soybean against listed insect pests.

Insecticide

COMMERCIAL

Suspension



WARNING

POISON

ACTIVE INGREDIENT: Imidacloprid ..... 600 g/L

Contains 1,2-benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 31068 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 0.25 - 1000 L

Bayer CropScience Inc.  
Suite 200, 160 Quarry Park Blvd. S.E.  
Calgary, Alberta T2C 3G3

In case of spills, poisoning, fire or other emergencies,  
call: 1-800-334-7577 (24 hours a day)

For product information, call: 1-800-283-6847

**SPECIAL USE RESTRICTIONS:** This product contains no colourant. Seed treated with ACCELERON IX-409 Insecticide Seed Treatment must be conspicuously coloured. Regulations pertaining to the *Seeds Act* must be strictly adhered to when using this product.

Table of Contents:		Section Number
General Information	Notice to User	1
	The Product	2
Safety and Handling	Precautions, Protective Clothing and Equipment	3
	First Aid and Toxicological Information	4
	Environmental Precautions	5
	Storage	6
	Disposal	7
Directions For Use	Crops and Pests	8
	Soybean	8.1
	Tank Mixtures	8.2
	Use Restrictions and Limitations	9
	Use Restrictions	9.1
	Seed Quality	9.2
	Labeling Treated Seed	9.3
	Special Use Restrictions	9.4
	Mixing Instructions	10
	Resistance Management Recommendations	11

## GENERAL INFORMATION

### Section 1: Notice to User

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

### Section 2: The Product

ACCELERON IX-409 Insecticide Seed Treatment is a systemic seed treatment insecticide that provides protection of certain crops from damage caused by listed chewing and sucking insects through contact and systemic activity. Thorough seed coverage is required for maximum protection of seed. Under high insect pressures, a foliar insecticide may be required, therefore monitor crops regularly for insect infestation levels. Do not apply any subsequent application of a Group 4 Insecticide (i.e., in-furrow or foliar application) following treatment with ACCELERON IX-409 Insecticide Seed Treatment.

## SAFETY AND HANDLING

### Section 3: Precautions, Protective Clothing and Equipment

#### PRECAUTIONS

##### KEEP OUT OF REACH OF CHILDREN.

Harmful or fatal if swallowed and harmful if inhaled. Avoid breathing vapour or spray mist. When handling ACCELERON IX-409 Insecticide Seed Treatment or seed treated with ACCELERON IX-409 Insecticide Seed Treatment, work in a well-ventilated area. When treating seeds, handling and planting treated seeds, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear. In commercial facilities, worker involved with cleaning seed treatment equipment must wear chemical-resistant coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides. Use good personal hygiene, washing hands and exposed skin before eating, drinking or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or [www.cropscience.bayer.ca](http://www.cropscience.bayer.ca).

### Section 4: First Aid and Toxicological Information

#### FIRST AID:



Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed	Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

#### TOXICOLOGICAL INFORMATION:

Treat symptomatically.

Section 5: Environmental Precautions
--------------------------------------

Toxic to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

Toxic to aquatic organisms.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites).

Imidacloprid is toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatment applications. When used according to label directions minimal exposure or risk is expected. Dust generated during planting of treated seed may be harmful to bees and other pollinators. To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at [www.canada.ca/pollinators](http://www.canada.ca/pollinators) .

When using a seed flow lubricant with this treated seed, only a dust-reducing seed flow lubricant is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for soybean seed treated with this insecticide. Carefully follow use directions for this seed flow lubricant. Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds. When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies. Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Section 6: Storage
--------------------

Store product in original container only, away from other pesticides, fertilizer, food or feed.

Keep container closed.

Do not store ACCELERON IX-409 Insecticide Seed Treatment in direct sunlight.

Do not store ACCELERON IX-409 Insecticide Seed Treatment above 35°C.

Section 7: Disposal
---------------------

#### Disposal of Container

For non-returnable containers:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Triple rinse instructions for container disposal (for recyclable and disposable containers only): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment provided the total volume of water does not exceed 6 % of the container size. Add 1/3 of the rinse water to the container and swish the contents thoroughly. Empty the rinsate into the seed treater holding tank and repeat this process two more times making sure the total volume of rinsate does not exceed 6%. Make sure rinsate is thoroughly mixed with seed treatment before treating. Close the container.

**CAUTION:** Do not dilute beyond 6% or excess seed wetness may result in seed handling difficulties. Using excess water can also cause ACCELERON IX-409 Insecticide Seed Treatment to thin resulting in settling of solids in the product. Be sure to adjust the undiluted application rate up by 6% to compensate for the dilution of the product caused by the addition of rinsate.

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements. Do not dispose of seed by double-sowing in headlands. Broadcast seeding is prohibited.

Section 8: Crops and Pests
----------------------------

### 8.1 SOYBEAN:

ACCELERON IX-409 Insecticide Seed Treatment is a systemic insecticide seed treatment that provides early season protection against soybean aphid, crop stand injury caused by wireworm, larvae of European chafer and Japanese beetle and seedcorn maggot, and early season defoliation caused by the over-wintering generation of bean leaf beetle.

Do not apply any subsequent application of a Group 4 Insecticide (i.e., in-furrow or foliar application) following treatment with ACCELERON IX-409 Insecticide Seed Treatment.

Crop	Pest	Use rate/100 kg seed
------	------	----------------------

		mL product	grams a.i.
Soybeans	Soybean aphid, Bean leaf beetle, Seedcorn maggot, Wireworm, Larvae of European chafer and Japanese beetle	104	62.5

Ensure product is thoroughly mixed prior to application or use entire container at one time. Apply ACCELERON IX-409 Insecticide Seed Treatment through slurry applicator seed treaters which provide uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect control. Maintain constant agitation of the slurry during application. Allow the seed to dry before bagging or storing in bulk containers. This product contains no colourant. All seed must be conspicuously coloured at the time of treatment in accordance with *Seed Act and Regulations*.

## 8.2 TANK MIXTURES:

For control of certain seed and soil-borne pathogens in soybean seeds and seedlings, ACCELERON IX-409 Insecticide Seed Treatment may be mixed with the following seed treatment fungicides: Trilex AL, Trilex AL Concentrate, Trilex FS, Allegiance FL, ACCELERON D-309, EverGol Energy, EverGol Xtend, Apron Maxx RFC and Apron Maxx RTA. Follow all appropriate directions and precautions as specified on the fungicide labels. Make sure that the specific legume crop to be treated is registered on the fungicide partner as well. When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

## SECTION 9: USE RESTRICTIONS AND LIMITATIONS

### 9.1 USE RESTRICTIONS:

Do not use treated seed for food, feed or oil processing.  
Do not graze or feed livestock on treated areas for four weeks after planting.

### 9.2 SEED QUALITY:

Laboratory and field studies have shown that ACCELERON IX-409 Insecticide Seed Treatment will not adversely affect germination of treated seeds. Treatment of highly mechanically damaged seed or seed of known low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. If seed lot quality is unknown, conduct a germination test on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

### 9.3 LABELLING TREATED SEED:

All treated soybean seed for sale or use in Canada must be labelled with the following information:

- This seed has been treated with ACCELERON IX-409 Insecticide Seed Treatment seed protectant which contains imidacloprid.
- Do not use for feed, food or oil processing. Store away from feeds and other foodstuffs.
- Toxic to birds, small wild mammals and aquatic organisms.
- Imidacloprid is toxic to bees. Dust generated during planting of treated seed may be harmful to bees and other pollinators.
- To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at [www.canada.ca/pollinators](http://www.canada.ca/pollinators) .
- When using a seed flow lubricant with this treated seed, only a dust-reducing fluency agent is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for soybean seed treated with this insecticide. Carefully follow use directions for the seed flow lubricant.
- Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds.
- When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.
- Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.
- Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements. Do not dispose of seed by double-sowing in headlands.
- Broadcast seeding is prohibited.
- Closed cabs must be used for planting commercially treated and bagged seeds.
- Do not plant treated seeds by hand.

#### 9.4 SPECIAL USE RESTRICTIONS:

This product contains no colourant. Seed treated with this product must be conspicuously coloured. Regulations pertaining to the *Seeds Act* must be strictly adhered to when using this product.

<b>SECTION 10: MIXING INSTRUCTIONS</b>
--

Storage of ACCELERON IX-409 Insecticide Seed Treatment at low temperatures is not recommended. Prior to and during application, ACCELERON IX-409 Insecticide Seed Treatment must be thoroughly agitated to ensure uniform mixing of the product. Due to the viscosity of the material, it should be kept above 10°C prior to and during application. Do not apply direct heat to container.

## SECTION 11: RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that ACCELERON IX-409 Insecticide Seed Treatment contains a Group 4A insecticide. Any insect population may contain individuals naturally resistant to ACCELERON IX-409 Insecticide Seed Treatment and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of ACCELERON IX-409 Insecticide Seed Treatment or other Group 4 insecticides with different groups that control the same pests in a field.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Bayer CropScience Inc. via internet at [www.cropscience.bayer.ca](http://www.cropscience.bayer.ca) or phone 1-888-283-6847.