

2023-1185
2023-12-15

This label has been updated according to the re-evaluation decision of Corn Gluten Meal RVD2023-02. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 01-16-2025 in accordance with the phase out period set out in the re-evaluation decision of Corn Gluten Meal RVD2023-02. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.



Ingredion

Prairie Gold™ Corn Gluten Meal D

Granular

(pre-emergent weed seed germinator inhibitor with corn gluten)

For the inhibition of dandelion and smooth crabgrass seed germination in lawn where established Kentucky bluegrass is predominant.

DOMESTIC

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Corn Gluten Meal 99.7%

Contains calcium propionate at 0.3% as a preservative

WARNING: Contains the allergen Sulfites

REG. NO. 29639 P.C.P. ACT

CAUTION EYE IRRITANT
POTENTIAL SENSITIZER

NET CONTENTS: 20 Kg

Ingredion Canada Incorporated
1600-90 Burnhamthorpe Road West
Mississauga, Ontario
L5B 0H9

Application Rate:

SPRING APPLICATION:

Apply to established Kentucky bluegrass turf in early spring application before weed seed germination at a rate of 9.7 kg/100 m² (20lbs. /1000ft²).

SUMMER APPLICATION:

Apply to established Kentucky bluegrass turf in summer before weed seed germination at a rate of 9.7 kg/100 m² (20lbs. /1000ft²).

FALL APPLICATION

Apply to established Kentucky bluegrass turf in autumn before weed seed germination, at a rate of 9.7 kg/100 m² (20lbs. /1000ft²).

Directions:

1. Measure the area to be applied by taking the length multiplied by the width of the total area in metres. One normal step is approximately one metre. Measure the size of any buildings, swimming pool or driveway in the same manner and subtract this amount from the total area. The remaining total will represent the grass area to be treated.
2. Apply Prairie Gold™ Corn Gluten Meal D (pre-emergent weed seed germinator inhibitor) in a criss-cross pattern. Use one half the rate in each direction. Apply when soil is moist and when rain is forecasted within 2 days of treatment. If rainfall does not occur within 2 days of treatment, irrigation is required. Excessive moisture at time of treatment may reduce the effectiveness of the product.
3. Apply Prairie Gold™ Corn Gluten Meal D (pre-emergent weed seed germinator inhibitor) using a rotary spreader for large areas. For smaller areas, apply Prairie Gold™ (pre-emergent weed seed germinator inhibitor) using a small hand spreader.

Prairie Gold™ must not be applied to newly seeded grass until after the first mowing when the root systems are established, since the product inhibits seeds from germinating. For best results: Apply to established Kentucky bluegrass turf three times a year; once in the early spring, once in summer and once in fall.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May cause sensitization. Prairie Gold™ Corn Gluten Meal D (pre-emergent weed seed germinator inhibitor) (pre-emergent weed seed germination inhibitor) should not be applied if a member of the household has a sensitivity or allergy to corn.

Avoid contact with skin, eyes or clothing.

Avoid inhaling dusts.

For good hygiene practice, wear a long-sleeved shirt, long pants, closed footwear, waterproof gloves, and a dust mask when handling and applying the product.

Keep pets off the treated area until a thorough watering or rainfall moves the applied product into the thatch layer.

NOTE: Do not apply the product on newly seeded grass as it may inhibit seeds from germinating, wait until after first mowing when root systems are established. Do not apply if allergic to corn. Do not apply the product in any season in which overseeding is being carried out as the grass seed may not germinate.

FIRST AID:

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN/CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

2023-1185
2023-12-15

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Take the container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically

STORAGE:

Store this product away from food or feed

DISPOSAL:

Do not reuse empty container. Dispose the empty container with household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites.