

2020-1059
2020-06-12

GROUP	9	HERBICIDE
-------	---	-----------

CHEMINOVA GLYPHOSATE II™
Soluble Concentrate Herbicide
Solution AGRICULTURAL AND INDUSTRIAL

ACTIVE INGREDIENT: glyphosate 356 g/L (acid equivalent) present as the isopropylamine salt

WARNING: EYE IRRITANT

REGISTRATION NO. 28925 PEST CONTROL PRODUCTS ACT
READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

IN CASE OF EMERGENCY, CALL 1-800-331-3148 (24 HOURS)

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

Label

NET CONTENTS: 10 L

READ ENTIRE LABEL CAREFULLY BEFORE USE

CHEMINOVA GLYPHOSATE II is a non-selective, non-residual herbicide containing 356 g/L glyphosate as isopropylamine salt, formulated as a water soluble liquid. It is used for the control of most herbaceous weeds in agricultural and industrial sites. The product is absorbed through the foliage and translocated throughout the plant down to the root system. Visible symptoms such as gradual wilting and yellowing are usually obvious within 2 to 4 days of application to annual weeds, but may not be apparent for 7 to 10 days on perennial weeds.

GENERAL PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN
- CAUSES EYE IRRITATION
- HARMFUL IF SWALLOWED
- AVOID CONTACT WITH EYES AND SKIN
- WASH HANDS AND EXPOSED SKIN BEFORE EATING, DRINKING, OR SMOKING, AND AFTER WORK

FOR GOOD AGRICULTURAL PRACTICE:

- DURING MIXING, LOADING, CLEANUP, AND REPAIR PROCEDURES WEAR GLOVES, COVERALLS, AND EYE PROTECTION
- WASH SPLASHES FROM SKIN AND EYES IMMEDIATELY

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

EMERGENCY TELEPHONE NUMBER

IN CASE OF A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL TOLL FREE, DAY OR NIGHT, 1-800-331-3148.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL HAZARDS

Avoid direct application to any body of water populated with fish or used for domestic purposes. Do not use in areas where adverse impact on domestic water or aquatic species is likely. Do not contaminate water by disposal of waste or cleaning of equipment. Avoid all drift or contact with vegetation for which treatment is not intended as damage or destruction may occur.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic, or plastic-lined containers. **DO NOT MIX, STORE, OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or the spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.

STORAGE

KEEP AWAY FROM FOOD, DRINK, AND ANIMAL FEEDSTUFFS. KEEP ONLY IN ORIGINAL CONTAINER, TIGHTLY CLOSED.

DISPOSAL AND DECONTAMINATION

RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

RETURNABLE-REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

TMCHEMINOVA GLYPHOSATE II and FMC are trademarks of FMC Corporation or an affiliate.

GROUP

9

HERBICIDE

CHEMINOVA GLYPHOSATE™ II
Soluble Concentrate Herbicide

Solution AGRICULTURAL AND INDUSTRIAL

WARNING: EYE IRRITANT

NET CONTENTS : 10 L

ACTIVE INGREDIENT: glyphosate 356 g/L (acid equivalent) present as the isopropylamine salt

READ THE LABEL AND THIS BOOKLET BEFORE USING
REGISTRATION NO. 28925 PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY, CALL 1-800-331-3148 (24 HOURS)

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

Booklet

CHEMINOVA GLYPHOSATE II - LABEL INDEX

GENERAL PRECAUTIONS; FIRST AID; EMERGENCY TELEPHONE NUMBER	
ENVIRONMENTAL HAZARDS; PHYSICAL OR CHEMICAL HAZARDS · STORAGE · DISPOSAL	
PRECAUTIONS FOR USE	
GENERAL PRODUCT INFORMATION	
GENERAL APPLICATION NOTES	
TANK MIXES	
VEGETATION CONTROLLED (LISTS)	
APPLICATION EQUIPMENT & MIXING INSTRUCTIONS	
GROUND BOOM & BOOMLESS SPRAYERS	
KNAPSACK SPRAYERS, HAND-HELD & HIGH VOLUME EQUIPMENT	
MIST BLOWERS	
WIPER, WICK & ROLLER EQUIPMENT	
AERIAL APPLICATION	
AGRICULTURAL & CROPLAND USES	
PREPLANT TREATMENT	
SUMMER FALLOW	
MINIMUM & ZERO TILLAGE SYSTEMS	
MINIMUM & ZERO TILLAGE TANK MIXES	
FALL STUBBLE	
SPOT TREATMENT (IN CROP)	
FORAGE GRASSES & LEGUMES	
PASTURE RENOVATION	
FORAGE SEED PRODUCTION(FOR SPOT TREATMENT)	
PREHARVEST CONTROL / HARVEST MANAGEMENT	
TREE, VINE, & BERRY CROPS	
TREE PLANTING - SHELTERBELTS, NURSERY STOCK, WOODY ORNAMENTALS	
NONCROPLAND & INDUSTRIAL USES	
TURFGRASS	
TREE INJECTION APPLICATIONS	
CUT STUMP APPLICATIONS	

WOODY BRUSH & TREES (FOLIAR APPLICATIONS)	
INDUSTRIAL SITES, RIGHTS-OF-WAY, RECREATIONAL & PUBLIC AREAS	
PURPLE LOOSESTRIFE CONTROL	

RATES AND USES CHARTS:

TANK MIXES FOR ANNUAL WEED CONTROL	
TANK MIXES FOR PERENNIAL WEED CONTROL	
GUIDELINES FOR TIMING OF PREHARVEST APPLICATIONS	
WEED CONTROL IN NONCROPLAND, INDUSTRIAL USES	
WEED CONTROL IN TREE, VINE, AND BERRY CROPS	
ANNUAL WEED CONTROL	
PERENNIAL WEED CONTROL	

READ ENTIRE LABEL CAREFULLY BEFORE USE

CHEMINOVA GLYPHOSATE II is a non-selective, non-residual herbicide containing 356 g/L glyphosate as isopropylamine salt, formulated as a water soluble liquid. It is used for the control of most herbaceous weeds in agricultural and industrial sites. The product is absorbed through the foliage and translocated throughout the plant down to the root system. Visible symptoms such as gradual wilting and yellowing are usually obvious within 2 to 4 days of application to annual weeds, but may not be apparent for 7 to 10 days on perennial weeds.

GENERAL PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN
- CAUSES EYE IRRITATION
- HARMFUL IF SWALLOWED
- AVOID CONTACT WITH EYES AND SKIN
- WASH HANDS AND EXPOSED SKIN BEFORE EATING, DRINKING, OR SMOKING, AND AFTER WORK

FOR GOOD AGRICULTURAL PRACTICE:

- DURING MIXING, LOADING, CLEANUP, AND REPAIR PROCEDURES WEAR GLOVES, COVERALLS, AND EYE PROTECTION
- WASH SPLASHES FROM SKIN AND EYES IMMEDIATELY

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

EMERGENCY TELEPHONE NUMBER

IN CASE OF A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL TOLL FREE, DAY OR NIGHT, 1-800-331-3148.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL HAZARDS

Avoid direct application to any body of water populated with fish or used for domestic purposes. Do not use in areas where adverse impact on domestic water or aquatic species is likely. Do not contaminate water by disposal of waste or cleaning of equipment. Avoid all drift or contact with vegetation for which treatment is not intended as damage or destruction may occur.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic, or plastic-lined containers. **DO NOT MIX, STORE, OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or the spray solutions of this product react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.

STORAGE

KEEP AWAY FROM FOOD, DRINK, AND ANIMAL FEEDSTUFFS. KEEP ONLY IN ORIGINAL CONTAINER, TIGHTLY CLOSED.

DISPOSAL AND DECONTAMINATION

RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

RETURNABLE-REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS FOR USE

Avoid contact with desirable vegetation by direct application or spray drift as severe injury or destruction may result. Avoid drift or overspray to non-target vegetation and wildlife habitats.

DO NOT USE IN GREENHOUSES.

Drain and clean sprayer and parts immediately after using this product.

Do not contaminate water sources by disposal of wastes or cleaning of equipment.

Reduced results may occur if water containing soil is used such as water from ponds and ditches. Poor control may also occur when treating weeds heavily covered with dust.

GENERAL PRODUCT INFORMATION

CHEMINOVA GLYPHOSATE II is a water soluble herbicide for non-selective weed control. CHEMINOVA GLYPHOSATE II is applied as a foliar spray for the control of

most herbaceous plants. It may be applied through most standard industrial or field type sprayers after dilution and thorough mixing with water in accordance with the booklet instructions.

CHEMINOVA GLYPHOSATE II moves through the plant from the point of foliage contact into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur until 7 to 10 days. Extremely cool or cloudy weather at treatment time may slow down the activity of this product and delay visible effects of control. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts.

CHEMINOVA GLYPHOSATE II does not provide residual weed control. For subsequent residual weed control, apply a registered residual herbicide. Read and carefully observe cautionary statements and all other information appearing on the labels of all herbicides used.

Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

GENERAL APPLICATION NOTES

Results are best when weeds are actively growing. If weeds have been mowed, allow to return to recommended growth stage.

Delay application until vegetation has emerged to the stage described for the control of such vegetation under the **ANNUAL** and **PERENNIAL WEED CONTROL** charts of this booklet to provide adequate leaf surface to receive the spray. Unemerged plants arising from underground rhizomes or root stocks of perennials will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds is obtained when the treatment is made at the late growth stages approaching maturity. Always use higher rates of CHEMINOVA GLYPHOSATE II per hectare within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought, flooding, frost, high temperatures, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

CHEMINOVA GLYPHOSATE II should only be mixed with products recommended on this label. Do not mix with any surfactant, pesticide, herbicide oils or any other material other than water unless specified.

TANK MIXES

CHEMINOVA GLYPHOSATE II may be used with the following surfactants:

Agra[®] 90, AgSurt[®], Companion[™], Frigate[®]. See charts on **TANK MIXES** for **ANNUAL** and for **PERENNIAL WEED CONTROL**.

CHEMINOVA GLYPHOSATE II may be used with the following herbicides:

Banvel[®], Pardner[®], Pursuit[®], 2,4-D low volatile ester or amine formulations: See section on **MINIMUM AND ZERO TILLAGE**

TANK MIXES.

Princep[®] *Nine-T*[®], *Simadex*[®]: See section on **TREE, VINE, AND BERRY CROPS.**

DyCleer[®] 480, *Simazine 80W*, *Simadex*[®] *Flowable*, *2,4-D amine*: See section on **NONCROPLAND AND INDUSTRIAL USES.**

Always refer to the surfactant and herbicide labels for specific instructions regarding the use of that product.

Trade Name

Agral, Frigate
AgSurf
Banvel,
Pursuit
Companion
DyCleer, Princep,
Nine-T Pardner
Simadex

Trademark

of Zeneca
Agro IPCO
BASF
Rohm & Haas Co.
Novartis
Rhone-Poulenc
AgrEvo

VEGETATION CONTROLLED

CHEMINOVA GLYPHOSATE II controls many annual and perennial grasses, broadleaf weeds and woody brush and trees when applied as recommended and under the conditions described. For information on how to control specific weeds, including herbicide rate, refer to the **ANNUAL WEED CONTROL** and **PERENNIAL WEED CONTROL** charts of this label. The following is a partial list of the weeds controlled:

Annual weeds:

Annual bluegrass
Barnyard grass
Chickweed
Cleavers
Cocklebur
Corn spurry
Cow cockle
Crab grass
Dodder
Downy brome
Eastern black flowering nightshade
Fleabane (Canada)
Flixweed
Giant foxtail
Green foxtail
Green smartweed
Hempnettle
Kochia
Lady's- thumb
Lamb's- quarters
Narrow-leaved hawk's- beard
Narrow-leaved vetch
Nightflowering catchfly
Pennsylvania smartweed
Persian darnel
Prickly lettuce
Ragweed
Redroot pigweed
Russian thistle
Shepherd's purse
Smooth pigweed
Sowthistle (annual)
Stinkweed
Velvetleaf
Volunteer barley
Volunteer canola
Volunteer corn
Volunteer flax
Volunteer wheat
Wild buckwheat
Wild mustard
Wild oats
Wild tomato

Poa annua
Echinochloa crus-galli
Stellaria media
Galium aparine
Xanthium strumarium
Spergula arvensis
Saponaria vaccaria
Digitaria sanguinalis
Cuscuta spp.
Bromus tectorum
Solanum ptycanthum
Erigeron canadense
Descurainia sophia
Setaria faberii
Setaria viridis
Polygonum scabrum
Galeopsis tetrahit
Kochia scoparia
Polygonum persicaria
Chenopodium album
Crepis tectorum
Vicia angustifolia
Silene noctiflora
Polygonum pennsylvanicum
Lolium persicum
Lactuca serriola
Ambrosia artemisiifolia
Amaranthus retroflexus
Salsola kali
Capsella bursa-pastoris
Amaranthus hybridus
Sonchus oleraceus
Thlaspi arvense
Abutilon theophrasti
Hordeum spp.
Brassica spp.
Zea mays
Linum spp.
Triticum spp.
Polygonum convolvulus
Brassica kaber
Avena fatua
Solanum triflorum

Perennial weeds:

Alfalfa
Bluegrass (Canada)
Bluegrass (Kentucky)

Medicago savita
Poa compressa
Poa pratensis

Bromegrass (smooth)
Canada thistle
Common cattail
Common milkweed
Cottontop
Curled dock
Dandelion
Field bindweed
Foxtail barley
Hemp dogbane
Hoary cress
Japanese knotweed
Perennial sowthistle
Poison ivy
Purple loosestrife
Quackgrass
Toad flax
Wormwood (Absinth)
Yellow nutsedge

Bromus inermis
Cirsium arvense
Typha latifolia
Asclepias syriaca
Eriophorum chamissionis
Rumex crispus
Taraxacum officinale
Convolvulus arvensis
Hordeum jubatum
Apocynum cannabinum
Cardaria draba
Polygonum cuspidatum
Sonchus arvensis
Rhus radicans
Lythrum salicaria
Elytrigia repens
Linaria vulgaris
Artemisia absinthium
Cyperus esculentus

Woody weeds, brush, and trees:

Alder
Birch
Broad-leaf meadowsweet
Canadian rhododendron
Cedar
Cherry
Douglas fir
Hemlock
Maple
Milkweed
Mountain-fly honeysuckle
Pine
Poison Ivy
Poplar
Raspberry
Salmonberry
Sheep laurel
Snowberry (western)
Sweet fern
Willow
Withrod

Alnus spp.
Betula spp.
Spiraea lattifolia
Rhododendron canadense
Thuja spp.
Prunus spp.
Pseudotsuga spp.
Tsuga spp.
Acer spp.
Asclepias syriaca
Lonicera villosa
Pinus spp.
Rhus radicans
Populus spp.
Rubus spp.
Rubus spp.
Kalmia angustifolia
Symphoricarpos occidentalis
Comptonia peregrina
Salix spp.
Viburnum cassinoides

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, CHEMINOVA GLYPHOSATE II is a Group 9 herbicide. Any weed population may contain or develop plants naturally resistant to CHEMINOVA GLYPHOSATE II and other Group 9 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance- management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of CHEMINOVA GLYPHOSATE II or other Group 9 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is

- permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
 - Monitor treated weed populations for resistance development.
 - Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
 - Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
 - For further information or to report suspected resistance, contact FMC of Canada Limited at 1-833-362-7722 or at www.fmccrop.ca.

APPLICATION EQUIPMENT AND MIXING

INSTRUCTIONS GROUND BOOM AND BOOMLESS SPRAYERS

Mixing: For field or industrial type sprayers, fill the spray tank with one half the required amount of water. Add the proper amount of CHEMINOVA GLYPHOSATE II herbicide (see appropriate chart) and mix well before adding the remaining portion of water. Placing the filling hose below the surface of the liquid solution will prevent any excessive foaming. Remove the hose from the tank immediately after filling to avoid back siphoning into water source (a one-way valve should be installed to prevent back siphoning). Use of mechanical agitators may cause excessive foaming. By-pass lines should terminate at the bottom of the tank.

Application: Use flat fan nozzles in boom sprayers. To control perennial weeds, woody brush, and trees as listed, apply CHEMINOVA GLYPHOSATE II in 50 to 300 L of water per hectare as a broadcast spray. Use no more than 275 kPa pressure.

To control annual weeds as listed, apply CHEMINOVA GLYPHOSATE II in 50 L to 100 L of water per hectare as a broadcast spray. Use no more than 275 kPa pressure.

KNAPSACK SPRAYERS. HAND-HELD & HIGH VOLUME EQUIPMENT

High volume spraying utilizes handguns or other suitable nozzle arrangements to apply a **directed spray** to weeds, woody brush, and trees. Use coarse sprays only.

Mixing: Mix the proper amount of CHEMINOVA GLYPHOSATE II with water in a large container. Fill the sprayer with the mixed solution. Unless otherwise stated, make a 1% solution of CHEMINOVA GLYPHOSATE II in water (1 L of CHEMINOVA GLYPHOSATE II in 100 L of water). A 2% solution (2 L of CHEMINOVA GLYPHOSATE II in 100 L of water) should be used on harder to control perennials.

Application: Spray coverage should be uniform and complete. Apply on a spray-to-wet basis. Do not spray to the point of runoff. Hand gun application should be properly directed to avoid spraying desirable plants.

MIST BLOWERS

For control of woody weeds, brush, and trees listed in the **VEGETATION CONTROLLED** list, use the recommended rate of CHEMINOVA GLYPHOSATE II in at least 200 L of water per hectare.

WIPER, WICK AND ROLLER EQUIPMENT

These applicators apply CHEMINOVA GLYPHOSATE II solution directly onto the weeds by contacting the weed with an absorbent material containing the herbicide solution.

Weeds should be a minimum of 15 cm above the desired vegetation to prevent contact of CHEMINOVA GLYPHOSATE II with the desired vegetation.

Mixing: Mix the proper amount of CHEMINOVA GLYPHOSATE II with water in a large container. Use this mixed solution in the wiper, wick or roller equipment.

Application: These applicators can be used to control weeds in:

- Industrial sites, tree plantings, and non-crop sites as specified.
- The following agricultural crops:
 - apple, cherry, peach, pear and plum orchards, grape vineyards, soybeans, dry beans, strawberries, and cranberries (note: applications must be made before initial pod set in soybeans and dry beans).

The applicator should be adjusted so that the contact point of the wiper, roller, or wick is at least 5 cm above the desirable vegetation. Droplets or foam of the CHEMINOVA GLYPHOSATE II solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Best results are obtained when more of the weed is exposed to the herbicide solution. It is recommended that two applications be made in opposite directions, if possible.

Weeds not contacted will not be affected. This may occur in dense clumps, severe infestation, or when the height of the weeds varies so that not all weeds are contacted. In these instances, a repeat treatment may be necessary.

AVOID CONTACT WITH DESIRABLE VEGETATION

Wiper, Wick, Roller Application Notes:

- Maintain wiper equipment in good operating condition. Care must be taken with all types of wipers to ensure that the absorbent material does not become over-saturated, causing the herbicide to drip onto desirable vegetation.
- Avoid leakage or dripping onto desirable vegetation.
- Adjust height of wiper applicator to ensure proper contact with weeds.
- Keep wiping surfaces clean.
- Maintain recommended roller speed on roller applicators while in use.
- DO NOT use wiper equipment when weeds are wet.
- DO NOT operate equipment at ground speeds less than 4 or greater than 10 km/h. Weed control may be affected by speed of application equipment. As weed density increases, reduce equipment ground speed to ensure good coverage of weeds.
- Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying on the upper end of the wiper applicator.
- Variation in equipment design may affect weed control. With wiper applicators, the wiping material and its orientation must allow delivery of sufficient quantities of the recommended CHEMINOVA GLYPHOSATE II herbicide solution directly to the weed.
- Mix only the amount of solution to be used during a one day period, as reduced activity may result from use of leftover solution. Thoroughly drain and clean all equipment immediately after use.

AERIAL APPLICATION

Directions for Use (for additional information see section on Aerial Application For Industrial Rights-of-Way ONLY)

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of

aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices. The use of a spotter plane is recommended.

Aerial Use Precautions

Apply only when weather conditions at the treatment site allow for complete and even crop coverage.

Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls, and goggles or face shield during mixing/loading, cleanup, and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit, and vehicle cabs must be decontaminated regularly.

Product-Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at (973) 305-6600, or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following: Volume: Apply the recommended rate in a spray volume of 30-100 L/ha

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of CHEMINOVA GLYPHOSATE II accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38412 may prevent corrosion.

AGRICULTURAL AND CROPLAND USES

The following are use situations for CHEMINOVA GLYPHOSATE II herbicide. The type of vegetation present and the use situation will dictate the choice of application equipment. Information on the equipment selected to apply CHEMINOVA GLYPHOSATE II can be found in the **APPLICATION EQUIPMENT AND MIXING INSTRUCTIONS** section. Use rates can then be selected from the **ANNUAL** and **PERENNIAL WEED CONTROL** charts.

PREPLANT TREATMENT

CHEMINOVA GLYPHOSATE II can be applied prior to planting of all crops for control of emerged weeds listed on the label. Ensure weeds are at the recommended growth stage at the time of application. Apply BEFORE seeding or transplanting crop.

SUMMER FALLOW

CHEMINOVA GLYPHOSATE II may be applied in summer fallow to control weeds listed on the label. Ensure weeds are at the recommended growth stage and actively growing at the time of application. Reduced control may result if weeds are drought stressed. Repeat treatments may be necessary to control later germinating weeds.

MINIMUM AND ZERO TILLAGE SYSTEMS (ALL FIELD CROPS INCLUDING CEREALS, OILSEEDS, PULSES, FORAGES AND CORN)

CHEMINOVA GLYPHOSATE II may be applied before or after seeding but before crop emerges for control of emerged weeds in minimum and zero tillage cropping systems for all field crops. Weeds should be treated at the growth stage according to the **ANNUAL** and **PERENNIAL WEED CONTROL** charts.

DO NOT APPLY AFTER CROP EMERGENCE.

Since CHEMINOVA GLYPHOSATE II does not provide residual control, application too far in advance of seeding may allow weeds to germinate between application and crop emergence.

MINIMUM AND ZERO TILLAGE TANK MIXES

CHEMINOVA GLYPHOSATE Herbicide II plus Pardner (bromoxynil) can be applied prior to seeding or after seeding, but before crop emergence in **wheat, barley, and oats**. See chart on **TANK MIXES for ANNUAL WEED CONTROL**.

CHEMINOVA GLYPHOSATE Herbicide II plus Pursuit can be applied before or after seeding, but prior to crop emergence in **soybeans**. CHEMINOVA GLYPHOSATE herbicide II will control emerged weeds listed on this label when applied as directed (see **VEGETATION CONTROLLED** lists). Pursuit will control weeds germinating from seed. Add the recommended rates of both products in 100 L of water/ha following the instructions on the Pursuit herbicide label.

Refer to the Pursuit label for further information on weeds controlled, application directions, and use precautions. Only SOYBEANS, FIELD CORN, SPRING BARLEY, SPRING WHEAT and WINTER

WHEAT may be planted the season following a Pursuit application. Winter wheat may be planted the same year as a Pursuit application to soybeans, but not earlier than 120 days after the application.

DO NOT APPLY AFTER CROP EMERGENCE.

CHEMINOVA GLYPHOSATE II TANK MIXES for ANNUAL WEED CONTROL SUMMER FALLOW & MINIMUM TILLAGE SYSTEMS

TANK MIXTURES	RATE L/ha	WEEDS CONTROLLED++	COMMENTS: (Apply in 50-100 L/ha water; add 350 mL/ha
----------------------	------------------	---------------------------	-------------------------------------------------------------

<p>CHEMINOVA GLYPHOSAT E II</p> <p>+</p> <p>Banvel</p>	<p>0.75 - 1.0</p> <p>+</p> <p>0.2</p> <p>9</p>	<p>Volunteer cereals, wild oats, green foxtail, volunteer canola (rapeseed), wild mustard, flixweed*, lamb's-quarters, lady's-thumb, stinkweed, kochia, Russian thistle, cow</p>	<p>This tank mix for summer fallow use only. Weeds should be less than 15 cm tall and actively growing. Use higher rate if weeds are taller than 8 cm.</p> <p>*CHEMINOVA GLYPHOSATE II</p>
<p>CHEMINOVA GLYPHOSAT E II</p> <p>+</p> <p>Pardner</p>	<p>0.75 - 1.0</p> <p>+</p> <p>1.2</p> <p>5</p>	<p>Volunteer cereals, green foxtail, volunteer canola (rapeseed), wild mustard, lady's- thumb, stinkweed, wild buckwheat*</p>	<p>This tank mix for summer fallow use; and prior to planting wheat, oats, and barley in minimum tillage systems. Weeds should be less than 15 cm tall and actively growing. Use higher rate if weeds are taller than 8 cm.</p> <p>*Use CHEMINOVA GLYPHOSATE at 1.0 L/kg rate for wild</p>
<p>CHEMINOVA GLYPHOSAT E II</p>	<p>0.75 - 1.0</p> <p>+</p>	<p>Volunteer cereals, wild oats*, green foxtail*, volunteer canola (rapeseed), wild</p>	<p>This tank mix for summer fallow use only. Weeds should be less than 15 cm</p>

2,4-D#	1.2	flixweed, redroot pigweed, lady's-thumb, stinkweed, kochia Lamb's-quarters**,	are taller than 8 cm. *Use CHEMINOVA GLYPHOSATE II at 1.0 L/ha rate only for wild oat and green foxtail control
<p>#0.56 kg ai/ha of 2,4-D. Adjust rates accordingly for other 2,4-D formulations. Use only low volatile ester or amine formulations of 2,4-D.</p> <p>++ For foxtail barley suppression, refer to chart on ANNUAL WEED CONTROL.</p> <p>NOTE: All CHEMINOVA GLYPHOSATE II herbicide tank mixtures for annual weed control require the addition of a non-ionic surfactant registered for this use, such as</p>			

CHEMINOVA GLYPHOSATE II TANK MIXTURES for PERENNIAL WEED CONTROL, SUMMER FALLOW OR FALL STUBBLE

TANK MIXTURES	RATE L/ha	WEEDS CONTROLLED	COMMENTS: Apply in 100-200 L/ha water; add 350 ml/ha surfactant (see NOTE)
CHEMINOVA GLYPHOSATE + Ranvel	1.7 L/ha + 1.25 L/ha	Canada thistle, perennial sow thistle	Summer fallow: Cultivate in the and apply when majority of thistles to 25 cm tall, and before the bud Fall stubble: Apply to actively growing thistles at least 2 weeks frost.
<p>NOTE: All CHEMINOVA GLYPHOSATE II herbicide tank mixtures for perennial weed control require the addition of a non-ionic surfactant registered for this use, such as Agral 90, AgSurf, or Companion. Grow only cereals, canola (including rapeseed), soybeans, field corn, sweet corn, or white beans after application of this tank mix.</p>			

FALL STUBBLE

Apply in the fall as a postharvest stubble treatment for control of perennial weeds including quackgrass and Canada thistle. Allow the Canada thistle and quackgrass to regrow to 20-25 cm tall. Straw should be removed or evenly spread to allow for proper regrowth and spray coverage. Heavy frost prior to treatment may decrease control.

SPOT TREATMENT (IN CROP)

CHEMINOVA GLYPHOSATE II may be applied for the control of Canada thistle, quackgrass and other perennial weeds in forage crops, barley, wheat, oats, soybeans and legumes, including seed production. Treatments may be made up to heading of small grain, initial pod set on soybeans and legumes, silking of corn, and emergence of seedheads. Avoid drift beyond the treated area.

Application can be made using a boom sprayer, knapsack, or high volume equipment (see **APPLICATION AND MIXING INSTRUCTIONS** section). Applications should be made using the same growth stages as listed in the **ANNUAL** and **PERENNIAL WEED CONTROL** charts. Or, use a 1% solution for annual weeds and quackgrass and a 2% solution for other perennial weeds (a 1% solution equals 1 litre CHEMINOVA GLYPHOSATE II herbicide in 100 litres of spray solution). The 1% and 2% solutions should be applied to wet, but not to run off.

NOTE: THE CROP IN THE TREATED AREA WILL BE KILLED BY THE TREATMENT. DO NOT APPLY IF CROP GROWTH HAS ADVANCED BEYOND SEED SET. ALLOW 3 TO 5 DAYS BEFORE GRAZING IN, OR HARVESTING TREATED AREAS AS

FORAGES.

FORAGE GRASSES AND LEGUMES

Use CHEMINOVA GLYPHOSATE II to control or suppress existing vegetation prior to emergence of legumes and grasses. If legumes and grasses are underseeded with a cover crop, CHEMINOVA GLYPHOSATE II must be applied prior to planting any cover crop.

PASTURE RENOVATION

CHEMINOVA GLYPHOSATE II may be used to control or suppress existing vegetation for zero tillage seeding of legume or grass pasture into established sod for renovation. Weed growth should be at least 20 cm high and most weed seeds should have germinated at the time of spraying.

FORAGE SEED PRODUCTION (FOR SPOT TREATMENT)

CHEMINOVA GLYPHOSATE II may be applied as a spot treatment for control of perennial weeds such as quackgrass and Canada thistle in seed fields. Apply to weeds at least 20-25 cm in height but before emergence of seedhead.

The crop in the treated area will be killed. For this reason, take particular care to avoid drift outside the treated area.

PREHARVEST CONTROL OF QUACKGRASS, CANADA THISTLE, DANDELION, TOADFLAX and MILKWEED; SEASON-LONG CONTROL OF PERENNIAL SOW THISTLE, AND HARVEST MANAGEMENT

For control of quackgrass, Canada thistle, dandelion, toadflax and common milkweed, and season-long control of perennial sow thistle, CHEMINOVA GLYPHOSATE II can be applied prior to harvest of wheat, barley (including malting barley), oats, canola (rapeseed), flax (including low-linoleic acid varieties), lentils, peas, dry beans and soybeans. **DO NOT** apply to crops grown for seed production.

This treatment may also provide harvest management benefits, by drying down crop and weed vegetative growth, for example, where late flushes of annual weeds, green vegetative crop growth, or late tilling may interfere with harvest operations.

EXTREMELY COOL, WET AND/OR CLOUDY WEATHER CONDITIONS BETWEEN THE TIME OF APPLICATION AND THE ANTICIPATED HARVEST DATE MAY SLOW THE ACTIVITY OF THIS PRODUCT, THEREBY DELAYING CROP DRYDOWN AND HARVEST DATE.

CHEMINOVA GLYPHOSATE II should be applied pre-harvest at 2.5 L/ha in 50 to 100 L/ha of clean water, by **GROUND APPLICATION ONLY**.

When to Apply: Apply only when the crop has 30% or less grain moisture content. This stage typically occurs 7 to 14 days before harvest (or 3-7 days for forage applications).

Consult the **GUIDELINES FOR TIMING OF PREHARVEST APPLICATIONS** chart for visible indicators of this stage in each crop. For the best weed control results quackgrass should be actively growing and have at least 4 to 5 green leaves.

Canada thistle and perennial sow thistle should be actively growing and at or beyond the bud stage for best results. Common milkweed should be at the bud to bloom stage and actively growing for best results. Applications for weed control (not for harvest management) must be made at the correct stage of both weed and crop growth.

Apply only during the period 7-14 days before harvest to ensure best weed control and to maximize harvest management benefits. Earlier application may reduce crop yield and/or quality, and may lead to excess glyphosate residues in the crop.

Use Precautions: Overspray or drift to important wildlife habitats such as bodies of

water, wetlands (e.g., sloughs), shelterbelts, woodlots and other cover on the edges of fields frequented by wildlife, should be avoided. Leave a 15 metre buffer zone between the last spray swath and the edge of any of these habitats.

Do not expose or contaminate any body of water or non-target vegetation by direct application, spray drift, or when cleaning and rinsing spray equipment.

DO NOT APPLY BY AIRCRAFT

GUIDELINES FOR TIMING OF PREHARVEST APPLICATIONS

CROP(S)	PERCENT GRAIN MOISTURE	VISIBLE INDICATORS
WHEAT, BARLEY, OATS	Less than 30	Hard dough stage; a thumbnail impression remains on seed.
CANOLA	Less than 30	Pods are green to yellow; most seeds are yellow
FLAX (including low- linoleic acid varieties)	Less than 30	Majority (75%-80%) of bolls are brown.
PEAS	Less than 30	Majority (75%-80%) of pods are brown.
LENTILS	Less than 30	Lowermost pods (bottom 15%) are brown and seeds rattle
DRY BEANS	Less than 30	Stems are green to brown in colour, pods are mature (yellow to brown in colour); 80%-90% leaf drop (original leaves)
SOYBEANS	Less than 30	Stems are green to brown in colour; pod tissue is dry and brown in appearance; 80%-90% leaf

TREE, VINE, AND BERRY CROPS

CHEMINOVA GLYPHOSATE II controls annual and perennial weeds in established vineyards or orchards, in blueberry, cranberry, strawberry and sugar beets, or for site preparation prior to transplanting tree or vine crops. See chart on **WEED CONTROL IN TREE, BERRY, AND VINE CROPS** for rate and time of application information.

This product does not provide residual or pre-emergent weed control. Repeat applications may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. For subsequent weed control, follow a program using residual herbicides or use repeated applications of CHEMINOVA GLYPHOSATE II. **DO NOT APPLY MORE THAN 35 L OF CHEMINOVA GLYPHOSATE II HERBICIDE PER HECTARE PER YEAR.**

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THE HERBICIDE SOLUTION, SPRAY DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, CANES OF BLUEBERRY BUSHES OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURE BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

Allow annual and perennial weeds that have been mowed, grazed, or cut, time to regrow to recommended growth stage for treatment.

Applications may be made with boom sprayer, shielded sprayers, hand-held and high volume orchard guns, or with wiper, wick, or roller equipment (orchards, vineyards, cranberry and strawberry only).

TREE PLANTING - Shelterbelts, Nursery Stock, Woody Ornamentals

CHEMINOVA GLYPHOSATE II may be applied to control annual and perennial weeds listed on this label. This may be used for site preparation prior to establishing plantations, or as a post directed spray in established plantations of the following species:

Deciduous

Ash - *Fraxinus spp.*
Caragana - *Caragana spp.*
Cherry - *Prunus spp.*
Elm - *Ulmus spp.*
Lilac - *Syringa spp.*
Maple - *Acer spp.*
Mountain ash - *Sorbus spp.*
Poplar - *Populus spp.*
Russian olive - *Elaeagnus spp.*
Willow - *Salix spp.*

Juniper

Coniferous

Fir - *Abies spp.*
- *Juniperus spp.*
Pine - *Pinus spp.*
Spruce - *Picea spp.*
Yew - *Taxus spp.*

SPRAY MAY CONTACT MATURE BROWN BARK ONLY.

Avoid contact with non-target plants, foliage, or suckers of established plantations.

NOTE: This product is not recommended for use as an over-the-top broadcast spray in forest tree nurseries or in Christmas tree plantations. Application in such sites should be limited to directed sprays. DO NOT treat Christmas tree plantations in the year of anticipated harvest.

NONCROPLAND AND INDUSTRIAL USES

When applied as recommended under the conditions described, CHEMINOVA GLYPHOSATE II will control weeds in the noncropland and industrial uses as listed in the **WEED CONTROL IN NONCROPLAND, INDUSTRIAL USES** chart.

TURFGRASS

CHEMINOVA GLYPHOSATE II may be applied to control existing vegetation prior to turfgrass establishment or renovation. DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE TREATMENT.

Where existing vegetation is growing under field or unmowed conditions, apply CHEMINOVA GLYPHOSATE II to actively growing weeds at the growth stages given in the charts on **ANNUAL** and **PERENNIAL WEED CONTROL**. Where the vegetation is growing under mowed turfgrass management, apply CHEMINOVA GLYPHOSATE after omitting at least one regular mowing to allow sufficient growth for good spray interception and translocation into underground plant parts.

Tillage or renovation techniques, such as vertical mowing, coring or slicing, should be delayed for 7 days after application to allow proper translocation into the underground plant parts. Delay establishment of the turfgrass to determine if regrowth from escaped underground plant parts occurs. When repeat treatments are necessary, sufficient weed regrowth must be attained prior to application.

AVOID **ALL** CONTACT WITH DESIRABLE VEGETATION IN THE VICINITY OF THE RENOVATION OR ESTABLISHMENT AREA.

TREE INJECTION APPLICATIONS

See **VEGETATION CONTROLLED** lists for species controlled.

Trees may be controlled if CHEMINOVA GLYPHOSATE II is injected directly into the trunk using suitable equipment that penetrates into the living tissue.

CHEMINOVA GLYPHOSATE II is to be used at a rate of 1 mL (undiluted product) per 10 cm of trunk diameter at chest height. The injections should be spaced evenly around the tree and below any major branches. Application may be done during periods of active growth and full leaf expansion.

Control of trees greater than 20 cm may not be acceptable. Total control may not be evident for 1-2 years following treatment. This treatment will only provide suppression of big-leaf maple; late fall application will provide optimum suppression of big-leaf maple.

CUT STUMP APPLICATIONS

See **VEGETATION CONTROLLED** lists for species controlled.

Woody vegetation may be controlled by the application of this product to freshly cut stumps to prevent regrowth. Application must be made using low-pressure equipment (i.e., squirt bottle).

Apply CHEMINOVA GLYPHOSATE II immediately to the surface of the freshly cut stump (i.e., within 5 minutes) at a rate of 0.5 mL CHEMINOVA GLYPHOSATE II for every 5 cm of trunk diameter at chest height. Treat only the cambial tissues (outer edge) of the cut surface. Do not treat the central area of the stump, or exposed roots or bark.

This treatment may be made at any time of year, except during heavy sap flow or when freezing temperatures prevent application of CHEMINOVA GLYPHOSATE II.

A water soluble dye added to the solution may be used as a treatment indicator.

Total control may not be apparent until 1-2 years after treatment.

WOODY BRUSH AND TREES (FOLIAR APPLICATIONS)

Spray coverage should be uniform and complete. Do not spray to the point of run off.

Do not allow spray drift to contact desirable vegetation as severe injury or destruction may occur. For woody brush and trees, early season applications may take 30-45 days for symptoms to develop on the target species. Late season application may be made to species that have some autumn colours provided no major leaf drop has occurred.

Control will be observed the following spring.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

For woody brush and trees, apply 3 to 6 litres of CHEMINOVA GLYPHOSATE II per hectare. Use groundboom or boomless equipment, or apply as a 1 to 2% solution using hand-held high volume equipment. Use the 6 L/ha rate for maple, alder and willow* species, as well as hard to control perennial weed species. (*suppression only).

INDUSTRIAL SITES, RIGHTS-OF-WAY, RECREATIONAL AND PUBLIC AREAS

CHEMINOVA GLYPHOSATE II may be applied to control brush, trees, and annual and perennial weeds listed on this label in **industrial** and **rights-of-way areas**, such as:

railways
highways

forest
roadsides
pumping
stations

pipelines
petroleum tank farms

telephone and power rights-of-way,
etc. and in **recreational** and **public**

areas, such as:

parks, golf courses, schoolyards, airports and other public areas.

NOTE: For all industrial sites, rights-of-way, recreational and public areas, repeat treatment may be necessary to control regeneration or new growth.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Ground application for all noncropland uses: For woody brush and trees, apply CHEMINOVA GLYPHOSATE II at 3 to 6 L/ha using ground boom, or boomless, or mist blower equipment. Or, apply as a 1 to 2% solution using hand-held high volume equipment. Use the higher rate for maple, alder and willow* species, and for hard to

control perennial weeds (*suppression only).

Apply as directed to foliage of actively growing vegetation. Spray coverage should be uniform and complete. Do not spray to the point of runoff, or allow spray drift to contact desirable vegetation as severe injury or destruction may occur.

Mowed or tilled weeds should be allowed to reach optimum growth stage at time of application. **DO NOT APPLY UNDER WIND OR OTHER CONDITIONS THAT ALLOW DRIFT.**

Aerial Application For Industrial Rights-of-Way ONLY: For woody brush and trees, apply 3-6 L/ha CHEMINOVA GLYPHOSATE II in 30-100 L of water. Use 6 L/ha for maple, alder and willow* species, and for hard to control perennial weed species (*suppression only). As density of vegetation increases, spray volume should be increased within the allowed range to ensure complete coverage.

Apply only in wind conditions in compliance with local and/or provincial regulations. Do not apply when other climatic conditions, including lesser wind velocities, will allow significant drift to occur.

Coarse sprays are less likely to drift, therefore, do not use nozzles or nozzle configurations that disperse spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of CHEMINOVA GLYPHOSATE II accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38412 may prevent corrosion.

PURPLE LOOSESTRIFE CONTROL

- **DO NOT TREAT PLANTS OVER OPEN WATER.** CHEMINOVA GLYPHOSATE II herbicide is not registered for direct application to bodies of water.
- Treat when plants are actively growing at or beyond the bloom stage. If using hand-held equipment, spray-to-wet.
- For wiper applications, see **WIPER, WICK AND ROLLER EQUIPMENT** section.
- Where feasible, remove flower heads before treatment to ensure prevention of seed set.
- For large (>1.6 ha) monocultures of loosestrife, work from the periphery inward in successive years to allow competing vegetation to invade the treated area.
- A long-term control strategy should include measures to control both established plants and seedlings. Sprayed areas should be monitored to determine the appropriate follow-up management. Early detection and treatment of second and third generation seedlings is important to prevent re-infestation of purple loosestrife. Desirable native plant communities will then have a chance to become re-established.

WEED CONTROL IN NONCROPLAND, INDUSTRIAL USES

WEEDS	GROUND APPLICATION			COMMENTS
	BOOM APPLICATION		Hand-Held High Volume Application	
	Rate	Water Vol		
Annual grasses and	2.25 - 3.5	50 - 100	1	Actively growing weeds
Perennial weeds	2.5	50 - 300	1	Actively growing weeds. Add 0.5% v/v of a recommended surfactant when using more than 150 L of water (see MINIMUM AND ZERO TILLAGE TANK MIXES). Use higher rate for heavy infestations and for long term control. See PURPLE LOOSESTRIFE CONTROL section for
Quackgrass	4.75 - 7.0	50 - 300	2	
Canada thistle (bud stage)	4.75 - 7.0	100 - 300	2	
Purple loosestrife	6.0	300 - 600	1-2 (or 33% for wiper application)	
Brush and trees	3.0 - 6.0	100 - 300	1 - 2	Summer through early fall.
Birch, cherry, poplar, western snowberry, willow	6.0	100 - 300	2	Late summer through fall. Fall is optimum.
Maple,				
Turfgrass renovation Annual & perennial	2.5 - 12.0	100 - 300	1 - 2	Use higher end of rate range for perennials.

Roadside vegetation (1-2 metres wide along shoulder)	1) 0.75 - 1.0 + 1.25 - 2.5 L DyClear 480 or 2) 0.75 - 1.0 + 0.30 L DyClear 480	25 - 150	-	Refer to Tank Mix sections on product labels for specific weeds controlled. Refer to chart on ANNUAL WEED CONTROL for rates for specific weeds. For different 2,4-D amine formulations, adjust the rate accordingly. Do not apply to standing water.
Residual Control Annual & perennial weeds	2.5 - 12 + 1) 2.5 - 5.6 kg Simazine 80W or 2) 4.0 - 9.0 L Simadex Flowable	200 - 400	-	The simazine part of this tank mix will provide season-long control of most germinating broadleaf weeds and grasses, and may also provide post-emergent control of certain annual weeds. Do not apply to coarse, sandy soil or gravelly soil. One application per year. Use the most restrictive label directions for each product in the mix.

WEED CONTROL IN TREE, VINE AND BERRY CROPS

Crop	Rate (L/ha)	Pre-Harvest	Max. App. per Year	Weed Controlled	Comments
Apples Apricot Cherry (sweet/sour) Peaches	2.25 - 12 (directed spray)	30	3	Annual and perennial weeds	Apply as directed spray with no more than 275 kPa pressure.
Apples Grapes	Tank Mix 2.25 - 12 + Simazine 2.0 - 4.5 kg ai/ha	-	1	Annual and perennial weeds	Will provide season-long pre-emergent control. Do not apply to coarse, sandy or gravelly soil. Use the more restrictive label directions for each product in the mix. DO NOT apply to orchards established less than 1 year
Grapes	2.25 - 12 (directed spray)	14	3	Annual and perennial	Remove all sucker growth from the spray zone before spraying, except for the

					grape. Suckering should be conducted within 2 weeks prior to application. Do not apply to vines that have
Highbush blueberry (cultivated)	2.8 - 5.6 (directed spray)	30	1	Quackgrasses	Use as a directed spray, with no more than 275 kPa pressure
Lowbush blueberry	1 - 2% solution (spot treatment)	Apply in non-bearing year only	1	Wood	Apply as directed spray in summer of the vegetative bearing year. See AGRICULTURAL AND CROPLAND USES section instructions on spot
Filberts Hazelnut (established plantations)	2.25 - (directed spray)	14	-	Annual weeds	Use as directed spray, with more than 275 kPa
Walnut Chestnut Japanese chestnut	2.25 - 12 (directed spray)	-	2	Annual perennial weeds	Apply late spring and fall, harvest but prior to a killing Apply in 200 - 300 L water directed spray, using no than 275 kPa pressure, alternatively as a 2% wiper solution. See EQUIPMENT AND MIXING INSTRUCTIONS section for instructions on wiper
Cranberry	20% Solution L CHEMIN O VA ATÉII L water)	30	1	Annual perennial weeds	Apply using wick or wiper applicators. See EQUIPMENT AND MIXING INSTRUCTIONS section for instructions on wiper
Strawberry	1 - 2% solution (spot treatment) 33% solution (wiper applicator)	30	1	Emerging perennial weeds	Apply when weeds are at a susceptible growth stage. AGRICULTURAL AND CROPLAND USES section instructions on spot See APPLICATION EQUIPMENT AND MIXING INSTRUCTIONS section for instructions on wiper applications.
Sugar beets	1 - 2% solution (spot treatment)	Treated crop MUST NOT be harvested	1	Dodder species	Apply when dodder is growing but before AGRICULTURAL AND CROPLAND USES section instructions on spot

ANNUAL WEED CONTROL

Equipment	Weeds Controlled	Growth	Rate	Water Vol	Comments
Boom or boomless	Wild oats, green foxtail, volunteer barley, volunteer wheat,	Weeds up to 8 cm in height	0.75	50 - 100	For wild oats apply at 1 to 3 leaf stage. Add 350 mL of a surfactant registered for use such as Agral 90, AgSurf, and Companion. For heavy wild oat infestations use 1.0
	All annual grasses listed above plus foxtail barley* (suppression only) All annual broadleaf weeds listed above plus flixweed** and kochia**	Weeds 8 cm to 15 cm	1.0	50 - 100	Add 350 mL of surfactant registered for use as listed above. * Apply before initiation of seedhead or senescence of the lower leaves. ** Suppression only. Refer to higher rates of this table.
	All annual grasses listed above plus downey brome, giant foxtail and Persian darnel. All annual broadleaf weeds listed above plus lamb's-quarters, redroot pigweed, hempnettle, flixweed, Russian	Weeds up to 15 cm in height	1.25 - 1.9	50 - 100	No additional surfactant required. * DO NOT use these rates on plants greater than 8 cm in height. ** For 3 to 4 leaf stage use 1.9 L/ha rate. *** For weeds 8 cm to 15 cm in height use 1.9 L/ha.
	All annual grasses listed above plus crab grass and annual bluegrass. All annual broadleaf weeds listed above plus kochia, prickly lettuce	Weeds up to 15 cm in height	2.25	50 - 100	

	All annual grasses and broadleaf	Weeds over 15 cm in	3.5	50 - 100	
Wipers and wicks	Annual weeds	Weeds to be at least 15 cm above desirable vegetation	1	2	This mixture is a 33% solution. Contact point for wiper or wick must be at least 5 cm above desirable vegetation. In severe weed infestations, reduce ground speed to ensure adequate control. See APPLICATION EQUIPMENT AND MIXING INSTRUCTIONS
Rollers	Annual weeds	Weeds to be at least 15 cm above desirable vegetation	0.5 - 1.0	10	This mixture is a 5-10% solution. Roller speed 50-150 rpm. See APPLICATION EQUIPMENT AND MIXING INSTRUCTIONS section for instructions on

PERENNIAL WEED CONTROL

Equipment	Weeds Controlle	Growt h	Rat e	Water Vol.	Commen ts
Boom or boomless	Fall application (after harvest)	3-5 green leaves (approx. 20 cm height)	2.5 2.5 - 7	50 - 300 50 - 300	For season-long control the following year. Do not till between harvest and application. Allow 5 days or more after application before tillage. Long Term Control: Reduced control may result if rhizomes have become dormant due to poor sod or land has not been tilled for several years. Treatment after a mild frost is possible if 3 to 4 leaves are
	Alfalfa	Early bud to full bloom stage. Fall applications only.	3.7 - 5	50 - 300	Allow 5 or more days after treatment before tillage. Use the higher rates when alfalfa populations are high or when heavy grass infestations are also present. Tank mix with

Canada thistle	Bud stage or beyond	4.75 - 7	100 - 300	Allow 5 days after application before tillage. Heavy frost prior to application may
Canada thistle	Rosette stage (summer fallow)	2.5	50 - 100	Ensure proper growth stage by performing last summer fallow tillage between July 5 and August 1st. Allow regrowth for a minimum of 5 weeks to reach rosette stage and a minimum of 15 cm in diameter. Allow 10 days after application before tillage. Treatment after a mild frost is
Dandelion	Up to 15 cm in height	2.5	50 - 100	Allow 3 or more days after treatment before tillage for all rates. Use the higher rates when infestations are heavy. Allow 7 or more days after treatment before tillage.
	Over 15 cm in height	3.7 - 5	50 - 300	
	Rosette to full bloom	2.5		
Field bindweed	Full bloom or beyond	7 - 12	100 - 300	Allow 7 days or more after application before tillage
Foxtail barley	Seedling to heading	2.5 - 5	50 - 300	Allow a minimum of 1 day after treatment before tillage or seeding. Use higher rates for larger more established plants, heavy infestations, or
Common Milkweed	Bud to full bloom for most shoots	12	100 - 300	Spot treatment rate is 120 mL per 5 L water/100 m ² and spray to wet not runoff. Reduced results may occur if sprayed after full bloom. Repeat treatment may be
Toadflax	Vegetative stage (summer fallow), bud to full bloom (pre-	2.5	50 - 100	Apply in clean water using flat fan nozzles. Allow 7 or more days after treatment before tillage in summer fallow. For higher water volumes 150-300L an
Quackgrass control, light to moderate infestations	3 to 4 green leaves or more	2.5	50 - 300	Apply in clean water using flat fan nozzles. Allow 3 or more days after treatment before tillage. For higher water volumes 150-300L an approved surfactant must be added at

	Quackgrasses (long term control, heavy infestation)	4 to 5 green leaves or more	2.5 - 7	50 - 300	Allow 3 or more days after treatment before tillage. Rates higher than 2.5 L will provide more consistent, longer term control especially with heavy infestations
	Quackgrasses - spring application (no fall)	3 to 4 green leaves (approx. 20 cm height)	2.5	50 - 300	Season-long control. At higher water volumes use approved surfactant at 0.5% v/v (0.5 L per 100 L water).
	Quackgrasses - spring application (fall-)	4 to 5 green leaves (approx. 20 cm height)	2.5	50 - 100	Season-long control. Apply in spring prior to seeding. Growth stage usually reached 1 to 4 weeks later on land that has been fall-
	Other perennial weeds	Early heading or early bud stage (See VEGETATION)	7 - 12	100 - 300	Use higher rate for weeds beyond 8 cm in height or in heavy weed infestation. Allow 7 days after application before tillage. CHEMINOVA GLYPHOSATE
	Woody brush and trees	Actively growing from June	3 - 6	100 - 300	Use higher rate for maple, alder, <i>Rubus</i> species and willow*. Spray to wet.
High volume or knapsack	Woody brush and trees	Actively growing from June through August	1 - 2.0	10 - 0	This mixture is a 1 to 2% solution. Use higher rate for maple, alder, <i>Rubus</i> species and willow*. Spray to wet. See APPLICATION EQUIPMENT AND MIXING INSTRUCTIONS section for
Wipers and wicks	Perennial weeds	Weeds to be at least 15 cm above desirable	1	2	See APPLICATION EQUIPMENT AND MIXING INSTRUCTIONS section for instructions on
Rollers	Annual and perennial weeds	Weeds to be at least 15 cm above desirable vegetation	0.5 - 1.0	10	This mixture is a 5-10% solution. See APPLICATION EQUIPMENT AND MIXING INSTRUCTIONS section for instructions on roller applications. <i>This treatment will only</i>
Tree injection	Trees*	During periods of active growth and	1 mL per 10 cm of	None	Suitable equipment must be used to penetrate to living tissue. Space applications evenly

		periods of heavy sap flow.	er at chest heigh		Control of trees with trunk diameters greater than 20 cm not be acceptable. See APPLICATION AND MIXING section for instructions on INJECTION APPLICATIONS.
--	--	----------------------------	-------------------	--	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------

* Suppression only for willow.

TMCHEMINOVA GLYPHOSATE II and FMC are trademarks of FMC Corporation or an affiliate. All other products listed are registered trademarks or trademarks of their respective companies.