

GEFG3001  
GEFG10-31A

**Wilson® WeedOut® Preventer**

Pre-emergent Dandelion, Crabgrass (smooth and large) and  
White Clover Weed Seed Germination Inhibitor with Corn Gluten

For the inhibition of dandelion and crabgrass (smooth and large) and white clover seed  
germination in lawns where established perennial ryegrass and Kentucky bluegrass predominate.

DOMESTIC

REG. NO. 30495 P.C.P. ACT

GUARANTEE: liquid corn gluten.....99.81%  
Contains potassium sorbate at 0.19% as a preservative

CAUTION EYE IRRITANT  
POTENTIAL SENSITIZER

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

*1 L to 10 L*

Sure-Gro IP Inc. /  
Lic. Premier Tech Home & Garden Inc.,  
P.O. Box 21001  
Brantford, ON N3R 7W9  
1-800-268-2806

Wilson® WeedOut® Preventer may inhibit dandelion, crabgrass (smooth and large) and white clover seed germination when used in conjunction with a sound lawn maintenance program. Established dandelion, crabgrass (smooth and large) and white clover at the time of application will not be inhibited. This product is formulated for use in residential lawns where established perennial ryegrass and Kentucky bluegrass are the predominant grass species.

**DIRECTIONS FOR USE:** Apply this product to dry or moist soil. Do not apply to excessively saturated soil. Do not apply if rain is forecasted within 24 hours after application. Do not irrigate for 24 hours after application. Excessive moisture at the time of application may reduce the effectiveness of the product. Do not apply under windy conditions. Application dates may vary for both spring and late summer/early fall applications from year to year according to weather conditions. **DO NOT** apply to any body of water. **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. Not recommended for use with a fertilizer containing nitrogen.

This product is to be applied to a mature lawn having a well developed root mass. Do not apply the product on newly seeded grass as it may inhibit seeds from germinating, wait until after first mowing when root systems are established. If over-seeding or re-sodding in the spring, do not apply the product in the spring. If overseeding or re-sodding in the fall, do not apply the product in the fall. For best results: Apply to established turf twice a year; once in the early spring and once in the late summer/early fall.

**SPRING APPLICATION:** Dandelion, Crabgrass (smooth and large) and White Clover Seed Germination Inhibition: Apply to established perennial ryegrass and Kentucky bluegrass turf in early spring before crabgrass (smooth and large) seed germination.

**LATE SUMMER/EARLY FALL APPLICATION:** Dandelion, Crabgrass (smooth and large) and White Clover Seed Germination Inhibition: Apply to established perennial ryegrass and Kentucky bluegrass turf in late summer or early fall after heat stress has passed.

**APPLICATION RATE:**

Apply at a rate of 1.08 L / 100 m<sup>2</sup> At this rate:

1 L container is calibrated to treat 93 m<sup>2</sup> (1,000 ft<sup>2</sup>)

2 L container is calibrated to treat 186 m<sup>2</sup> (2,000 ft<sup>2</sup>)

4 L container is calibrated to treat 372 m<sup>2</sup> (4,000 ft<sup>2</sup>)

5 L container is calibrated to treat 465 m<sup>2</sup> (5,000 ft<sup>2</sup>)

10 L container is calibrated to treat 930 m<sup>2</sup> (10,000 ft<sup>2</sup>)\*

\* No hose and sprayer included with 10 L container. See instruction below for more details.

**METHOD OF APPLICATION:**

**READ THE COMPLETE LABEL FOR APPLICATION INSTRUCTIONS AND PRECAUTIONS PRIOR TO USE**

**INSTRUCTIONS FOR HOSE END ATTACHMENT CONTAINERS**

1. Shake container thoroughly, then attach to garden hose.
2. Turn water supply on at the source. Water pressure will build in hose since the sprayer is already in the "OFF" position.
3. To apply product, bend safety tab and turn the control valve clockwise to the "ON" position.
4. For best results, hold sprayer and bottle level, and direct spray in a sweeping side-to-side movement. Spray the contents of bottle evenly over the lawn surface. Refer to the "Application Rate" section of this label for the application rate and treatment areas. Do not overlap areas already sprayed.
5. To spray water only, turn the control valve clockwise so that the valve indicator arrow is pointing to "WATER" position on the sprayer.

6. When spraying is complete, turn the control valve to “OFF” position. Turn water off. Disconnect hose from applicator and store product in original container.

#### INSTRUCTIONS FOR 10 L CONTAINER REQUIRING HAND-HELD OR BACKPACK SPRAYER

Open container and slowly pour 1 part Wilson® WeedOut® Preventer into any standard hand-held or backpack sprayer containing 3 parts water (example: pour 1 L product into 3 L water). Apply the mixed solution at a rate of 43 mL/m<sup>2</sup>. At this rate, a 10 L container of product will treat 930 m<sup>2</sup> (10,000 ft<sup>2</sup>). Direct the spray in a sweeping side-to-side movement over the lawn surface. Do not overlap areas already sprayed.

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.** May cause sensitization. This product should not be applied if a member of the household has a sensitivity or allergy to corn. May irritate eyes and skin. Avoid contact with skin, eyes or clothing. May cause respiratory irritation. Avoid breathing spray mist. For good hygiene practice, wear a long-sleeved shirt, long pants, closed footwear and gloves when handling the product. Use freshly prepared spray solution each time when using handheld or backpack sprayers. Do not re-enter or allow re-entry of adults, children or domestic pets into treated areas until the spray is dried.

**FIRST AID: If swallowed,** call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If inhaled,** move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. **If in eyes,** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical advice.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**STORAGE:** Keep in cool, dry conditions, away from seed, fertilizer and other pesticides. To prevent contamination, store this product away from food or feed.

**DISPOSAL:** DO NOT reuse the empty containers. Dispose in household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal site.

**For additional product information or in case of emergency, spills or fire call toll-free 1-800-268-2806.  
[www.pthomeandgarden.com](http://www.pthomeandgarden.com)**

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.