

2019-3292
2020-01-07

This product has been cancelled and has expired as of December 22, 2021, but is authorized under paragraph 21(5)(a) of the Pest Control Products Act. Manufacturing and importing of the product are no longer permitted. The existing stocks in Canada are to be depleted according to the phase-out schedule, as listed in the Pesticide Product Information Database (<https://pesticide-registry.canada.ca>).



GROUP	4	INSECTICIDE
-------	---	-------------

STRESS SHIELD® for cereals

Stress Shield for cereals is a systemic seed treatment insecticide for protection of wheat, barley and oats against wireworm

Insecticide

COMMERCIAL

Suspension

ACTIVE INGREDIENT: Imidacloprid 480 g/L

REGISTRATION NO. 29609 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 0.25 – 1,000 L

BAYER CROPSCIENCE INC.
200-160 Quarry Park Blvd, SE
Calgary, Alberta T2C 3G3

24 HOUR EMERGENCY PHONE: 1-800-334-7577

GROWER INFORMATION: 1-888-283-6847

SPECIAL USE RESTRICTIONS: This product contains no colourant. Seed treated with Stress Shield for cereals must be conspicuously coloured. Regulations pertaining to the *Seeds Act* must be strictly adhered to when using this product.

® Stress Shield is a registered trademark of Bayer CropScience

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapour or spray mist. When handling Stress Shield for cereals or seed treated with Stress Shield for cereals, work in a well-ventilated area and wear a long-sleeved shirt, long pants, chemical-resistant gloves, and shoes plus socks. DO NOT use leather or cloth gloves. Wear goggles and a suitable dust mask approved by NIOSH/MSHA when handling this product. Use good personal hygiene, washing hands and exposed skin before eating, drinking or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed	Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS: This product is highly toxic to birds and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present. Do not contaminate water when disposing of equipment wash waters. Cover or incorporate spilled treated seeds. Left over treated seed should be double-sown around the headland, or buried away from water sources.

Toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatment applications. When used according to label directions minimal exposure or risk is expected. Dust generated during planting of treated seed may be harmful to bees and other pollinators. To help minimize the dust generated during planting, refer to the “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators . Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds. When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies. Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep container closed.

Do not store Stress Shield for cereals in direct sunlight.

Do not store Stress Shield for cereals above 35°C.

DISPOSAL OF CONTAINER

FOR NON-RETURNABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS FOR CONTAINER DISPOSAL (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment provided the total volume of water does not exceed 6 % of the container size. Add 1/3 of the rinse water to the container and swish the

contents thoroughly. Empty the rinsate into the seed treater holding tank and repeat this process two more times making sure the total volume of rinsate does not exceed 6%. Make sure rinsate is thoroughly mixed with seed treatment before treating. Close the container.

CAUTION: Do not dilute beyond 6% or excess seed wetness may result in seed handling difficulties. Using excess water can also cause Stress Shield for cereals to thin resulting in settling of solids in the product. Be sure to adjust the undiluted application rate up by 6% to compensate for the dilution of the product caused by the addition of rinsate.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

® Stress Shield is a registered trademark of Bayer CropScience



GROUP 4 INSECTICIDE

STRESS SHIELD® for cereals

Stress Shield for cereals is a systemic seed treatment insecticide for protection of wheat, barley and oats against wireworm

Insecticide

COMMERCIAL

Suspension

ACTIVE INGREDIENT: Imidacloprid 480 g/L

REGISTRATION NO. 29609 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 0.25 – 1,000 L

BAYER CROPSCIENCE INC.
200-160 Quarry Park Blvd, SE
Calgary, Alberta T2C 3G3

24 HOUR EMERGENCY PHONE: 1-800-334-7577

GROWER INFORMATION: 1-888-283-6847

SPECIAL USE RESTRICTIONS: This product contains no colourant. Seed treated with Stress Shield for cereals must be conspicuously coloured. Regulations pertaining to the *Seeds Act* must be strictly adhered to when using this product.

® Stress Shield is a registered trademark of Bayer CropScience

Table of Contents:	Section Number
GENERAL INFORMATION	Notice to User.....1
	The Product.....2
SAFETY AND HANDLING	Precautions, Protective Clothing and Equipment.....3
	First Aid and Toxicological Information.....4
	Environmental Precautions and Information.....5
	Storage.....6
	Disposal.....7
DIRECTIONS FOR USE	Crops and Pests.....8
	Wheat, Barley, Oat.....8.1
	Tank Mixtures.....8.2
	Use Restrictions and Limitations.....9
	Use Restrictions.....9.1
	Seed Quality.....9.2
	Labeling Treated Seed.....9.3
	Special Use Restrictions.....9.4
	Mixing Instructions.....10
	Resistance Management Recommendations.....11

GENERAL INFORMATION

SECTION 1: NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

SECTION 2: THE PRODUCT

Stress Shield for cereals is a systemic seed treatment insecticide that provides early season protection of wheat, barley and oats from wireworms. Thorough seed coverage is required for maximum protection of seed. Use the higher rate when insect pressure is expected to be high. Under high insect pressures, a foliar insecticide may be required, therefore monitor crops regularly for insect infestation levels. Do not apply any subsequent application of a Group 4 Insecticide (i.e. in-furrow or foliar application) following treatment with Stress Shield for cereals.

SAFETY AND HANDLING

SECTION 3: PRECAUTIONS, PROTECTIVE CLOTHING AND EQUIPMENT

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapour or spray mist. When handling Stress Shield for cereals or Stress Shield for cereals-treated seed, work in a well-ventilated area and wear a long-sleeved shirt, long pants, chemical-resistant gloves, and shoes plus socks. DO NOT use leather or cloth gloves. Wear goggles and a suitable dust mask approved by NIOSH/MSHA when handling this product.

Use good personal hygiene, washing hands and exposed skin before eating, drinking or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

SECTION 4: FIRST AID AND TOXICOLOGICAL INFORMATION

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed	Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

SECTION 5: ENVIRONMENTAL PRECAUTIONS AND INFORMATION

This product is highly toxic to birds and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present. Do not contaminate water when disposing of equipment wash waters. Cover or incorporate spilled treated seeds. Left over treated seed should be double-sown around the headland, or buried away from water sources.

Toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatment applications. When used according to label directions minimal exposure or risk is expected. Dust generated during planting of treated seed may be harmful to bees and other pollinators. To help minimize the dust generated during planting, refer to the “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators. Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds. When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies. Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

SECTION 6: STORAGE

Store product in original container only, away from other pesticides, fertilizer, food or feed.
Keep container closed.

Do not store Stress Shield for cereals in direct sunlight.

Do not store Stress Shield for cereals above 35°C.

SECTION 7: DISPOSAL

DISPOSAL OF CONTAINER

FOR NON-RETURNABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

2. Triple- or pressure-rinse the empty container. Add the rinsings to mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS FOR CONTAINER DISPOSAL (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment provided the total volume of water does not exceed 6 % of the container size. Add 1/3 of the rinse water to the container and swish the contents thoroughly. Empty the rinsate into the seed treater holding tank and repeat this process two more times making sure the total volume of rinsate does not exceed 6%. Make sure rinsate is thoroughly mixed with seed treatment before treating. Close the container.

CAUTION: Do not dilute beyond 6% or excess seed wetness may result in seed handling difficulties. Using excess water can also cause Stress Shield for cereals to thin resulting in settling of solids in the product. Be sure to adjust the undiluted application rate up by 6% to compensate for the dilution of the product caused by the addition of rinsate.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DIRECTIONS FOR USE

SECTION 8: CROPS AND PESTS

8.1 DURUM WHEAT, WINTER WHEAT, SPRING WHEAT, BARLEY & OAT

To provide early season protection against crop stand injury caused by wireworm apply Stress Shield for cereals at 21-63 ml/100 kg of seed. For fields with a history of moderate to high wireworm pressure, treat crops at 42 – 63 ml / 100 kg seed. Use the higher rate when infestation pressures are expected to be heavy. Do not apply any subsequent application of a Group 4 Insecticide (i.e. in-furrow or foliar application) following treatment with Stress Shield for cereals.

For use in commercial and on-farm seed treatment equipment. Mix thoroughly before use, or use entire container at one time. All seed must be conspicuously coloured at the time of treatment in accordance with *Seed Act and Regulations*. Stress Shield for cereals-treated seed may reduce seed flow in the seed drill. Recalibration of the seed drill may be required to obtain correct seeding rate before planting.

Crop	Insect Pest	Use Rate / 100 kg seed		Remarks
		ml product	grams a.i.	
Wheat (Durum, spring, winter), Barley, Oats	Wireworms	21 – 63	10 – 30	Dilute in sufficient liquid to achieve uniform distribution on the seed.

Pre-test the germination of a small sample of each seed lot with Stress Shield for cereals prior to commercial application to the whole lot. Stress Shield for cereals can be used as an over-treatment.

8.2 TANK MIXTURES WITH FUNGICIDES

For control of certain seed and soil-borne pathogens in wheat, barley and oat seed and seedlings, Stress Shield for cereals may be mixed with Raxil T or Raxil MD Seed Treatment Fungicides. Follow all appropriate directions and precautions as specified on the fungicide labels. Do not tank mix Stress Shield for cereals with pesticides, fertilizers or any other chemical additives unless recommended on this label.

Crop	Seed Treatment Fungicide	Application Rate ml product / 100 kg
Wheat (spring, winter & durum), barley, oats	Raxil T	225

Wheat (spring, winter & durum), barley, oats	Raxil MD	300
--	----------	-----

SECTION 9: USE RESTRICTIONS AND LIMITATIONS

9.1 USE RESTRICTIONS:

Do not use treated seed for food or feed.

Do not graze or feed livestock on treated areas for four weeks after planting.

9.2 SEED QUALITY:

Treatment of highly mechanically damaged seed, or seed of known low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. If seed lot quality is unknown, conduct a germination test on a small portion of seed before committing the total seed lot to a selected chemical treatment.

9.3 LABELLING TREATED SEED:

All treated wheat, oat and barley seed for sale or use in Canada must be labelled with the following information:

- This seed has been treated with Stress Shield for cereals seed protectant which contains imidacloprid.
- Do not use for feed, food or oil processing. Store away from feeds and other foodstuffs.
- Imidacloprid is toxic to bees. Dust generated during planting of treated seed may be harmful to bees and other pollinators.
- To help minimize the dust generated during planting, refer to the “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators .
- Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds.
- When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.
- Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

9.4 SPECIAL USE RESTRICTIONS:

This product contains no colourant. Seed treated with this product must be conspicuously coloured. Regulations pertaining to the *Seeds Act* must be strictly adhered to when using this product.

SECTION 10: MIXING INSTRUCTIONS

Storage of Stress Shield for cereals at low temperatures is not recommended. Prior to and during application, Stress Shield for cereals must be thoroughly agitated to ensure uniform mixing of the product. Due to the viscosity of the material, it should be kept above 10 °C prior to and during application. Do not apply direct heat to container.

SECTION 11: RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that Stress Shield for cereals contains a Group 4 insecticide. Any insect population may contain individuals naturally resistant to Stress Shield for cereals and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of Stress Shield for cereals or other Group 4 insecticides with different groups that control the same pests in a field.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Bayer CropScience Inc. via internet at www.cropscience.bayer.ca or phone 1-888-283-6847.

® Stress Shield is a registered trademark of Bayer CropScience