

BASE

GROUP	4	HERBICIDE
-------	----------	-----------

OPTICA™ TRIO

Solution

COMMERCIAL

**BROADLEAF HERBICIDE FOR USE IN WHEAT (Spring, durum and winter),
BARLEY, OATS AND NON-CROP LAND/INDUSTRIAL SITES**



CAUTION

POISON

DANGER - CORROSIVE TO EYES

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING.
PROTECT FROM FREEZING.**

Warning, contains the allergen, soy

ACTIVE INGREDIENT:

MCPA, present as dimethylamine salt . . . 160 g/L
Dichlorprop-P, present as dimethylamine salt . . . 310 g/L
Mecoprop-P, present as dimethylamine salt . . . 130 g/L

REGISTRATION No.: 29662 PEST CONTROL PRODUCTS ACT

Nufarm UK Limited
Wyke Lane, Wyke, Bradford
West Yorkshire, BD12 9EJ
England

Canadian Agent:
Nufarm Agriculture Inc.
5101, 333 - 96th Ave NE
Calgary, Alberta T3K 0S3

For product use information, please contact:
United Agri Products Canada, Inc.
789 Donnybrook Drive
Dorchester, ON
N0L 1G5
1-800-265-4624

24 Hour Emergency Response Number 1-800-424-9300

NET CONTENTS 10 L

12-2048

OPTICA™ TRIO

Solution

COMMERCIAL

**BROADLEAF HERBICIDE FOR USE IN WHEAT (Spring, durum and winter),
BARLEY, OATS AND NON-CROP LAND/INDUSTRIAL SITES**



CAUTION

POISON

DANGER - CORROSIVE TO EYES

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING.
PROTECT FROM FREEZING.**

Warning, contains the allergen, soy

ACTIVE INGREDIENT:

MCPA, present as dimethylamine salt . . . 160 g /L
Dichlorprop-P, present as dimethylamine salt . . . 310 g/L
Mecoprop-P, present as dimethylamine salt . . .130 g/L

REGISTRATION No.: 29662 PEST CONTROL PRODUCTS ACT

Nufarm UK Limited
Wyke Lane, Wyke, Bradford
West Yorkshire, BD12 9EJ
England

Canadian Agent:
Nufarm Agriculture Inc.
5101, 333 - 96th Ave NE
Calgary, Alberta T3K 0S3

For product use information, please contact:
United Agri Products Canada, Inc.
789 Donnybrook Drive
Dorchester, ON
N0L 1G5
1-800-265-4624

24 Hour Emergency Response Number 1-800-424-9300

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Store with container tightly closed, and away from seeds, fertilizers, plant food-stuffs and greenhouses. Protect from freezing. Avoid spray drift on to susceptible crops. Wash sprayer thoroughly, before and after use. After use wash hands and skin contaminated by spray or concentrate.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

If this pest control product is to be used on a commodity that may be exported to the U. S. and you require information on acceptable residue levels in the U. S., visit CropLife Canada's website at: www.croplife.ca.

PROTECTIVE CLOTHING AND EQUIPMENT (PPE):

The following PPE is for all agricultural uses. For PPE required when treating non-crop land/industrial sites, use the PPE listed in the "Non-Crop Lands and Industrial Sites" section.

Handling the concentrate (mixing and loading): Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (face shield or safety goggles). Rinse gloves before removal. Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5L.

Handling the dilute spray solution (during application or repairing or cleaning equipment): Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor.

If OPTICA TRIO is applied when crop is under stress, shortening of straw may occur, but yield will not be affected. Protect from freezing.

Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 12 hours after application for agricultural uses.

Do not enter or allow entry into treated non-crop land/industrial sites until residues have dried.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

ENVIRONMENTAL HAZARDS:

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loamy soils) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient.

High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low-blood pressure and unconsciousness. Treat symptomatically.

EMERGENCY TELEPHONE NUMBERS:

For spills or transportation accidents, Chemtrec, 1-800-424-9300.

For health or environmental emergencies, ProSAR, 1-877-325-1840.

For product and use information, UAP Canada Inc. 1-800-265-4624.

STORAGE:

Protect from freezing

DISPOSAL:**For Recyclable Containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site.

Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Returnable-Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE:

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 12 hours after application for agricultural uses.

Do not enter or allow entry into treated non-crop land/industrial sites until residues have dried.

DO NOT APPLY BY AIR

DO NOT APPLY BY MECHANICALLY-PRESSURIZED HANDHELD EQUIPMENT

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management OPTICA TRIO is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to OPTICA TRIO and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of OPTICA TRIO or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage, cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact UAP Canada Inc. at 1-800-265-4624 or at www.uap.ca.

GENERAL INFORMATION

For systemic control of many broadleaf weeds in: Wheat (spring, durum and winter), Barley and Oats and Non-cropland and Industrial sites.

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 7 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Observe a minimum plant-back interval of 30 days.

BUFFER ZONES:

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer*	Wheat, barley, oats, non-cropland/industrial sites	1	1	1	1	2

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

WEEDS CONTROLLED:

1.5 L/ha

- Stinkweed
- Wild Mustard
- Lamb's Quarters
- Volunteer Canola

2.5 L/ha

- Pigweed (Redroot)
- Common chickweed
- Kochia
- Common Ragweed
- Wild Buckwheat
- Cleavers¹
- Lady's Thumb (suppression only)

Best results are obtained when weeds are in the 2-3 leaf stage and actively growing. Less than satisfactory control may result if weeds are not actively growing.

Notes:

Top Growth Suppression Only - 2.5 L/ha:

Canada thistle

Notes:

Cleavers¹: Spray in 1-2 whorl stage

TIMING OF APPLICATION:

CROP	CROP STAGE
Wheat (spring, durum) Oats Barley	2 to 5 leaf
Winter Wheat	For Winter Wheat, apply in the spring before the crop is more than 30 cm high (top leaf extended). Winter annual weeds should be sprayed as soon as new growth starts.
Non-cropland /industrial sites	See notes above for specific weed staging recommendations.
Note: Application made later than the recommended timing may result in shortening of the straw (particularly under stress conditions).	

VOLUME: Apply the recommended rate in a minimum spray volume of 50 litres per hectare.

APPLICATION INFORMATION:

Rate 1.5 – 2.5 litres per hectare depending on broadleaf weeds that are present. Use 50-200 L/ha of water by ground.

Do not apply more than one treatment per year.

For Non-Crop Lands and Industrial Sites

(e.g. for use on rights-of-way for transportation, rights-of-way for utility lines, and in airports, wastelands, industrial parks, etc.):

1. Treat weeds when they are in the 2-3 leaf stage and actively growing.
2. SPOT TREATMENT: For knapsack application of Optica Trio for spot treatment of weeds such as thistles, mix 50 mL of Optica Trio in one litre of water. For easy-to-control weeds such as stinkweed, wild mustard, lamb’s quarters and volunteer canola, mix 30 mL of Optica Trio in one litre of water. Mix well and wet all foliage thoroughly.
3. For low pressure handwand application of Optica Trio, use 1.5 - 2.5 litres of Optica Trio per hectare depending on broadleaf weeds that are present. Use 50-200 L/ha of water.
4. For broadcast treatment of large areas use 1.5 - 2.5 litres of Optica Trio per hectare depending on broadleaf weeds that are present. Use 50-200 L/ha of water.

5. For broadcast treatment of small areas use 15 – 25 mL of Optica Trio mixed in 2 L of water per 100 m², depending on broadleaf weeds that are present.

Optica Trio only controls broadleaved weeds. Optica Trio will not control annual or perennial grasses.

Personal Protective Equipment (PPE) for Non-Crop Lands and Industrial Sites

When applying with ground boom equipment:

- When handling 208 L or less of Optica Trio:
Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. During mixing and loading, protective eyewear must also be worn.
- When handling more than 208 L of Optica Trio:
Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes when mixing, loading, application, clean-up and repair. During mixing and loading, protective eyewear must also be worn. Application must be made using ground boom equipment with an enclosed cab. Gloves are not required during application when applicator is in an enclosed tractor.

When applying with right-of-way sprayers:

- When handling 41 L or less of Optica Trio:
Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. During mixing and loading, protective eyewear must be worn, and a closed mixing/loading system must be used.
- When handling more than 41 L of Optica Trio:
Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. During mixing and loading, protective eyewear must be worn, and a closed mixing/loading system must be used. Application must be made using ground boom equipment with an enclosed cab. Gloves are not required during application when applicator is in an enclosed tractor.

When applying with a low pressure handwand:

- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. During mixing and loading, protective eyewear must be worn.

When applying with backpack equipment:

- Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. During mixing and loading, protective eyewear must be worn.

PRODUCT SPECIFIC PRECAUTIONS:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-265-4624 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

BROADLEAF WEEDS, WILD OATS, GREEN AND YELLOW FOXTAIL CONTROL IN SPRING WHEAT ONLY IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION OF BRITISH COLUMBIA

To control broadleaf weeds, wild oats, green and yellow foxtail in spring wheat only in the following tank mix:

<p>OPTICA TRIO HERBICIDE</p> <p>1.5 or 2.5 L/ha</p> <p>+</p> <p>230 mL/ha of HORIZON[®] 240 EC OR Signal[™] Herbicide</p> <p>+</p> <p>0.8% volume/volume* of SCORE[®] Or Signal[®] Adjuvant</p> <p>OR</p> <p>230 mL/ha of Signal Herbicide OR Nufarm Clodinafop Herbicide</p> <p>+</p> <p>0.25% volume/volume** of Nufarm Enhance[®]</p>	<p>Weeds Controlled</p> <p>Green Foxtail Wild Oats Yellow Foxtail</p> <p>And Broadleaf Weeds listed on the OPTICA TRIO HERBICIDE label</p>
---	--

***800 mL** of SCORE[®] or Signal Adjuvant with 100 L of water per hectare

****250 mL** of Nufarm Enhance with 100 L of water per hectare

Note:

1. DO NOT use OPTICA TRIO HERBICIDE/HORIZON or SIGNAL Herbicide or Nufarm Clodinafop Herbicide tank mix on barley or on oats.
2. When tank mixing, read and observe all directions and restrictions and follow the most stringent precautions of the tank mix partners labels.
3. Apply when spring wheat is in the 2-5 leaf stage.
4. DO NOT APPLY THIS TANK MIX BY AIR.
5. Use a minimum of 100 L of water per hectare.
6. Observe a minimum interval to harvest of 60 days after treatment

7. Observe a minimum of 7 days before grazing livestock on crops treated with the tank mix.
8. Apply when broadleaf weeds are in the 2-3 leaf stage, when wild oats are in the 1-6 leaf stage and when foxtails are in the 1-5 leaf stage.
9. Do not contaminate streams, rivers, lakes, ponds, wetlands, sloughs, irrigation waters, vegetated ditch banks and other water used by livestock or for domestic purposes. Do not apply within 15 meters of a wetland area or any of the aforementioned bodies of water.
10. Do not apply more than one treatment per year.

OPTICA TRIO HERBICIDE WITH EVEREST® TANKMIX:

CROP	WEEDS	RATE	REMARKS
Spring Wheat Including: Hard red spring, Canada Prairie spring, soft white spring and extra strong (utility) wheat	Wild Oats* and Green Foxtail (Wild Millet) Broad Leaf Weeds listed on the OPTICA TRIO HERBICIDE label. *Reduction in wild oat control may be observed with this tank mixture at high populations	EVEREST SOLUPAK 70 DF, 43 g/ha + OPTICA TRIO HERBICIDE 1.5 or 2.5 L/ha. + Non-ionic surfactant such as: LI 700, Super Spreader, Agral 90, Ag-Surf, or Surf 92 at 0.25% v/v or 0.25L/100L.	Follow all directions and precautions on the tank-mix partner labels.
	Green Foxtail ONLY (Wild Millet) Broadleaf Weeds listed on the OPTICA TRIO HERBICIDE label.	EVEREST SOLUPAK 70 DF, 21.5 g/ha + OPTICA TRIO HERBICIDE 1.5 or 2.5 L/ha. + Non-ionic surfactant such as: LI 700, Super Spreader, Agral 90, Ag-Surf, or Surf 92 at 0.25% v/v or 0.25L/100L.	

MIXING AND SPRAYING INSTRUCTIONS:

Spray Pressure: 200-300 kPa

Spray Volume: minimum of 50 litres of water per hectare

Use mesh screen and filters

1. Fill the sprayer ½ full with water. Start gentle agitation.
2. Add tank mix partner, if applicable, and agitate thoroughly.
3. Fill tank ¾ full with water and add OPTICA TRIO HERBICIDE.
4. Complete filling of sprayer and maintain gentle agitation while mixing and spraying.
5. Do not allow contents to stand without agitation.
6. Do not apply more than one treatment per year.

®All product names are trademarks of their respective owners.

FRONT

GROUP 4 HERBICIDE

OPTICA™ TRIO

Solution

COMMERCIAL

**BROADLEAF HERBICIDE FOR USE IN WHEAT (Spring, durum and winter),
BARLEY, OATS AND NON-CROP LAND/INDUSTRIAL SITES**



CAUTION

POISON

DANGER - CORROSIVE TO EYES

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING.
PROTECT FROM FREEZING.**

Warning, contains the allergen, soy

ACTIVE INGREDIENT:

MCPA, present as dimethylamine salt . . . 160 g/L
Dichlorprop-P, present as dimethylamine salt . . . 310 g/L
Mecoprop-P, present as dimethylamine salt . . . 130 g/L

REGISTRATION No.: 29662 PEST CONTROL PRODUCTS ACT

Nufarm UK Limited
Wyke Lane, Wyke, Bradford
West Yorkshire, BD12 9EJ
England

Canadian Agent:
Nufarm Agriculture Inc.
5101, 333 - 96th Ave NE
Calgary, Alberta T3K 0S3

For product use information, please contact:
United Agri Products Canada, Inc.
789 Donnybrook Drive
Dorchester, ON
N0L 1G5
1-800-265-4624

24 Hour Emergency Response Number 1-800-424-9300

NET CONTENTS 10 L

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Store with container tightly closed, and away from seeds, fertilizers, plant food-stuffs and greenhouses. Protect from freezing. Avoid spray drift on to susceptible crops. Wash sprayer thoroughly, before and after use. After use wash hands and skin contaminated by spray or concentrate.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields and public buildings.

If this pest control product is to be used on a commodity that may be exported to the U. S. and you require information on acceptable residue levels in the U. S., visit CropLife Canada's website at: www.croplife.ca.

PROTECTIVE CLOTHING AND EQUIPMENT (PPE):

The following PPE is for all agricultural uses. For PPE required when treating non-crop land/industrial sites, use the PPE listed in the "Non-Crop Lands and Industrial Sites" section.

Handling the concentrate (mixing and loading): Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (face shield or safety goggles). Rinse gloves before removal. Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5L.

Handling the dilute spray solution (during application or repairing or cleaning equipment): Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor.

If OPTICA TRIO is applied when crop is under stress, shortening of straw may occur, but yield will not be affected. Protect from freezing.

Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 12 hours after application for agricultural uses.

Do not enter or allow entry into treated non-crop land/industrial sites until residues have dried.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

ENVIRONMENTAL HAZARDS:

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient.

High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low-blood pressure and unconsciousness. Treat symptomatically.

EMERGENCY TELEPHONE NUMBERS:

For spills or transportation accidents, Chemtrec, 1-800-424-9300.

For health or environmental emergencies, Prosar, 1-877-325-1840.

For product and use information, UAP Canada Inc. 1-800-265-4624.

STORAGE:

Protect from freezing

DISPOSAL:

For Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site.

Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Returnable-Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

All product names are trademarks of their respective companies.