

GROUP	7	FUNGICIDE
--------------	----------	------------------

Vertisan® Fungicide

EMULSIFIABLE CONCENTRATE

FOR USE ON CANOLA, DRY LEGUMES, CEREALS, CORN, SORGHUM, SOYBEANS,
SUNFLOWER, TUBEROUS AND CORM VEGETABLES, SUGAR BEETS AND TURFGRASS

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Penthiopyrad 200 g/L

REGISTRATION NO. 30332 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1L – 1,500 LITRES

DANGER – EYE IRRITANT
SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

Belchim Crop Protection Canada, Inc.
104 Cooper Dr., Unit 3
Guelph, ON
N1C 0A4
1-866-613-3336

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Severely irritating to the eye. **DO NOT** get in eyes.
- May irritate the skin. Avoid contact with skin.
- Potential skin sensitizer.
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- For corn detasseling do not enter or allow worker entry into treated areas during the restricted entry interval of 3 days. For all other crops and activities, do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.
- Sod Farms: **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Golf Courses: **DO NOT** enter or allow entry into treated areas until sprays have dried.
- See the Aerial Application Label Instructions section of this label for additional precautions specific to aerial application.
- Apply only when the potential for drift to areas of human habitation and human activity (other than golf courses) such as parks, school grounds, and playing fields, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

ENVIRONMENTAL PRECAUTIONS:

- Toxic to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination, store this product away from food or feed.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

In case of emergency involving a poisoning or major spill call 1-866-336-2983.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

<<DFU Booklet>>

GROUP	7	FUNGICIDE
--------------	----------	------------------

Vertisan® Fungicide

EMULSIFIABLE CONCENTRATE

FOR USE ON CANOLA, DRY LEGUMES, CEREALS, CORN, SORGHUM, SOYBEANS,
SUNFLOWER, TUBEROUS AND CORM VEGETABLES, SUGAR BEETS AND TURFGRASS

COMMERCIAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

ACTIVE INGREDIENT: Penthiopyrad 200 g/L

REGISTRATION NO. 30332 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1L - 1000 LITRES

DANGER – EYE IRRITANT
SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

Belchim Crop Protection Canada, Inc.
104 Cooper Dr., Unit 3
Guelph, ON
N1C 0A4
1-866-613-3336

GENERAL INFORMATION

Vertisan® Fungicide is a broad-spectrum fungicide, recommended for control or suppression of foliar and soil-borne plant diseases and has curative and locally systemic activity. Vertisan® Fungicide must be applied in a regularly scheduled protective spray program in rotation with other fungicides. See directions below for specific crop/disease recommendations.

Vertisan® Fungicide can be applied with ground or air, except as otherwise directed, using sufficient water to obtain thorough coverage of plants. Minimum aerial application volume is 40 L/ha and minimum ground application volume is 110L/ha.

DIRECTIONS FOR USE

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) fine classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) fine classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Spray Buffer zones:

Use of the following spray methods or equipment DO NOT require a spray buffer zone: hand-held or backpack sprayer, spot treatment, inter-row hooded sprayer, soil drench and soil incorporation.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands). and estuarine/marine habitats.

Method of application	Crop		Spray Buffer Zones (meters) Required for the Protection of:	
			Freshwater Habitats of Depths:	
			Less than 1 m	Greater than 1 m
Field Sprayer	Chickpeas, lentils, beans, peas, sunflower seed, sugar beets, rye, sorghum, triticale, wheat, barley, corn (all), oats, soybeans, canola, potatoes		4	2
Aerial	Chickpeas, lentils, beans, peas	Fixed-wing	65	5
		Rotary-wing	55	1
	Sunflower seed, sugar beets	Fixed-wing	50	1
		Rotary-wing	40	1
	Rye, sorghum, triticale, wheat, barley, corn(all) Oats	Fixed-wing	35	1
		Rotary-wing	30	1
Soybeans, canola	Fixed-wing	25	1	
	Rotary-wing	20	1	

	Potatoes	Fixed-wing	55	4
		Rotary-wing	45	1

Method of application	Crop	Spray Buffer Zones (meters) Required for the Protection of:			
		Freshwater Habitats of Depths:		Estuarine/Marine Habitats of Depths:	
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field Sprayer	Turf	10	3	1	1

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

NOTE: Applicators may recalculate a site-specific spray buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) fine classification.' To access the Spray Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

CANOLA

Crop	Disease Controlled	Application Rate	SPECIFIC DIRECTIONS
Canola	Sclerotinia stem rot (white mold – <i>Sclerotinia sclerotiorum</i>)	1.25 to 1.5 L/ha	Begin application at 20-50% bloom prior to disease development to control white mold. Under high disease pressure, make a second application 7-14 days later. Use higher rate and shorter interval when disease pressure is high.

RESTRICTIONS AND PRECAUTIONS: Canola

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum seasonal use rate is 3 L/ha.
- Minimum time (PHI) between application and harvest is 21 days.
- For any of the diseases listed above, use the high rate under heavy pest pressure.

DRY LEGUMES

Crop	Disease Controlled	Application Rate	SPECIFIC DIRECTIONS
Dry Legumes (dry shelled beans and peas) chickpea (garbanzo); lentil; guar; lablab bean; lentil; broad bean (dry); pigeon pea; (<i>Lupinus</i>) grain lupin, sweet lupin, white lupin, white sweet lupin; (<i>Phaseolus</i>) field bean, kidney bean, lima bean, navy bean, pinto bean, tepary bean, (<i>Vigna</i>) adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, (<i>Pisum</i>) field pea	Ascochyta blight (<i>Ascochyta spp.</i>)	1 to 1.5 L/ha	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
	Gray mold (<i>Botrytis cinerea</i>)	1.25 to 1.5 L/ha	
	Suppression of Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	1 to 1.75 L/ha	

RESTRICTIONS AND PRECAUTIONS: Dry Legumes

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum total seasonal use rate is 3 L/ha.
- Minimum time (PHI) between application and harvest is 21 days.
- For any of the diseases listed above, use the high rate under heavy pest pressure.

Cereal Grains

Crop	Disease Controlled	Application Rate	SPECIFIC DIRECTIONS
Small grains:			
Wheat, Barley, triticale	Suppression of Septoria leaf spot (<i>Septoria tritici</i>)	1.2 to 1.75 L/ha	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high. To optimize yields in cereals, it is important to protect the flag leaf from foliar diseases. For optimizing yield and flag leaf disease control, apply Vertisan® Fungicide at Feeke's 9, 'flag leaf out'.
Wheat, oats, triticale, barley	Black stem rust (<i>Puccinia graminis f. sp. tritici</i>)		
Wheat, rye, triticale	Brown leaf rust, orange leaf rust (<i>Puccinia recondita f. sp. tritici</i>)		

RESTRICTIONS AND PRECAUTIONS: Cereal grains

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum seasonal use rate is 3.5 L/ha.
- Crop may be used for grazing, forage or hay within 0 days after the last application. Do not apply after flowering (Feekes 10.5.1).
- For any of the diseases listed above, use the high rate under heavy pest pressure.

Corn and sorghum

Crop	Disease Controlled	Application Rate	SPECIFIC DIRECTIONS
Corn, sorghum:			
corn, field corn, sweet corn, seed popcorn, sorghum (milo), sorghum (sudangrass and hybrids)	Suppression of Grey leaf spot (<i>Cercospora zeae-</i> <i>maydis</i>) Common rust (<i>Puccinia sorghi</i>) – on corn; field, sweet, seed and popcorn only	1 to 1.75 L/ha	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high. For suppression of grey leaf spot, add a non-ionic surfactant (NIS) at a rate of 0.125% v/v (1.25 L per 1,000 L of spray solution).

RESTRICTIONS AND PRECAUTIONS: Corn and sorghum

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum seasonal use rate is 3.5 L/ha.
- Minimum time (PHI) between application and grain and stover harvest is 7 days for corn. Minimum time (PHI) between application and grain and stover harvest is 30 days for sorghum.
- Corn and sorghum may be used for grazing or forage within 0 days after the last application.
- For any of the diseases listed above, use the high rate under heavy pest pressure.

SOYBEANS

Crop	Disease Suppressed	Application Rate	SPECIFIC DIRECTIONS
Soybean	Asian Soybean Rust <i>(Phakospora pachyrhizi)</i> Brown Spot <i>(Septoria glycines)</i> Frogeye leafspot <i>(Cercospora sojina)</i>	1 to 1.75 L/ha	Begin applications prior to disease development and continue on a 7 to 14-day interval. Use higher rate and shorter interval when disease pressure is high.

RESTRICTIONS AND PRECAUTIONS: Soybean

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum total seasonal use rate is 3 L/ha.
- Soybeans may be used for forage and hay 0 days after application.
- Minimum time (PHI) between application and harvest is 14 days.
- For any of the diseases listed above, use the high rate under heavy pest pressure.

SUNFLOWER

Crop	Disease Suppressed	Application Rate	SPECIFIC DIRECTIONS
Sunflower	Rust <i>(Puccinia helianthi)</i> Sclerotinia head rot <i>(Sclerotinia sclerotiorum)</i>	1.75 L/ha	Begin applications prior to disease development and continue on a 7 to 14-day interval. Use higher rate and shorter interval when disease pressure is high.

RESTRICTIONS AND PRECAUTIONS: Sunflower

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum total seasonal use rate is 4.5 L/ha.
- Minimum time (PHI) between application and harvest is 14 days.
- For any of the diseases listed above, use the high rate under heavy pest pressure.

TUBEROUS AND CORM VEGETABLES

Crop	Disease Controlled	Application Rate	SPECIFIC DIRECTIONS
Tuberous and Corm Vegetables			
Potato; sweet potato	Suppression of early blight (<i>Alternaria solani</i>)	1 to 1.75 L/ha	Begin applications prior to disease development and continue on a 7 to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
Potato; Artichoke, Chinese; Artichoke, Jerusalem; Canna, edible; Chufa, Sweet potato	Gray mold (<i>Botrytis cinerea</i>)	1.25 to 1.5 L/ha	
Potato, sweet potato	Suppression of stem rot and transplant rot (sweet potato) (<i>Rhizoctonia solani</i>)	15.5 to 31 mL per 100 m row	Apply in furrow at planting, using 1.4-1.75 L water/100 row meters. Do not exceed 1.75 L/ha.

RESTRICTIONS AND PRECAUTIONS: Tuberous and corm vegetables

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum total seasonal use rate is 5 L/ha.
- Minimum time (PHI) between application and harvest is 7 days.
- For any of the diseases listed above, use the high rate under heavy pest pressure.

SUGARBEET

Crop	Disease Controlled	Application Rate	SPECIFIC DIRECTIONS
Sugarbeet	Suppression of Powdery mildew (<i>Erysiphe betae</i>)	1 to 1.5 L/ha	Begin applications prior to disease development and continue on a 7 to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
	Crown and root rots (<i>Rhizoctonia solani</i>)	15.5 to 31 mL per 100 m row	Apply in furrow at planting, using 1.4-1.75 L water/100 row meters. Do not exceed 2.2 L/ha per application.

RESTRICTIONS AND PRECAUTIONS: Sugarbeet

- Make no more than 2 sequential applications of Vertisan® Fungicide before switching to a fungicide with a different mode of action.
- Maximum seasonal use rate is 4.5 L/ha.
- Sugarbeet tops may be used for grazing or forage 7 days after the last application.
- Minimum time (PHI) between application and harvest is 7 days.
- For any of the diseases listed above, use the high rate under heavy pest pressure.

CROP ROTATION

Crops and crop groups that are on this label and the following list of crops and crop groups may be planted immediately after harvest:

alfalfa, low-growing berries crop subgroup (strawberries), *brassica* (cole) leafy vegetables crop group, bulb vegetables crop group (onion), cucurbit vegetables crop group (cucumber, melons, squash), fruiting vegetables crop group (tomato, pepper), leafy vegetables crop group (lettuce, celery, spinach), legume vegetables crop group (succulent), peanuts, root vegetables crop subgroup (carrot, radish, turnip)

All other crops cannot be planted until 12 months after the last application of Vertisan® Fungicide.

TURFGRASS

Vertisan® Fungicide may only be used on turfgrasses that are being grown for golf courses (tee box areas, roughs, fairways, greens, collars etc.) and commercial sod farms.

Vertisan® Fungicide is a broad spectrum fungicide recommended for the control of many important diseases of turfgrass. For maximum performance Vertisan Fungicide should be applied preventatively but also may be used curatively for some diseases.

USE Vertisan Fungicide IN A PREVENTATIVE DISEASE CONTROL PROGRAM

Apply in sufficient water to ensure thorough coverage.

Apply the recommended rates as indicated in the directions for use in spray volumes up to 40 L/100 m² or 4,000 L/ha. Apply with a properly calibrated sprayer.

Apply after mowing OR allow sprayed area to completely dry before mowing.

For control of foliar diseases, allow sprayed area to completely dry before irrigation.

Under conditions optimum for high disease pressure, use the higher rate.

For optimum turf quality and disease control, use Vertisan Fungicide in conjunction with turf management practices that promote good plant health and optimum disease control.

Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

For broader spectrum control, Vertisan Fungicide can be tank mixed with other fungicides registered for this use. Check compatibility before tank mixing. When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

Turf	
Disease Controlled	Dollar Spot (<i>Sclerotinia homeocarpa</i> syn. <i>Clariireedia homeocarpa</i>)
Location	Golf course tees and greens, fairways and all other turfgrass
Use rate/100m ²	22.5-37.5 mL
Use rate/ha	2.25-3.75 L
Application Interval (days)	14 days for golf course tees and greens 14-21 days for golf course fairways and all other turfgrass.
Comments	Begin application prior to or at the early stages of disease

	development. Use the high rate for moderate to high disease pressure and/or for curative situations
Disease Controlled	Brown Patch (<i>Rhizoctonia solani</i>)
Location	Golf course tees and greens, fairways and all other turfgrass
Use rate/100m ²	22.5-37.5 mL
Use rate/ha	2.25-3.75 L
Application Interval (days)	14
Comments	Begin application at the early stages of disease development. Use high rate for moderate to high disease pressure or for curative situations
Disease Controlled	Anthrachnose foliar blight (<i>Colletotrichum cereale</i>)
Location	Golf course tees and greens, fairways and all other turfgrass
Use rate/100m ²	22.5-37.5 mL
Use rate/ha	2.25-3.75 L
Application Interval (days)	14
Comments	Begin applications at the early stages of disease development. Vertisan Fungicide may be applied in a tank mix with Daconil 2787 or Heritage fungicide, which are registered for anthracnose control on turf. Use high rate of Vertisan Fungicide for high disease pressure
Disease Controlled	Grey snow mould (<i>Typhula incarnata</i> , <i>Typhula ishikariensis</i>)
Location	Golf course tees and greens, fairways and sod farms
Use rate/100m ²	53.5 mL
Use rate/ha	5.35 L
Application Interval (days)	Apply once per season.
Comments	Apply in late fall before snow cover. Do not apply on top of snow.

USE RESTRICTIONS FOR TURFGRASS

- DO NOT apply this product through any type of irrigation system.
- DO NOT apply more than 170 mL of Vertisan Fungicide per 100m² per year or 17 L of Vertisan Fungicide per hectare per year.
- Keep people and pets away from areas treated with Vertisan Fungicide until treated area has dried
- DO NOT apply using aerial application equipment

APPLICATION INFORMATION

Mixing Instructions

1. Fill clean spray tank 1/4 - 1/2 full of water.
2. While agitating, add the required amount of Vertisan® Fungicide, continuing agitation until the product is completely dispersed.
3. Continue filling the tank, with agitation, adding desired additives

Mix thoroughly to fully disperse the fungicide; once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Belchim Crop Protection Canada Inc. at 1-866-613-3336 or www.belchimcanada.com for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses

that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Belchim Crop Protection Canada Inc.

Adjuvants

Vertisan® Fungicide may be used with nonionic surfactants at typical agricultural use rates for these adjuvants.

SPRAYER CLEANUP:

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

AERIAL APPLICATION INSTRUCTIONS:

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions for Aerial Application

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the **“National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides”**.

Operator Precautions for Aerial Application

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions for Aerial Application

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-866-613-3336 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Apply the recommended rate in a minimum spray volume of 40 litres per hectare.

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Severely irritating to the eye. **DO NOT** get in eyes.
- May irritate the skin. Avoid contact with skin.
- Potential skin sensitizer.
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- For corn detasseling do not enter or allow worker entry into treated areas during the restricted entry interval of 3 days. For all other crops and activities, do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.
- Sod Farms: **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Golf Courses: **DO NOT** enter or allow entry into treated areas until sprays have dried.
- See the Aerial Application Label Instructions section of this label for additional precautions specific to aerial application.
- Apply only when the potential for drift to areas of human habitation and human activity (other than golf courses) such as parks, school grounds, and playing fields, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

ENVIRONMENTAL PRECAUTIONS:

- Toxic to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that Vertisan® Fungicide contains a Group 7 Fungicide. Any fungal population may contain individuals naturally resistant to Vertisan® Fungicide and other group 7 Fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- ◆ Where possible, rotate the use of Vertisan® Fungicide or other Group 7 fungicides with different groups that control the same pathogens. Use tank mixtures with fungicides from different groups when such use is permitted.
- ◆ Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- ◆ Where possible, make use of predictive disease models to effectively time fungicide applications.
- ◆ Monitor treated fungal populations for resistance development.
- ◆ If disease continues to progress after treatment with this product, do not increase the use rate, Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- ◆ For further information and to report suspected resistance, contact Belchim Crop Protection Canada Inc. representatives at 1-866-613-3336.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination, store this product away from food or feed.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

In case of emergency involving a poisoning or major spill call 1-866-336-2983.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

"®/™ Trademarks of E.I. du Pont de Nemours and Company, used under license

*All other products mentioned are trademarks of their respective companies.